DOI: 10.2225/vol12-issue1-fulltext-7

RAPD analysis of seized marijuana (Cannabis sativa L.) in Turkey

Emine Pinarkara

Department of Animal Sciences Biometry-Genetics Unit Faculty of Agriculture Selcuk University, 42075 Campus, Konya, Turkey Tel: 903322232811 Fax: 903322410108 E-mail:eminepinarkara@hotmail.com

Seyit A. Kayis*

Department of Animal Sciences Biometry-Genetics Unit Faculty of Agriculture Selcuk University, 42075 Campus, Konya, Turkey Tel: 903322232830 Fax: 903322410108 E-mail: skayis@selcuk.edu.tr

Erdogan E. Hakki

Department of Field Crops Faculty of Agriculture Selcuk University, 42075 Campus, Konya, Turkey Tel: 903322232863 Fax: 903322410108 E-mail: eehakki@selcuk.edu.tr

Ayla Sag

Institute of Forensic Sciences Istanbul University Istanbul, Turkey Tel: 902124143000 Fax: 902125880011 E-mail: aylasag@yahoo.com

Financial support: The Scientific And Technical Research Council of Turkey (TUBITAK-Kariyer-1040547) and the Scientific Research Coordination Center of Selcuk University (BAP-07201006) are gratefully acknowledged for their supports.

Keywords: AMOVA, cluster analysis, individualization of seized marijuana, principle coordinate analysis, randomly amplified polymorphic DNA, UPGMA.

Abbreviations:AFLP: amplified fragment length polymorphisms
AMOVA: analysis of molecular variance
CBD Cannabidiol
PCoA: principal coordinate analysis
PCR: polymerase chain reaction
RAPD: randomly amplified polymorphic DNA
RAPD: randomly amplified polymorphic DNA
STR: short tandem repeat
THC: Δ^9 -tetrahydrocannabinol
UPGMA: unweighted pair-group method using arithmetic average

Cannabis sativa L. is a multiple-use plant. However, its cultivation is strictly controlled due to its psychoactive nature and usage in producing drugs such as marijuana, and hashish. In this study, psychoactive type *Cannabis* samples, which were seized from 29 different

locations of Turkey, were used. Interests were to identify the genetic relatedness of the seized samples and to partition molecular variance between and within populations. Randomly Amplified Polymorphic DNAs were employed for analysis based on single plant

^{*}Corresponding author

material and bulked samples of them. Data were analysed via cluster and principal coordinate analyses (PCoA). Analysis of molecular variance (AMOVA) was performed to obtain variations between and within populations. *Cannabis* accessions were basically separated into two main groups by PCoA and cluster analyses according to geographical regions. One of them was made up of Cannabis plants, which were seized from mostly western part of Turkey (group 1). The other one was made up of Cannabis plants that were seized from mostly eastern part of Turkey (group 2). It is found that 20.23% of the genetic variation is due to differences between accessions groups while 79.77% of the genetic variation is due to between accessions within accessions groups. Compared to group 1, group 2 showed more variation.

Cannabis sativa L. is thought to have originated from the Central Asia region and has since been distributed worldwide by humans (Small and Cronquist, 1976). It is a plant that provides food and oil from its seeds, fiber for rope, fabric from its stems, and psychoactive drugs from its flowers and leaves. Hemp seeds oil can also be used for fuel and as raw material for plastics (Ranalli and Venturi, 2004) as well as feed for livestock or as a fertilizer (Karus and Vogt, 2004).

Beside the economical properties, mentioned above, some varieties of *Cannabis* have psychoactive potency as well. *Cannabis* plants that contain low Δ^9 -tetrahydrocannabinol (THC), a low THC:Cannabidiol (CBD) ratio and are cultivated for fiber and/or achenes (*e.g.* seeds) are called hemp. On the other hand, *Cannabis* plants that have high THC content, high THC:CBD ratio and are used for their psychoactive potency are known as marijuana (Alghanim and Almirall, 2003; Elsohly and Slade, 2005; Hillig, 2005). For hemp, EU has assigned the upper levels of THC and CBD to THC ratio as 0.2% and 2%, respectively.

In many countries, including Turkey, possession and cultivation of *Cannabis* was either ceased or limited because of its potential use as a drug. When samples of suspect materials are recovered, they must be tested for the presence of controlled substances (*e.g.* marijuana). In addition to the identification of marijuana samples, it is desirable but difficult to link individual growers and distributors to specific illicit field and greenhouse operations. Molecular genetics may offer solution in identification and individualisation via investigating the genetic relatedness between individuals/populations.

Jagadish et al. (1996) were able to distinguish between the samples from distinct sources in a randomly amplified polymorphic DNA (RAPD) assay conducted with 51 *C. sativa* samples. Genetic analysis using in combinations of RAPD and restriction fragment length polymorphism (RFLP) methods were also found to be useful in distinguishing between drug type, fiber type and intermediate drug type strains (Shirota et al. 1998). Hakki

et al. (2003) used RAPD and amplified fragment length polymorphisms (AFLP) markers to fingerprint the 18 different *Cannabis* individuals from five different locations representing 3 geographical regions of Turkey. In another study, it was reported that it was possible to discriminate illegal, potent marijuana cultivars from hemp plants by using AFLP markers (Datwyler and Weiblen, 2006). In a preliminary work conducted with three strains of *C. sativa* from different sources, Kojoma et al. (2002) reported that different samples were identified by means of inter simple sequence repeats (ISSR). In a recent study by using ISSRs marijuana (*Cannabis sativa* L.) was separated efficiently from hemp (Hakki et al. 2007).

Gillan et al. (1995) reported the differentiation of C. sativa samples with the use of RAPDs when HPLC analysis was inefficient. Faeti et al. (1996) assessed genetic diversity of C. sativa cultivars/accessions (from 5 European countries, and one accession from Korea) by using RAPD markers and high levels of polymorphism were reported. In a study of genetic structure and degree of variability of six C. sativa L. varieties via RAPD markers, it was reported that 5 varieties were properly identified with the scored loci (Forapani et al. 2001). Hsieh et al. (2003) investigated the usage of short tandem repeat (STR) loci in identification of Cannabis samples and predicting their genetic relationship. Alghanim and Almirall (2003) were developed STR markers for Cannabis. They reported that STR markers were very effective in uniquely identifying 27 profiles of the Cannabis samples tested and useful for DNA typing and genetic relatedness analyses. Gilmore and Peakall (2003) isolated microsatellite markers in Cannabis sativa L. which have utility for characterizing genetic diversity in cultivated and naturalized Cannabis populations. Gilmore et al. (2003) reported that STR markers are capable of discriminating among individuals and varieties of Cannabis.

RAPD markers were used to individualize Palo Verde tree in a criminal case (Yoon, 1993) and strawberry in a civil case (Congiu et al. 2000). In both cases the method has been accepted in court although, in the Palo Verde tree case the statistical significance was not used since the representative population consists of too few samples. Congiu et al. (2000) employed RAPD markers for individualization of strawberry because of its two main advantages: it allows random sampling of markers over whole genomic DNA and does not require any previous information on the genome of the organism under investigation.

Although RAPD marker analysis has reproducibility problem, it is inexpensive, simple to perform, and has moderate ability to distinguish between unrelated individuals compared to AFLPs and STRs (Coyle et al. 2003). Therefore, the method can still be useful for individualization of Cannabis samples in the developing countries that have very limited lab facilities but majority of *Cannabis* production occur.

Sample ID ^a	Paralel ID ^b	Settlement seized ^c	Province ^d	
C1	3102	Tekirdag	Tekirdag	
C2	492784175	Geyve	Sakarya	
С3	04-62927/5432 ND	Tekirdag	Tekirdag	
C4	05/002598/315	Edirne	Edirne	
C5	04/409/5782	Susehri	Sivas	
C6	058576/5057	Tekirdag	Tekirdag	
C7	056012/4833	Kocaeli	Kocaeli	
C8	065364/5645	Golcuk	Istanbul	
C9	4243-2	Ferizli	Sakarya	
C10	4243	Ferizli	Sakarya	
C11	2075/1	Salihli	Manisa	
C12	758/9	Izmir	Izmir	
C13	847/1-C-1	Denizli	Denizli	
C14	677/2	Didim	Aydin	
C15	315/2	Aydin	Aydin	
C16	676/2 A	Didim	Aydin	
C17	04 4047	Osmaniye	Osmaniye	
C18	AT 05/1458	Kadirli	Osmaniye	
C19	AT 05/678	Gaziantep	Gaziantep	
C20	AT 04/4114	Dortyol	Hatay	
C21	AT 04/3933	Gaziantep	Gaziantep	
C22		Bingol	Bingol	
C23		Elazig	Elazig	
C24		Malatya	Malatya	
C25		Rize	Rize	
C26		Ardesen	Rize	
C27		Akcaabat	Trabzon	
C28		Trabzon	Trabzon	
C29		Arsin	Trabzon	

Table 1. Cannabis accessions used in the study and their origin.

^a Sample name used in this study, ^bFile code in seized samples (if available), ^cThe region where sample was seized, ^dProvincial location where sample was seized.

Pinarkara, E. et al.

The objectives of the present study were: 1) to analyze the high number of seized *Cannabis* samples by means of RAPD, 2) to compare two different approaches (in the first, a single plant represents an accession and in the second, a set of ten different plants of the same accession bulked equally represent the specific accession) for individualizing *Cannabis* accessions, 3) to obtain information on the genetic variation and relatedness which might be a useful information about the sources and distribution networks of these illicit substances.

MATERIALS AND METHODS

Plant material

Psychoactive *Cannabis* samples used in this study were seized from 29 different locations (C1-C29) representing geographically distinct and problematic areas of Turkey (western and eastern parts). Some of the materials, used in this study, were previously also utilized for discriminating drug type *Cannabis* from hemp types (Hakki et al. 2007). All the information relevant to the seized samples and accessions are shown in Table 1. Provincial locations of the seized *Cannabis* accessions are shown on a map of Turkey in Figure 1. Ten seeds were planted from each accession to produce material for DNA extraction. Plants were grown in a fully automated greenhouse.

DNA extraction from leaf

Leaves collected from three week old seedlings were shock-frozen in liquid nitrogen and stored at -80°C until DNA isolations were performed. DNAs were extracted individually from a total of 290 samples. Total DNAs of the samples were extracted using a standard 2X CTAB protocol with minor modifications (Rogers and Bendich, 1988). For each accession, 100 mg of leaf sample from 10 different plants were used and DNAs were isolated individually. After concentrations were determined by an Eppendorf BioPhotometer, sample DNAs were diluted to the working concentration of 20 ng/ μ L.

PCR amplification of the DNA with RAPD primers

In this study, 22 arbitrary RAPD primers that gave the most informative patterns (in terms of repeatability, scorability and the ability to distinguish between individuals) were selected for identification (Table 2). Each reaction contained 2.5 mM MgCl₂; 10 mM Tris-HCl (pH 8.8); 50 mM KCl; 0.8% Nonidet P40; 200 mM of each of the dNTPs; 0.5 μ M primer; 20 ng DNA template and 0.3 units of Taq DNA Polymerase (Bioron) in a final reaction volume of 25 μ l. After a pre-denaturation step of 3 min at 94°C, amplification reactions were optimized for every individual primer and optimization was usually started by cycling the reaction 45 times at 94°C for 1 min, at annealing temperature (Table 2) for 50 sec and 72°C for 1 min in Eppendorf Mastercycler gradient thermocycler. A final extension was allowed for 10 min at 72°C. Upon completion of the reaction, amplified products were loaded onto a 2.0% agarose/1x Tris-Borate EDTA gel and electrophoresed at 4 V/cm.

In the RAPD assay, two sets of PCR amplifications were carried out. In the first set (SET1), each accession was represented by DNA of randomly selected an individual plant in that accession. In the second set (SET2), pooled DNA from all ten individuals in each accession was used (the one individual used in the first set was also included).

Every primer was amplified for the two sets simultaneously. Then the resultant products were run in agarose gel and evaluated. Amplifications were repeated at least twice (in different time periods) for each primer, using the same reagents and procedure.

Data collection and statistical analysis

Each DNA fragment generated was treated as a separate character and scored as a discrete variable, using 1 to indicate presence, and 0 for absence. Accordingly, a rectangular binary data matrix was obtained and statistical analysis was performed using the NTSYS-pc version 2.1 (Rohlf, 2000) statistical package. A pairwise similarity matrix was generated using simple matching coefficient (by means of SIMQUAL procedure of NTSYS-pc) and principal coordinate analysis (PCoA) was performed using a batch mode of NTSYS-pc both for SET1 and SET2. Then, cluster analysis was performed (by means of SAHN procedure of NTSYS-pc) via unweighted pair-group method using arithmetic average (UPGMA) to develop a dendrogram both for SET1 and SET2. To estimate the strength of the grouping, generated by cluster analysis, bootstrap analysis was performed with 2000 replications using the winboot computer program (Yap and Nelson, 1996). Also, a matrix comparison of Mantel Z test (Mantel, 1967), for the correspondence of the similarity matrices of SET1 and SET2, was performed (by means of MXCOMP procedure of NTSYS-pc) for the null hypothesis that there is no association between the similarity matrices of SET1 and SET2. To obtain significance level, 5000 permutations were performed. In addition, a genetic similarity matrix was calculated according to Nei and Li, (1979).

An analysis of molecular variance (AMOVA) was performed using GENALEX 6 (Peakall and Smouse, 2006) in SET2 to partition the total molecular variance between and within populations (clusters). Significance level was detected via permutation test (n = 1000). *Cannabis* clusters were defined according to the PCoA results (Figure 2). *Cannabis* accessions that do not cluster closely with any of the clusters were removed from data set before AMOVA.

RESULTS

RAPD amplification and analysis

The RAPD markers, used in the analysis of SET1 and SET2 allowed reproducible and informative

						Single plant analysis		ysis	Bulk analysis			
Primer	Primer sequence	Tm (°C)	bp	GC (%)	Annealing temp (°C)	Number of scored bands	Number of polymorphic bands	Percentage of polymorphic bands (%)	Number of Scored bands	Number of polymorphi c bands	Percentage of polymorphic bands (%)	
RAPD L2	5'- GTT TCG CTC C -3'	32	10	60	34	19	18	94.7	13	12	92.3	
RAPD L3	5'- GTA GAC CCG T -3'	32	10	60	33	6	4	66.6	8	6	75.0	
RAPD L4	5'- AAG AGC CCG T -3'	32	10	60	33	14	12	85.7	11	8	72.2	
RAPD L5	5'- AAC GCG CCG T -3'	32	10	60	34	13	9	69.2	15	12	80.0	
RAPD L6	5'- CCC GTC AGC A -3'	34	10	70	34	10	9	90.0	8	6	75.0	
RAPD B1	5'- CCC GCC GTT G -3'	36	10	80	35	32	32	100	32	32	100	
RAPD B2	5'- TGC GCC CTT C -3'	34	10	70	33	14	14	100	8	8	100	
RAPD B3	5'- GAT GAC CGC C -3'	34	10	70	34	10	9	90	8	8	100	
RAPD B4	5'- CTC ACC GTC C -3'	34	10	70	33	11	11	100	13	13	100	
RAPD B5	5'- GAC GGA TCA G -3'	32	10	60	31	20	20	100	18	17	94.4	
RAPD B6	5'- CCG ATA TCC C -3'	32	10	60	31	10	8	80.0	10	7	70.0	
RAPD B7	5'- TTG GTA CCC C -3'	32	10	60	31	10	9	90.0	7	6	85.7	
RAPD B8	5'- ACG GTA CCA G -3'	32	10	60	31	9	9	100	9	9	100	
RAPD B9	5'- CCA GCG TAT T -3'	30	10	50	29	8	8	100	11	11	100	
RAPD B10	5'- CTA CTG CGC T -3'	32	10	60	31	10	10	100	10	10	100	
RAPD B11	5'- CCT CTG ACT G -3'	32	10	60	31	6	6	100	7	7	100	
RAPD B12.2	5'- TCC GAT GCT G -3'	32	10	60	31	7	7	100	4	4	100	
RAPD B13	5'- TTC AGG GTG G -3'	32	10	60	31	6	6	100	8	8	100	
RAPD B14	5'- TCC TGG TCC C -3'	34	10	70	33	12	12	100	8	8	100	
RAPD B16	5'- AGT CGG GTG G -3'	34	10	70	33	11	9	81.8	11	9	81.8	
RAPD B17	5'- GTC GTT CCT G -3'	32	10	60	31	17	17	100	13	13	100	
RAPD B18	5'- GAG TCA GCA G -3'	32	10	60	31	9	9	100	9	9	100	
Total and averaged values				264	248	% 93	241	223	% 92			

Table 2. RAPD primers used in the study and the number and the type of fragments they amplified

Pinarkara, E. et al.

polymorphisms (pictures of gels that were run at different time were given in Figure 4 for illustration purpose). Selected primers yielded a total of 264 bands in SET1, and 241 bands in SET2, 93% and 92% of which were polymorphic, respectively. However, using bulked samples resulted in much more consistent and reliable amplifications. The RAPD B1 primer was the one that gave the highest number of polymorphisms in SET1 and SET2. In the accession C20, that is the accession seized from Hatay, 119 markers were missing. This was due to lack of material to produce DNA from that accession.

Results from statistical analysis of genotypic data

Results from PCoA of RAPD markers by using SET2 showed that *Cannabis* accessions are basically separated into two main groups by PCo axis 1 (Figure 2). In fact, this separation was in agreement with the geographical regions of Turkey. The first group (group 1) was made up of *Cannabis* plants, which were seized from mostly western part (costal regions, namely Mediterranean, Aegean, and Marmara) of Turkey and there were 18 accessions. The second one (group 2) was made up of *Cannabis* plants that were seized from mostly eastern part of Turkey and there were 8 accessions. Accessions C21 (Gaziantep21), C7 (Kocaeli7), and C20 (Hatay20) were not attributable to any group and we named them as outliers. Compared to group 1, group 2 showed more variation.

The genetic relationships among *Cannabis* accessions, in SET2, were presented in a dendrogram (Figure 3). Results of analysis of SET2 via cluster analysis and PCoA were slightly different. Group 1 *Cannabis* accessions in PCoA, were clearly grouped as one major branch with a similarity of 81% based on simple matching similarity index. However, part of *Cannabis* accessions in group 2 (Elazig23, Malatya24, Trabzon28, and Rize25) were attached to the cluster of group 1, and then the rest of group 2 were attached. Accession Gaziantep21, which was seen to be an outlier in the PCoA, took place in this group. Overall similarity, based on simple matching similarity index, was 71%.

Unlike SET2, there was no separation in the seized *Cannabis* accessions in SET1 via either PCoA or cluster analysis. Therefore, no result from PCoA or cluster analysis is presented in here from SET1.

Results from matrix comparison via Mantel Z test showed that there is a statistically significant association between the similarity matrices of SET1 and SET2 (P < 0.001). However, correlation between SET1 and SET2 was weak (r = 0.39).

Genetic similarity matrices, calculated from SET1 and SET2 according to Nei and Li, (1979) were given in Figures 5 and 6, respectively. Genetic similarity matrix for SET2 shows a similarity range from 0.05 to 0.32. The 0.05 level similarity was seen between the accessions C20 and

C2. We note that the accession C20 is the one which has large number of missing marker genotypes, which most likely caused low similarity. Similarity range between the accession C20 and other accessions changed from 0.05 to 0.11. When the C20 is removed from the data set, similarity range changed from 0.15 to 0.32 with a mean of 0.24. The minimum similarity, 0.15, was between accessions C2 and C22 while the maximum similarity, 0.32, was between accessions C10 and C16.

Genetic similarity matrix for SET1 shows a similarity range from 0.06 to 0.28. The 0.06 level similarities are seen between the accession C20 and accessions C2, C5, C9, and C21. When the C20 is removed from the data set, as in SET2, similarity range changed from 0.15 to 0.28 with a mean of 0.20. The minimum similarities are seen between accessions C4 - C5 and C5 - C7 while the maximum similarity, 0.28, was between accessions C27 and C28. Genetic distance matrices both from SET1 and SET2 show that the accessions tested in this study are divergent at the DNA level.

Results from AMOVA indicated that 20.23% of the genetic variation is attributable to differences among accessions groups while 79.77% of the genetic variation is attributable to between accessions within accessions groups. Sum of squares in group 1 and group 2 were found to be 315.94 ($n_1 = 18$) and 188.50 ($n_2 = 8$), respectively. Detailed results from AMOVA were given in Table 3.

DISCUSSION

This study was conducted on two SETs of *Cannabis* materials, by using RAPD markers, which were analyzed simultaneously. The degrees of polymorphism of the markers found were 93% for SET1 and 92% for SET2. Forapani et al. (2001) reported the degree of polymorphism of RAPD markers for different hemp varieties in a range from 31.1% to 97.1%, which includes our findings.

Clusters were observed between the seized *Cannabis* accessions in SET2 via both cluster analysis and PCoA but not in SET1. A weak correlation, resulted from matrix comparison via the Mantel test between similarity matrices of SET1 and SET2, might be interpreted as a reason for not having similar results from SET1 and SET2. This might be due to the template effect, where amplification of faint fragments failed in bulked samples. Hence, more consistent bands were amplified and the specific accessions were better represented.

In the PCoA of SET2, all accessions, except accessions Gaziantep21, Hatay20, and Kocaeli7, were clearly separated into two main groups. The first group was made up of *Cannabis* plants, which were seized from mostly western parts of Turkey, while the second group was made up of *Cannabis* plants that were seized from mostly eastern parts of Turkey. Visual examination of distribution of *Cannabis* accessions on the first two principal coordinates

Table 3. Analysis of molecular	variance (AMOVA).	. Statistics includ	es degrees	of freedom	(df), s	sum of squares,	estimated	variance
(EV), and percentage of total varia	ation (%).							

Source	df	Sum of Squares	EV	%
Among groups	1	80.066	5.33	20.23**
Within groups	24	504.44	21.02	79.77**
Total	25	584.50	26.35	100.00

**Significant at P < 0.01 level.

(Figure 2) indicates that there is more variation in group 2 compared to group 1. Averaged Nei and Li, (1979) genetic similarity, which were found to be 0.263 and 0.220 for group 1 and group 2 respectively, supported visual examination. Findings from AMOVA supported these results by having estimated variances as 18.58 (with 17 degrees of freedom) and 26.92 (with 7 degrees of freedom) for group 1 and group 2, respectively. Results from AMOVA indicated that 20.23% of the genetic variation was attributable to differences among accessions groups while 79.77% of the genetic variation was attributable to between accessions within accessions groups. Datwyler and Weiblen (2006) identified geographic sources of seized drugs by using AFLP markers. In that study, 27.2% of the genetic variation from AMOVA was due to differences between drug and hemp lines, while 20.9% differences among hemp varieties. Our among group variation is comparable with their among hemp varieties variation. In a comparison study of six hemp types Cannabis varieties with RAPD markers, it is reported that the proportion of among-cultivars variance changed dramatically ranging from 12.8% up to 76%. The later was observed between two highly selected, divergent cultivars (Forapani et al. 2001).

Results from UPGMA for group 2 was slightly different compared to PCoA, although accessions of the group 1 were clustered together in UPGMA. PCoA is a translation of similarities/dissimilarities between objects into the actual distances between objects in multidimensional space. On the other hand, in cluster analysis, once a group or cluster is formed from two or more objects, that group can not be broken later in the process. As a result, the dendrogram is not a representation of all pairwise dissimilarities between objects (Legendre and Anderson, 1999).

Examination of UPGMA Dendrogram accession by accession would give information about relatedness in individual level. For example, 0.91 genetic relatedness was found between accessions C3 and C11, and C5 and C15 with a bootstrapping P value of 58.6% and 50.2%, respectively. These results might lead us to speculation that there might be relationship between the individual illicit

Cannabis growers of accessions C3 and C11, and C5 and C15. It is worthy noting that these illicit products are not clonally propagated as they were grown from seeds. Therefore, we are not expecting accessions to have exactly the same genotypes.

ACKNOWLEDGMENTS

This study is derived from MSc thesis of the first author. The authors thank Prof. Sait Gezgin for providing the controlled greenhouse while growing the plants. Authors thank to Istanbul, Izmir, Adana, Malatya, and Trabzon forensic departments for providing research material.

REFERENCES

ALGHANIM, H.J. and ALMIRALL, J.R. Development of microsatellite markers in *Cannabis sativa* for DNA typing and genetic relatedness analyses. *Analytical and Bioanalytical Chemistry*, August 2003, vol. 376, no. 8, p. 1225-1233.

CONGIU, L.; CHICCA, M.; CELLA, R.; ROSSI, R. and BERNACCHIA, G. The use of random amplified polymorphic DNA (RAPD) markers to identify strawberry varieties: a forensic application. *Molecular Ecology*, 2000, vol. 9, no. 2, p. 229-232.

COYLE, H.M.; PALMBACH, T.; JULIANO, N.; LADD, C. and LEE, H.C. An overview of DNA methods for the identification and individualization of marijuana. *Croatian Medical Journal*, 2003, vol. 44, no. 3, p. 315-321.

DATWYLER, S.L. and WEIBLEN, G.D. Genetic variation in hemp and marijuana (*Cannabis sativa* L.) according to amplified fragment length polymorphisms. *Journal of Forensic Sciences*, March 2006, vol. 51, no. 2, p. 371-375.

ELSOHLY, M.A. and SLADE, D. Chemical constituents of marijuana: The complex mixture of natural cannabinoids. *Life Sciences*, December 2005, vol. 78, no. 5, p. 539-548.

FAETI, V.; MANDOLINO, G. and RANALLI, P. Genetic diversity of *Cannabis sativa* germplasm based on RAPD

markers. *Plant Breeding*, November 1996, vol. 115, no. 5, p. 367-370.

FORAPANI, S.; CARBONI, A.; PAOLETTI, C.; MOLITERNI, V.M.C.; RANALLI, P. and MANDOLINO, G. Comparison of hemp varieties using random amplified polymorphic DNA markers. *Crop Sciences*, 2001, vol. 41, no. 6, p. 1682-1689.

GILLAN, R.; COLE, M.D.; LINACRE, A.; THORPE, J.W. and WATSON, N.D. Comparison of *Cannabis sativa* by random amplification of polymorphic DNA (RAPD) and HPLC of cannabinoids: a preliminary study. *Science and Justice*, July-September 1995, vol. 35, no. 3, p. 169-177.

GILMORE, S.; PEAKALL, R. and ROBERTSON, J. Short tandem repeat (STR) DNA markers are hypervariable and informative in *Cannabis sativa*: implications for forensic investigations. *Forensic Science International*, January 2003, vol. 131, no. 1, p. 65-74.

GILMORE, S. and PEAKALL, R. Isolation of microsatellite markers in *Cannabis sativa* L. (marijuana). *Molecular Ecology Notes*, March 2003, vol. 3, no. 1, p. 105-107.

HAKKI, E.E.; UZ, E.; SAG, A.; ATASOY, S. and AKKAYA, M.S. Genotyping of Cannabis sativa L. accessions from Turkey using RAPD and AFLP markers. *Forensic Science International*, 2003, vol. 136, no. 1, p. 31.

HAKKI, E.E.; KAYIS, S.A.; PINARKARA, E. and SAG, A. Inter simple sequence repeats separate efficiently hemp from Marijuana (*Cannabis sativa* L.). *Electronic Journal of Biotechnology*, October 2007, vol. 10, no. 4, p. 570-581.

HILLIG, K.W. Genetic evidence for speciation in *Cannabis* (Cannabaceae). *Genetic Resources and Crop Evolution*, March 2005, vol. 52, no. 2, p. 161-180.

HSIEH, Hsing-Mei; HOU, Rur-Jyun; TSAI, Li-Chin; WEI, Chih-Sheng; LIU, Su-Wen; HUANG, Li-Hung; KUO, Yi-Chen; LINACRE, Adrian and LEE, James Chun-I. A highly polymorphic STR locus in *Cannabis sativa*. *Forensic Science International*, January 2003, vol. 131, no. 1, p. 53-58.

JAGADISH, Vidya; ROBERTSON, James and GIBBS, Adrian. RAPD analysis distinguishes *Cannabis sativa* samples from different sources. *Forensic Science International*, May 1996, vol. 79, no. 2, p. 113-121.

KARUS, Michael and VOGT, Dominik. European hemp industry: Cultivation, processing and product lines. *Euphytica*, January 2004, vol. 140, no. 1-2, p. 7-12.

KOJOMA, Mareshige; IIDA, Osamu; MAKINO, Yukiko; SEKITA, Setsuko and SATAKE, Motoyoshi. DNA fingerprinting of Cannabis sativa using inter-simple sequence repeat (ISSR) amplification. *Planta Medica*, January 2002, vol. 68, no. 1, p. 60-63.

LEGENDRE, P. and ANDERSON, M.J. Distance based redundancy analysis: testing multispecies responses in multifactorial ecological experiments. *Ecological Monographs*, February 1999, vol. 69, no. 1, p. 1-24.

MANTEL, N.A. The detection of disease clustering and a generalized regression approach. *Cancer Research*, February 1967, vol. 27, no. 2, p. 209-220.

NEI, Masatoshi and LI, Wen-Hsiung. Mathematical model for studying genetical variation in terms of restriction endonucleases. *Proceedings of the National Academy of Sciences of the United States of America*, October 1979, vol. 76, no. 10, p. 5269-5273.

PEAKALL, R. and SMOUSE, P.E. Genalex 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*, March 2006, vol. 6, no. 1, p. 288-295.

RANALLI, Paolo and VENTURI, Gianpietro. Hemp as a raw material for industrial applications. *Euphytica*, January 2004, vol. 140, no. 1-2, p. 1-6.

ROGERS, Scott O. and BENDICH, Arnold J. Extraction of DNA from plant tissues. In: GELVIN, S.B. and SCHILPEROORT, R.A. eds. *Plant Molecular Biology Manual*. Kluwer Academic Publisher, Boston, Massachuset, 1988, vol. A6, p. 1-10.

ROHLF, F.J. *NTSYS-pc:* Numerical Taxonomy System. Ver. 2.1. Exeter Publishing, Ltd. Setauket, NY. 2000.

SHIROTA, O.; WATANABE, A.; YAMAZAKI, M.; SAITO, K.; SHIBANO, K.; SEKITA, S. and SATAKE, M. Random amplified polymorphic DNA and restriction fragment length polymorphism analyses of *Cannabis sativa*. *Natural Medicines*, 1998, vol. 52, no. 2, p. 160-166.

SMALL, E. and CRONQUIST, A. Practical and natural taxonomy for cannabis. *Taxon*, 1976, vol. 25, no. 4, p. 405-435.

YAP, V.I. and NELSON, R.J. WINBOOT: A Program for Performing Bootstrap Analysis of Binary Data to Determine the Confidence Limits of UPGMA-Based Dendrograms. IRRI Discussion paper series No. 14, Manila, Philippines; International Rice Research Institute. 1996.

YOON, C.K. Botanical witness for the prosecution. *Science*, 1993, vol. 260, no. 5110, p. 894-895.

RAPD analysis of seized marijuana (Cannabis Sativa I.) in Turkey

APPENDIX

FIGURES



Figure 1. Provincial locations of seized Cannabis accessions on the map of Turkey.



Figure 2. Distribution of 29 *Cannabis* accessions by 2-dimensional principal coordinate analysis. PCo axis 1 and PCo axis 2 account for 13.9% and 7.7% of the variation, respectively.



Figure 3. UPGMA dendrogram Bulk RAPD SM 29 accessions.



Figure 4. The random amplified polymorphic DNA (RAPD) profile of (individual and bulked) *Cannabis* samples, using the random primer L2 which is listed in Table 2.



Figure 4. (continued). Repeat of the random amplified polymorphic DNA (RAPD) profile of (individual and bulked) *Cannabis* samples, using the random primer L2 which is listed in Table 2.

C1 C2 C3 C4 C5 C6 C7 C8 C9 C10 C11 C12 C13 C14 C15 C16 C17 C18 C19 C20 C21 C22 C23 C24 C25 C26 C27 C28 C29 C1 1.00 C2 0.20 1.00 C3 0.19 0.20 1.00 C4 0.16 0.16 0.20 1.00 C5 0.17 0.19 0.18 0.15 1.00 C6 0.19 0.21 0.20 0.18 0.21 1.00 C7 0.19 0.19 0.22 0.17 0.15 0.22 1.00 C8 0.21 0.23 0.24 0.19 0.21 0.24 0.25 1.00 C9 0.17 0.17 0.19 0.17 0.19 0.23 0.19 0.24 1.00 C10 0.18 0.19 0.20 0.17 0.18 0.21 0.20 0.22 0.20 1.00 C11 0.18 0.20 0.23 0.18 0.19 0.24 0.22 0.25 0.20 0.22 1.00 C12 0.19 0.18 0.23 0.18 0.18 0.23 0.23 0.22 0.22 0.20 0.23 1.00 C13 0.19 0.21 0.22 0.18 0.19 0.20 0.21 0.24 0.20 0.19 0.21 0.21 1.00 C14 0.17 0.20 0.22 0.18 0.20 0.21 0.20 0.24 0.21 0.21 0.23 0.21 0.23 1.00 C15 0.17 0.19 0.19 0.16 0.16 0.19 0.19 0.21 0.18 0.18 0.19 0.18 0.21 0.22 1.00 C16 0.18 0.20 0.23 0.18 0.19 0.21 0.19 0.24 0.21 0.21 0.25 0.21 0.23 0.23 0.20 1.00 C17 0.17 0.19 0.22 0.17 0.16 0.20 0.23 0.23 0.19 0.19 0.21 0.21 0.19 0.20 0.20 0.20 1.00 C18 0.18 0.19 0.23 0.19 0.17 0.21 0.23 0.24 0.20 0.21 0.22 0.21 0.22 0.23 0.23 0.24 0.21 1.00 C19 0.17 0.19 0.22 0.18 0.18 0.23 0.20 0.23 0.19 0.20 0.23 0.20 0.22 0.21 0.19 0.23 0.21 0.23 1.00 C21 0.17 0.19 0.19 0.20 0.18 0.19 0.20 0.22 0.19 0.19 0.20 0.17 0.21 0.20 0.19 0.19 0.19 0.21 0.20 0.06 1.00 C22 0.17 0.19 0.22 0.19 0.19 0.23 0.22 0.23 0.19 0.19 0.22 0.22 0.21 0.21 0.18 0.20 0.22 0.19 0.21 0.09 0.19 1.00 C23 0.16 0.17 0.21 0.17 0.17 0.19 0.19 0.23 0.18 0.18 0.21 0.17 0.20 0.20 0.19 0.20 0.17 0.18 0.20 0.08 0.19 0.21 1.00 C24 0.17 0.20 0.24 0.18 0.18 0.18 0.21 0.21 0.25 0.19 0.19 0.21 0.19 0.21 0.19 0.23 0.20 0.19 0.22 0.07 0.19 0.22 1.00 C25 0.19 0.20 0.20 0.17 0.17 0.20 0.22 0.24 0.19 0.17 0.23 0.19 0.21 0.19 0.19 0.20 0.20 0.19 0.20 0.20 0.19 0.21 0.22 1.00 C26 0.21 0.20 0.23 0.21 0.19 0.24 0.22 0.25 0.24 0.21 0.23 0.21 0.21 0.21 0.19 0.23 0.21 0.22 0.08 0.21 0.23 0.23 0.23 0.23 0.21 1.00 C27 0.18 0.20 0.22 0.18 0.20 0.20 0.19 0.24 0.21 0.20 0.22 0.19 0.21 0.22 0.19 0.23 0.19 0.21 0.23 0.08 0.21 0.20 0.23 0.23 0.20 0.28 1.00 C28 0.18 0.19 0.21 0.18 0.17 0.20 0.20 0.23 0.21 0.19 0.22 0.20 0.21 0.22 0.20 0.23 0.21 0.22 0.21 0.07 0.20 0.20 0.21 0.23 0.21 0.25 0.25 1.00 C29 0.16 0.19 0.20 0.17 0.16 0.19 0.19 0.20 0.16 0.18 0.20 0.17 0.18 0.19 0.19 0.19 0.19 0.19 0.19 0.07 0.19 0.20 0.19 0.19 0.19 0.19 0.19 0.17 1.00

Figure 5. Genetic similarity matrix among 29 *Cannabis* accessions, based on RAPD data from SET1 (computed using the Nei and Li's formula (Nei and Li, 1979)).

C1 1.00 C2 0.21 1.00 C3 0.26 0.21 1.00 C4 0.29 0.22 0.30 1.00 C5 0.25 0.20 0.26 0.28 1.00 C6 0.25 0.21 0.25 0.28 0.27 1.00 C7 0.20 0.16 0.24 0.23 0.22 0.22 1.00 C8 0.24 0.21 0.28 0.28 0.25 0.25 0.23 1.00 C9 0 25 0 20 0 26 0 28 0 25 0 26 0 20 0 24 1 00 C10 0.27 0.22 0.29 0.31 0.28 0.27 0.25 0.30 0.29 1.00 C11 0.26 0.21 0.30 0.29 0.27 0.26 0.24 0.30 0.26 0.31 1.00 C12 0.24 0.20 0.28 0.28 0.26 0.26 0.22 0.26 0.26 0.29 0.27 1.00 C13 0.23 0.21 0.26 0.27 0.25 0.25 0.20 0.28 0.25 0.28 0.26 0.26 1.00 C14 0.25 0.20 0.26 0.27 0.26 0.25 0.21 0.26 0.25 0.26 0.26 0.25 0.25 1.00 C15 0.26 0.20 0.27 0.29 0.28 0.26 0.24 0.28 0.26 0.30 0.28 0.28 0.26 0.27 1.00 C16 0.25 0.22 0.29 0.30 0.27 0.28 0.24 0.30 0.29 0.32 0.31 0.30 0.29 0.27 0.30 1.00 C17 0.22 0.18 0.26 0.26 0.24 0.25 0.25 0.28 0.24 0.27 0.28 0.26 0.23 0.25 0.27 0.28 1.00 C18 0.25 0.20 0.27 0.27 0.28 0.26 0.23 0.29 0.27 0.29 0.29 0.26 0.26 0.27 0.29 0.30 0.25 1.00 C19 0.26 0.21 0.28 0.29 0.26 0.27 0.23 0.29 0.27 0.30 0.28 0.28 0.28 0.28 0.28 0.30 0.26 0.30 1.00 C20 0.08 0.05 0.09 0.10 0.09 0.10 0.11 0.09 0.09 0.10 0.11 0.08 0.08 0.07 0.09 0.10 0.09 0.09 0.08 1.00 C21 0.22 0.18 0.24 0.26 0.25 0.24 0.22 0.25 0.23 0.26 0.24 0.24 0.25 0.25 0.25 0.25 0.22 0.26 0.27 0.07 1.00 C22 0.18 0.15 0.20 0.20 0.19 0.18 0.20 0.21 0.19 0.20 0.20 0.20 0.20 0.19 0.20 0.22 0.20 0.21 0.21 0.21 0.09 0.20 1.00 C23 0.21 0.17 0.22 0.24 0.22 0.22 0.22 0.24 0.21 0.24 0.23 0.21 0.22 0.22 0.22 0.23 0.23 0.22 0.23 0.11 0.21 0.23 1.00 C24 0.20 0.17 0.24 0.25 0.23 0.22 0.22 0.24 0.21 0.25 0.25 0.22 0.22 0.22 0.23 0.25 0.23 0.23 0.23 0.09 0.22 0.19 0.26 1.00 C25 0.21 0.18 0.23 0.24 0.22 0.21 0.24 0.24 0.22 0.26 0.25 0.22 0.23 0.20 0.23 0.25 0.23 0.22 0.23 0.09 0.22 0.21 0.23 0.23 1.00 C26 0.22 0.19 0.23 0.26 0.22 0.23 0.23 0.24 0.23 0.26 0.25 0.24 0.22 0.22 0.25 0.26 0.25 0.24 0.25 0.10 0.23 0.21 0.26 0.25 0.24 1.00 C27 0.18 0.16 0.22 0.22 0.21 0.21 0.22 0.24 0.20 0.24 0.25 0.21 0.21 0.21 0.23 0.26 0.22 0.24 0.22 0.09 0.21 0.19 0.22 0.22 0.23 0.26 1.00 229 0.18 0.17 0.22 0.22 0.22 0.22 0.22 0.22 0.19 0.23 0.22 0.21 0.20 0.19 0.21 0.22 0.21 0.21 0.23 0.08 0.23 0.17 0.20 0.21 0.22 0.21 0.22 0.20 1.00

Figure 6. Genetic similarity matrix among 29 *Cannabis* accessions, based on RAPD data from SET2 (computed using the Nei and Li's formula (Nei and Li, 1979)).