

***In vitro* regeneration and genetic fidelity of *Tigridia pavonia* (L.f.) DC.**

José Luis Piña-Escutia

Centro de Investigación y Estudios Avanzados en Fitomejoramiento
Facultad de Ciencias Agrícolas
Universidad Autónoma del Estado de México
Toluca, Estado de México, México

Luis Miguel Vázquez-García

Centro de Investigación y Estudios Avanzados en Fitomejoramiento
Facultad de Ciencias Agrícolas
Universidad Autónoma del Estado de México
Toluca, Estado de México, México

Amaury Martín Arzate-Fernández*

Centro de Investigación y Estudios Avanzados en Fitomejoramiento
Facultad de Ciencias Agrícolas
Universidad Autónoma del Estado de México
Carretera Toluca-Ixtlahuaca km 11.5
Campus Universitario "El Cerrillo" 50200
Toluca, Estado de México, México
E-mail: amaury1963@yahoo.com.mx

Financial support: We are grateful to Consejo Nacional de Ciencia y Tecnología (CONACYT) from México for the scholarship give to JLPE (2005-183935), and to the RED TIGRIDIA (SINAREFI) for covering the publishing payment.

Keywords: genetic integrity, micropropagation, oceloxóchitl.

Abbreviations: 2,4-D: 2,4-Dichlorophenoxyacetic acid

AC: activated charcoal

BAP: 6-Benzylaminopurine

ISSR: Inter Simple Sequence Repeats

MS: murashige and skoog

NAA: Naphtalen Acetic Acid

N₆: Chu basal medium

PCR: Polymerase Chain Reaction

Plants of *Tigridia pavonia* (L.f.) DC were regenerated from twin-scaling explants cultured on Murashige and Skoog and N₆ basal medium. The highest formation of shoots per responding explant was obtained on N₆ medium supplemented with 4.5 µM 2,4-dichlorophenoxy acetic acid in combination with 2.2 µM benzylaminopurine. Shoots rooted readily on N₆ basal medium supplemented with 1 g l⁻¹ activated charcoal and 2.6 µM naphtalenacetic acid. The rooted shoots achieved 100% survival. Inter Simple Sequence Repeat analysis was carried out to check for possible genetic alterations in plants obtained after two consecutive subcultures. The results revealed that the recovered plants did not exhibit any type of polymorphism.

Tigridia pavonia (L.f.) DC., also known as tiger flower or oceloxóchitl, is a native bulbous plant of Mexico and it was used by the Aztecs as an ornamental, and for food and medicinal purposes. Although in Mexico there is no record of its commercial production, it is commercialized as

garden plant in Europe, Asia and Australia where is known for its flowering longevity of only 10-14 hrs. This species have been strongly extracted, and its phylogenetic germplasm could be lost (Vázquez-García et al. 2001).

T. pavonia is usually propagated in one of two ways: by seed or by bulb. The use of seeds is not a practical propagation method because of the semi-compatibility and low germination rate. Moreover, to obtain stock bulbs, from two to three years is required because of the slow growth of the bulbs until flowering (Vázquez-García et al. 2001).

Tissue culture can be employed as an alternative to conventional methods of vegetative propagation with the goal of enhancing the rate of multiplication of desired genotypes (Paek and Murthy, 2002). The technique of twin-scaling has been effective for tissue culture propagation of *Iris hollandica* (Fidalgo et al. 2005) and other bulbous species, including *Eucrosia stricklandii* (Colque et al. 2002), *Narcissus* (Santos et al. 2002) and *Sternbergia clusiana* (Sawsan and Fattash, 2005).

*Corresponding author

Despite the advantages of *in vitro* propagation, factors such as growth regulators, time of culture and media composition appear to be capable of inducing *in vitro* genetic variability in the regenerated plantlets (Silvarolla, 1992). Several approaches have been applied for identification of variants among micropropagated plants including: phenotypic variation (Kancherla and Bhalla, 2003), karyotypic analysis of chromosomes (Bennici et al. 2004), and biochemical analysis (Punja et al. 2004). A major disadvantage of these techniques is the availability of only limited numbers of informative markers since the number of assays to detect them is relatively low. Therefore, the percentage coverage of the genome is inadequate to make a comprehensive study of genetic diversity. Furthermore, the markers can be influenced by environmental factors, and the expression differences can cause confusion in the interpretation of the results (Vicente and Fulton, 2003).

Inter Simple Sequence Repeat (ISSR) markers, which use anchored primers to amplify simple sequence repeats without the requirement for prior sequence information, is a very simple, fast, cost-effective, highly discriminative and reliable technique (Reedy et al. 2002), and they have been successfully applied to detect the genetic similarities or

dissimilarities in micropropagated material in *Prunus dulcis* (Martins et al. 2004), *Tulipa* spp. (Podwyszynska et al. 2006), *Musa* spp. (Lakshmanan et al. 2007), and *Vitex negundo* (Ahmad et al. 2008).

To date, there have been no reports on *in vitro* propagation of *T. pavonia*. Here, we describe an efficient protocol for direct plant regeneration from twin-scale explants of *T. pavonia* along with the ISSR analysis to determine the genetic fidelity of the regenerated plants.

MATERIALS AND METHODS

Plant material and disinfection

T. pavonia (L.f.) DC bulbs were obtained from plants grown in a greenhouse, and were used as source material to initiate *in vitro* cultures.

The outer scales, one-third of the upper part and the dry remains of the previous root disks of the bulbs were removed, and the bulbs were placed in a flask with 10 ml of antibacterial soap (Reckitt Benckiser Inc. USA) and rinsed in running tap water for 30 min. Bulbs were then soaked for 1 hr in a solution containing 1% (w/v) Benomyl (metil-1-




Figure 1. Plantlet regeneration of *Tigridia pavonia* (L.f) DC from twin-scales.

(a) Shoots (arrows) of 0.5-1 cm in length developing at the basal portion of the scale, two weeks after culture in MS medium supplemented with 4.5 μ M 2,4-dichlorophenoxyacetic acid (2,4-D) and 2.2 μ M 6-benzilaminopurine (BAP).
 (b) Shoot proliferation after 8 weeks culture in N_6 medium supplemented with 4.5 μ M of 2,4-D and 2.2 μ M of BAP.
 (c) Rooting of shoots, after 3 weeks on N_6 basal medium supplemented with 1 g l⁻¹ activated charcoal (AC) and 2.6 μ M naphtalenacetic acid (NAA).
 (d) A regenerated plantlet after 16 weeks in culture.
 (e) Plants during acclimation 20 weeks after culture initiation.
 (f) ISSR profiles generated by the primer ASSR-15 using *T. pavonia* individuals after two subcultures (lines 1-9) and from a mother plant grown under greenhouse conditions (line 10). M, represents the DNA ladder. bp, base pairs.

butylcarbomol-2-bencinidazol-carbamate a.i. 50% w/w) and 1% (w/v) Agry-Gent (gentamycin sulphate a.i. 2% w/w and Oxytetracyclin chloride a.i. 6% w/w). Afterwards, in a laminar flow cabinet, bulbs were surface disinfected with 70% ethanol for 2 min, then immersed in 5% sodium hypochlorite for 10 min followed by 5% calcium hypochlorite for 15 min; finally, they were washed three times with distilled sterilized water.

Each disinfected bulb was longitudinally sectioned to obtain explants formed by segments of twin-scales, 2-3 mm wide and 6-8 mm length, joined by a thick segment (2-3 mm) of basal tissue. The explants were cultured upright with the basal plate tissue inserted into the culture medium.

Medium culture and shoot induction

For shoot induction, two basal media: MS (Murashige and Skoog, 1962) and N₆ (Chu et al. 1975) and 0 and 2.2 µM concentrations of 6-benzylaminopurine (BAP) and 0, 4.5, 9 and 13.5 µM 2,4-dichlorophenoxyacetic acid (2,4-D) in all possible combinations, were tested. The basal media were supplemented with 30 g l⁻¹ sucrose and 1 g l⁻¹ activated charcoal. All media were adjusted to a pH of 5.7 ± 0.1 and then 2.5 g l⁻¹ gelrite was added before autoclaving at 121°C and 1.1 kg/cm² for 20 min. Eight explants were placed on each sterile Petri dish (90 mm diameter) containing 20 ml of medium, then sealed with ParafilmTM. Cultures were incubated at 25 ± 1°C under cool-white fluorescent tubes with a light intensity of 34 ± 5 µmol m⁻² s⁻¹ and 16 hrs photoperiod. Explants were subcultured every 30 days.

Rooting and acclimatization

Shoots of 4-6 cm in length were separated from the original explant, and subcultured in glass flasks (500 ml, covered with aluminum plugs) on N₆ basal medium supplemented with 1 g l⁻¹ activated charcoal and 2.6 µM naphthalenacetic acid (NAA) to induce rooting. The incubation conditions and light intensity were the same as stated above.

Plants with a mean height of 15 cm and a well-developed root system were removed from glass flasks and transferred to *ex vitro* conditions. For this purpose, they were rinsed free of tissue medium culture with distilled water, and potted in plastic pots (25 cm³) containing a sterilized mixture (1:1) of compost and agrolita (DICALITE®). Clear polyethylene bags (300 cm²) were inverted over the pots and secured with rubber bands. The bags were ventilated by removing the rubber bands after 2 weeks; bags were removed 1 week later and the plants were then transferred to a greenhouse.

Statistical analysis

To evaluate the effect of each treatment on shoot induction, a factorial analysis and LSD test were performed using the data of number of shoots per explant formed 8 weeks after

culture initiation. The treatments were arranged in a completely randomized design. Data were analyzed with Statistical Analysis System software version 8.0 (SAS Institute, 1999).

DNA extraction and ISSR analysis

DNA was extracted from 100 mg of fresh leaves derived from one to four shoots from each treatment after two subcultures (30 d each) and from a mother plant grown under greenhouse conditions. DNA extraction was carried out according to Arzate-Fernández et al. (2005). The DNA samples were stored at -20°C until be used.

Five anchored simple sequence repeat primers (3'-ASSR02, 3'-ASSR15, 3'-ASSR20, 3'-ASSR29 and 3'-ASSR35) (Yamagishi et al. 2002) were used for the Polymerase Chain Reaction (PCR). Amplification reactions were performed in 10 µl containing 1 µl of 10X PCR ammonium buffer (1.5 mM MgCl₂), 0.5 µl of MgCl₂ (25 mM), 1 µl of dNTPs (10 mM) (APPLIED BIOSYSTEMS®), 1 µl of the primer (20 µM) (INVITROGENTM), 0.1 U of Taq DNA polimerase (MERCURY REAGENTSTM) and 1 µl of DNA (10 ng). DNA amplification was performed in a thermal Mastercycler gradient (EPPENDORF®). The cycles for the primers 3'-ASSR02 and 3'-ASSR15 were according to Yamagishi et al. (2002) and Arzate-Fernández et al. (2005) with modifications comprising the following: an initial cycle of 9 min at 94°C, 1 min at 46°C and 1 min at 72°C; then 45 cycles of 1 min at 94°C, 1 min at 46°C and 1 min at 72°C followed by one final extension cycle of 9 min at 94°C, 1 min at 46°C and 10 min at 72°C. The cycles for the primers 3'-ASSR20, 3'-ASSR29 and 3'-ASSR35 were according to Yamagishi et al. (2002). Amplified products were separated by 1% agarose gel (w/v) (SIGMA®) electrophoresis at 100 V. The gel was stained with 3 µl of ethidium bromide and the DNA bands were photographed under ultraviolet light using a UV Transilluminator BioDoc-It Imaging System (UVPTM). The sizes of the amplification products were estimated with a DNA ladder (100 bp to 3.0 kb) (FERMENTASTM). The ISSR profiles were analyzed on the basis of the presence or absence of individual ISSR bands.

RESULTS AND DISCUSSION

Shoot induction

The response of explants cultured on MS and N₆ medium culture with 2,4-D and BAP on shoot induction of *T. pavonia* is shown in Table 1. After two weeks of culture, shoots arose from the basal portion of the bulb scales (Figure 1a) in some treatments. In those responding treatments, the original explant maintained its organogenic capacity for more than 12 weeks; during which adventitious shoot proliferation was observed.

The period for shoot induction reported here, is faster in comparison with other bulbous species such as *Eucrosia*

Table 1. Number of shoots induced in *T. pavonia* in response to plant growth regulators and medium. Values were obtained 8 weeks after culture initiation and represent the means \pm standard deviations.

PGR: 2,4-D : BAP (μ M)	MS	N ₆	
0.0 : 0.0	0.6 \pm 0.4 ef	1.6 \pm 0.4 cde	
4.5 : 0.0	2.6 \pm 0.4 bc	2.6 \pm 0.4 bc	
9.0 : 0.0	2.6 \pm 0.4 bc	2.3 \pm 0.9 bcd	
13.5: 0.0	2.6 \pm 0.4 bc	0.0 f	
0.0 : 2.2	0.6 \pm 0.4 ef	1.3 \pm 0.4 de	
4.5 : 2.2	3.0 \pm 0.0 ab	4.0 \pm 0.8 a	
9.0 : 2.2	1.6 \pm 0.4 cde	3.0 \pm 0.0 ab	
13.5: 2.2	0.6 \pm 0.4 ef	1.6 \pm 0.4 cde	
Source of variation	Df	Mean square	F
Media culture (A)	1	0.52	1.25ns
2,4-D (B)	3	10.40	24.98***
BAP (C)	1	0.02	0.05ns
AxB	3	1.46	3.52ns
AxC	1	7.52	18.05***
BxC	3	0.85	2.05ns
AxBxC	3	1.79	4.32ns
Error	32		
Total	47		

PGR = Plant growth regulators; 2,4-D = 2,4-dichlorophenoxyacetic acid; BAP = 6-benzilaminopurine. Values followed by different letters are significantly different (LSD, 0.05). ***P = \leq 0.001, ns = non significant.

stricklandii, where a response was obtained in four weeks (Colque et al. 2002), and *Narcissus asturiensis*, where it occurred at eight weeks (Santos et al. 2002).

In this study, up to 90% of the explants developed shoots and the number of shoots per explant varied between 0 and 4 (Table 1). Although the number of shoots per explants was low, it was similar to shoot numbers reported in other bulbous plants; for example, in *Eucumis autumnalis* 1.0 to 3.7 shoots per explant were obtained (Ault, 1995), and in *Eucrosia stricklandii* 0.7 to 2.2 shoots per explant was observed (Colque et al. 2002).

The effect of 2,4-D and of the interaction of medium and BAP on shoot induction, after eight weeks, was highly significant (Table 1). It has been reported that auxin is an effective PGR for shoot induction and in combination with a cytokinin, is essential for shoot induction (Paek and Murthy, 2002; Santos et al. 2002). In this study, the combination of 2,4-D with BAP stimulated shoot induction (Figure 1b). Our results agree with those reported by Boltenkov and Zarembo (2005) in *Iris ensata*, and Anbari et al. (2007) in *Narcissus papyraceus*, who reported that combinations of 2,4-D and BAP were necessary for shoot induction.

The combination of low levels of 2,4-D (4.5 μ M) with BAP (2.2 μ M), produced three and four shoots per explant in MS and N₆ media respectively (Table 1): but in high levels of 2,4-D (13.5 μ M) with BAP (2.2 μ M), the response was minor (0.6 and 1.6 shoots, respectively) and the shoots formed in these levels were smaller to those obtained in low levels. Similar results were observed by Aftab et al. (2008) who reported sporadic single shoot regeneration in *Gladiolus hybridus* using high levels of 2,4-D (9 μ M). In contrast, Mata-Rosas et al. (2006) obtained the best shoot formation in *Magnolia dealbata* using low concentrations of 2,4-D (0.5 μ M). The results of this study, and the above mentioned work, suggest that low levels of 2,4-D in combination with BA, are enough for *in vitro* shoot induction, but the use of 2,4-D at high levels affects the normal development of plant tissues, which was also observed by Filippi et al. (2001) and Steinitz et al. (2003) in *Musa* spp and *Capsicum annum*, respectively.

Although the effect of MS and N₆ on shoot induction was not significant, the rate of shoot induction was slightly higher in N₆ medium in comparison with the MS medium (Table 1). MS medium has been reported as one of the best medium for shoot induction (Colque et al. 2002; Santos et al. 2002; Sawsan and Fattash, 2005) while N₆ medium has been used in few species, for the same purpose. MS medium differs significantly from N₆ medium with respect to the content of ammonium and nitrate ions, and also to a lesser degree with the content of phosphate ions. Given that both elements (nitrogen and phosphorus) are consumed quickly by the explants cultivated *in vitro*, they can be limiting factors in tissue culture growth (Ramage and Williams, 2002). Also, it has been observed that high ammonium levels have a negative effect on growth and morphogenesis. Considering this, it is possible that the highest shoot formation response observed in the N₆ medium in this study, may have been the result of the higher quantity of phosphorous and the smaller quantity of ammonium present in the N₆ medium in comparison with MS medium.

It has been reported that germination of *T. pavonia* seeds is very low, while its propagation by bulb can produce four bulbs with same number of plants in a two to three year period (Vázquez-García et al. 2001). In contrast, the *in vitro* regeneration protocol described in this study may produce 48 plants from one bulb in a period of five months (an average of 12 scales per bulb and four plantlets per scale). This is the first report for *in vitro* culture of *T. pavonia* and the procedure described is an effective regeneration protocol, although further experiments on the explant size will be required to improve shoot induction.

Rooting and acclimatization

It has been reported that the AC improved shoot formation, plant recovery and rooting (Arzate-Fernández et al. 2007), and that low NAA concentrations (0.5-2.5 μ M) are optimal for the root induction (Colque et al. 2002; Aftab et al. 2008;

Torabi-Giglou and Hajieghrari, 2008). In this study, the addition of AC and NAA showed a positive effect because all adventitious plantlets (4-6 cm in length) continued developing on N₆ basal medium supplemented with 1 g l⁻¹ AC and 2.6 μ M NAA, and after 3 weeks culture, roots formed (Figure 1c). Our results are in agreement with Ault (1995), and Mata-Rosas et al. (2006) who also reported beneficial effects of AC and NAA on *in vitro* root stimulation in *Eucrosia autumnalis* and *Magnolia dealbata* respectively.

Pospíšilová et al. (1999) mentioned that, through *in vitro* culture, the plantlets can develop morphological, anatomical and physiological abnormalities. Also, regenerated plantlets often die during transfer from *in vitro* to the *ex vitro* conditions. In contrast, in this study, sixteen weeks after culture initiation, more than 100 regenerated plants (10-15 cm in length) with roots (4-6 cm) and 2-4 green leaves were obtained (Figure 1d) and these were established with a mixture of compost and agrolita and transferred to a greenhouse with a 100% success rate (Figure 1e). It is possible that the development of complete plants and the *ex vitro* handling protocol may have contributed to the adaptation success.

ISSR analysis

Genetic uniformity is one of the most important prerequisites for the successful micropropagation of any crop species. Nevertheless, a major problem encountered in cells grown *in vitro* is the occurrence of genetic variation. Although the origin of this variation is unclear, it is considered that there are two main factors: an intrinsic factor, which largely depends on the genetic stability of the explants, and an extrinsic factor, depending on culture media and particularly growth regulators (in particular 2,4-D) (Smith, 1998).

The ISSR technique has been used successfully for analyzing the genetic stability of species propagated through *in vitro* shoot formation (Martins et al. 2004; Podwyszynska et al. 2006; Lakshmanan et al. 2007; Ahmad et al. 2008). In this study, ISSR analysis did not reveal any type of polymorphism between randomly selected *in vitro* plants of *T. pavonia* and the mother plant, indicating the clonal nature of progeny (Figure 1f). This observation constitutes an advantage for the *in vitro* regeneration procedure because it minimizes the risk of genetic variation.

The five ASSR primers tested in this study generated 21 scorable band classes (all monomorphic) ranging in size from 500 bp (ASSR-20) to 1700 bp (ASSR-35). The number of bands for each primer was from 3 (ASSR-02 and ASSR-15), 4 (ASSR-29), 5 (ASSR-35) to 6 (ASSR-20) with an average of 4.2 bands per ASSR primer. An example of PCR amplification products obtained with the primer ASSR-15 is given in Figure 1f. This findings agree with those reported by others (Lakshmanan et al. 2007; Ahmad

et al. 2008; Bhatia et al. 2009) who have pointed out the efficacy of 3'-anchored primers for giving a higher number of clearly definable bands and being better than other markers such as RAPDs in the assessment of clonal fidelity of micropropagated plants (Martins et al. 2004).

CONCLUDING REMARKS

The best conditions for *in vitro* shoot induction using twin-scale explants of *T. pavonia* was the treatment with 4.5 μ M 2,4-D and 2.2 μ M BAP in N₆ medium supplemented with 1 g l⁻¹ of activated charcoal, and resulted in four shoots per explant. Thus, 48 plants can be produced from one bulb within five months of culture initiation. N₆ medium supplemented with 1 g l⁻¹ activated charcoal and 2.6 μ M NAA can be used to induce rooting. The protocol described is an effective regeneration procedure for *T. pavonia* because no somaclonal variation was observed among the regenerated plants based on ISSR analysis.

REFERENCES

- AFTAB, Faheem; ALAM, Memoona and AFRASIAB, Humera. *In vitro* shoot multiplication and callus induction in *Gladiolus hybridus* Hort. *Pakistan Journal Botanical* [on line]. April 2008, vol. 40, no. 2 [cited 20 December 2008]. Available from Internet: [http://www.pakbs.org/pjbot/PDFs/40\(2\)/PJB40\(2\)517.pdf](http://www.pakbs.org/pjbot/PDFs/40(2)/PJB40(2)517.pdf).
- AHMAD, N.; WALI, S.A. and ANIS, M. *In vitro* production of true-to-type plants of *Vitex negundo* L. from nodal explants. *Journal of Horticultural Science & Biotechnology*, 2008, vol. 83, no. 3, p. 313-317.
- ANBARI, Saghi; TOHIDFAR, Masoud; HOSSEINI, Ramin and HADDAD, Rahim. Somatic embryogenesis induction in *Narcissus papyraceus* cv. Shirazi. *Plant Tissue Culture and Biotechnology* [on line]. June 2007, vol. 6, no. 4 [cited 10 November 2008]. Available from Internet: <http://www.banglajol.info/index.php/PTCB/article/view/1119/1142>.
- ARZATE-FERNÁNDEZ, Amaury-M.; MIWA, Makoto; SHIMADA, Tomohide; YONEKURA, Tetsushi and OGAWA, Kazuo. Genetic diversity of miyamasukashi-yuri (*Lilium maculatum* Thunb. var. *Bukosanense*), an endemic endangered species at Mount Buko, Saitama, Japan. *Plant Species Biology*, April 2005, vol. 20, no. 1, p. 57-65.
- ARZATE-FERNÁNDEZ, Amaury-M.; MIWA, Makoto; SHIMADA, Tomohide; YONEKURA, Tetsushi and OGAWA, Kazuo. *In vitro* propagation of miyamasukashi-yuri (*Lilium maculatum* Thunb. var. *Bukosanense*), an endangered plant species. *Revista Fitotecnica Mexicana*, 2007, vol. 30, no. 4, p. 373-379.
- AULT, J.R. *In vitro* propagation of *Eucomis autumnalis*, *E. comosa*, and *E. zambesiaca* by twin scaling. *HortScience*, December 1995, vol. 30, no. 7, p. 1441-1442.
- BENNICI, Andrea; ANZIDEI, Maria and VENDRAMIN, Giovanni G. Genetic stability and uniformity of *Foeniculum vulgare* Mill. regenerated plants through organogenesis and somatic embryogenesis. *Plant Science*, January 2004, vol. 166, no. 1, p. 221-227.
- BHATIA, R.; SINGH, K.P.; JHANG, T. and SHARMA, T.R. Assessment of clonal fidelity of micropropagated gerbera plants by ISSR markers. *Scientia Horticulturae*, January 2009, vol. 119, no. 2, p. 208-211.
- BOLTENKOV, E.V. and ZAREMBO, E.V. *In vitro* regeneration and callogenesis in tissue culture of floral organs of the genus *Iris* (Iridaceae). *Biology Bulletin*, March 2005, vol. 32, no. 2, p. 138-142.
- CHU, C.C.; WANG, C.C.; SUN, C.S.; HSU, C.; YIN, K.C.; CHU, C.Y. and BI, F.Y. Establishment of an efficient medium for anther culture of rice through comparative experiments on the nitrogen sources. *Scientia Sinica*, 1975, vol. 18, p. 659-668.
- COLQUE, R.; VILLADOMAT, F.; BASTIDA, J. and CODINA, C. Micropropagation of the rare *Eucrosia stricklandii* (Amaryllidaceae) by twin-scaling and shake liquid culture. *Journal of Horticultural Science & Biotechnology*, 2002, vol. 77, no. 6, p. 739-743.
- FIDALGO, F.; SANTOS, A.; OLIVEIRA, N.; SANTOS, I. and SALEMA, R. Induction of somatic embryogenesis in *Iris hollandica* Hort. Cv. "Bronze Queen". *Journal of Horticultural Science & Biotechnology*, 2005, vol. 80, no.1, p. 135-138.
- FILIPPI, Silvia Balbão; APPEZZATO-DA-GLÓRIA, Beatriz and RODRIGUEZ, Adriana Pinheiro Martinelli. Histological changes in banana explants, cv. Nanicão (*Musa* spp., Group AAA), submitted to different auxins for induction of somatic embryogenesis. *Revista Brasileira de Botânica* [on line]. December 2001, vol. 24, no. 4, suppl. p. 595-602.
- KANCHERLA, L.S. and BHALLA, P.M. Phenotypic variations in micropropagated Australian ornamental climber *Pandorea pandorana*. *Acta Horticulturae*, 2003, no. 616, p. 463-466.
- LAKSHMANAN, Venkatachalam; REDDAMPALLI, Venkataramareddy Sreedhar and NEELWARNE, Bhagyalakshmi. Molecular analysis of genetic stability in long-term micropropagated shoots of banana using RAPD and ISSR markers. *Electronical Journal of Biotechnology*, 15 January 2007, vol. 10, no. 1, p. 106-113.
- MARTINS, M.; SARMENTO, D. and OLIVEIRA, M.M. Genetic stability of micropropagated almond plantlets, as assessed by RAPD and ISSR markers. *Plant Cell Reports*, December 2004, vol. 23, no. 7, p. 492-496.

- MATA-ROSAS, Martín; JIMÉNEZ-RODRÍGUEZ, Ángel and CHÁVEZ-AVILA, Víctor. Somatic embryogenesis and organogenesis in *Magnolia dealbata* Zucc. (Magnoliaceae), an endangered, endemic Mexican species. *HortScience*, 2006, vol. 41, no. 5, p. 1325-1329.
- MURASHIGE, Toshio and SKOOG, Folke. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, July 1962, vol. 15, no. 3, p. 473-497.
- PAEK, K.Y. and MURTHY, H.N. High frequency of bulblet regeneration from bulb scale sections of *Fritillaria thunbergii*. *Plant Cell Tissue and Organ Culture*, March 2002, vol. 68, no. 3, p. 247-252.
- PODWYSZYNSKA, M.; NIEDOBA, K.; KORBIN, M. and MARASEK, A. Somaclonal variation in micropropagated tulips determined by phenotype and DNA markers. *Acta Horticulturae*, 2006, no. 714, p. 211-220.
- POSPÍŠILOVÁ, J.; TICHÁ, I.; KADLEČEK, P.; HASEL, D. and PLZÁKOVÁ, Š. Acclimatization of micropropagated plants to *ex vitro* conditions. *Biologia Plantarum*, December 1999, vol. 42, no. 4, p. 481-494.
- PUNJA, Z.K.; FEENEY, M.; SCHLUTER, C. and TAUTORUS, T. Multiplication and germination of somatic embryos of american ginseng derived from suspension cultures and biochemical and molecular analyses of plantlets. *In vitro Cellular and Development Biology Plant*, May 2004, vol. 40, no. 3, p. 329-338.
- RAMAGE, Carl M. and WILLIAMS, Richard R. Mineral nutrition and plant morphogenesis. *In vitro Cellular and Development Biology-Plant*, March 2002, vol. 38, no. 2, p. 116-124.
- REDDY, M.P.; SARLA, N. and SIDDIQ, E.A. Inter simple sequence repeat (ISSR) polymorphism and its application in plant breeding. *Euphytica*, November 2002, vol. 128, no. 1, p. 9-17.
- SANTOS, A.; FIDALGO, F.; SANTOS, I. and SALEMA, R. *In vitro* bulb formation of *Narcissus asturiensis* a threatened species of the *Amaryllidaceae*. *Journal of Horticultural Science & Biotechnology*, 2002, vol. 77, no. 2, p. 149-152.
- SAS. *SAS User's guide: Statistics version 8.0*. Cary, NC: SAS Institute Inc. 1999. 1243 p. ISBN 1580254942.
- SAWSAN, A.O. and FATTASH I.A. *In vitro* propagation of an endangered medicinal bulbous plant *Sternbergia clusiana* Ker-Gawler (*Amaryllidaceae*). *Journal of Horticultural Science & Biotechnology*, 2005, vol. 80, no. 4, p. 399-402.
- SILVAROLLA, M.B. Plant genomic alterations due to tissue culture. *Journal of Brazilian Association Advanced Science*, 1992, no. 44, p. 329-335.
- SMITH, M.K. A review of factors influencing the genetic stability of micropropagated banana fruits. *Fruits*, 1998, vol. 43, no. 4, p. 219-223.
- STEINITZ, Benjamin; KÜSEK, Mustafa; TABIB, Yona; PARANI, Ilan and ZELCER, Aaron. Pepper (*Capsicum annum* L.) regenerants obtained by direct somatic embryogenesis fail to develop a shoot. *In vitro Cellular and Development Biology Plant*, May 2003, vol. 39, no. 3, p. 296-303.
- TORABI-GIGLOU, Mousa and HAJIEGHRARI, Behzad. *In vitro* study on regeneration of *Gladiolus grandiflorus* corm calli as affected by plant growth regulators. *Pakistan Journal Biological Sciences*, 2008, vol. 11, no. 8, p. 1147-1154.
- VÁZQUEZ-GARCÍA, L.M.; NORMAN, M.T.H. and CORONA, R.M.C. Oceloxochitl *Tigridia pavonia* (L. f.) DC. *Colección: Ciencias Naturales y Exactas, Serie: Ciencias Agrícolas*. Universidad Autónoma del Estado de México, Toluca, Estado de México. México, 2001. 69 p. ISBN 9-68-835691-3.
- VICENTE, M. Carmen and FULTON, Theresa. *Tecnologías de Marcadores Moleculares para Estudios de Diversidad Genética de Plantas: Módulo de Aprendizaje*. Illus. Nelly Giraldo. Instituto Internacional de Recursos Fitogenéticos (IPGRI), Roma, Italia. 2003. vol. 1, 52 p. ISBN 92-9043-666-2.
- YAMAGISHI, Masumi; ABE, Hiromi; NAKANO, Michihan and NAKATSUKA, Akira. PCR-based molecular markers in Asiatic hybrid lily. *Scientia Horticulturae*, December 2002, vol. 96, no. 1-4, p. 225-234.