

Thirteen Years' Experience of Diaphragmatic Injury in Children from the Post Graduate Institute of Medical Sciences (PGIMS), Rohtak, India

Kamal Nain RATTAN¹, Rajat NARANG¹, Seema ROHILLA², Sarita MAGGU², Dhara B DHAULAKHANDI³

Submitted: 6 Apr 2008
Accepted: 30 Aug 2010

¹ Department of Pediatric Surgery, Post Graduate Institute of Medical Sciences, Pt BD Sharma University of Health Sciences, Rohtak 124 001, Haryana, India

² Department of Radiodiagnosis and Imaging, Post Graduate Institute of Medical Sciences, Pt BD Sharma University of Health Sciences, Rohtak 124001, Haryana, India

³ Department of Biotechnology and Molecular Medicine, Post Graduate Institute of Medical Sciences, Pt BD Sharma University of Health Sciences, Rohtak 124 001, Haryana, India

Abstract

Background: Diaphragmatic hernia is migration of abdominal viscera into the thoracic cavity through a defect in the diaphragm. In children, it is mostly congenital; traumatic diaphragmatic hernia being less common. This study aimed to review our experience with traumatic diaphragmatic rupture (TDR) and to identify the clinical findings and diagnostic modality that may help in early diagnosis and prompt therapy.

Methods: The study involved 11 children (1–18 years old) with TDR who were hospitalised between 1993 and 2005. In addition to clinical examination, a plain X-ray of the chest and abdomen, an ultrasound, barium studies, and a computerised tomography (CT) scan were used to evaluate the patients.

Results: All of the diaphragmatic ruptures occurred on the left side, with 10 occurring in the posterolateral part and 1 near the oesophageal hiatus. Two of our patients presented 7 and 10 days after the injury, and 1 patient presented 1 year after the trauma.

Conclusion: TDR should remain a diagnostic possibility in children. These patients are best assessed using a CT scan. New research on stem cells and tissue-engineered bioprosthesis may pave the path for better future therapies in these cases.

Keywords: acute respiratory distress syndrome, child, diagnosis, diaphragmatic hernia, medical imaging, rupture, trauma

Introduction

Diaphragmatic rupture is an uncommon but well-recognised complication of trauma that consists of 1.0%–5.8% of admissions into a trauma unit (1). It occurs in 0.5%–8.0% of adult trauma patients (2). In paediatric patients, it is estimated to occur at a rate of 4%–6%, and its presence has been reported to indicate high impact. It is associated with other severe injuries in 44%–94% of cases (3).

Traumatic diaphragmatic injuries are usually caused by blunt abdominal trauma or penetrating injuries (2); they were first described by Ambrose Paire in 1579 (4). Due to their rarity in infants and children, such injuries can be overlooked if

unsuspected. Delayed presentation can lead to life-threatening complications as a result of organ herniation and strangulation (3).

Subjects and Methods

The study included children (1–18 years old) with traumatic diaphragmatic rupture (TDR) who were admitted to the Department of Pediatric Surgery of Pt BD Sharma Post Graduate Institute of Medical Sciences, Rohtak, Haryana, India, between 1993 and 2005. The following information was recorded for each patient: age, gender, duration between the trauma and hospital admission, type of injury, clinical/radiological

findings, relevant information regarding traumatic site, herniated organs into the thorax, associated injuries, and patient's outcome.

Result

In the study duration, 11 patients were treated for TDR in our hospital; 8 boys and 3 girls. Their ages ranged from 1–18 years (mean 9.5 years). The TDR occurred following blunt trauma in 9 cases and a penetrating injury in 2 cases (elaborated in Table 1). Road traffic accidents were the most common cause. The most common clinical findings were respiratory distress and abdominal pain.

All of the ruptures occurred on the left side. The ruptures were localised in the posterolateral parts in 10 cases and near the oesophageal hiatus in 1 case. Eight of the patients were admitted soon after the trauma, while 1 patient was admitted after 7 days of injury. One patient was referred to our hospital after 10 days of injury with bilious contents from the thoracic drainage site (admitted to a peripheral hospital in the initial days following the injury). The chest tube had been inserted into the stomach because it was thought to be a case of hydropneumothorax. Another patient presented with abdominal pain after 1 year of trauma with the small and large bowels herniated into the thoracic cavity. The large bowel had strangulated and required resection and anastomosis.

Abnormalities were found in all of the chest X-rays (Figure 1), and all included an elevated or indistinct diaphragm or pleural effusion. Computerised tomography (CT) scan confirmed a diaphragmatic injury with herniation of the abdominal contents in all cases (Figure 2).

After adequate investigations and resuscitation, all of these cases were treated surgically through an abdominal approach. The defect was localised, the abdominal contents were reduced, and the defect in the diaphragm was repaired. Post-operatively, all patients had an uneventful recovery with the exception of 2 cases of post-operative bronchopneumonia, which was managed medically.

Discussion

The diaphragm is a complex, musculotendinous, dome-shaped structure dividing the thoracic and abdominal cavities. In blunt trauma, a diaphragmatic rupture occurs when intra-abdominal forces overcome the



Figure 1: Chest X-ray (posteroanterior view) showing a cystic structure with an air-fluid level occupying the left haemithorax, causing collapse of the left lung and a slight mediastinal shift towards the right. The left haemidiaphragm is not well outlined (herniating stomach mimicking hydropneumothorax).



Figure 2: Axial noncontrast computerised tomography scan of the lower chest showing a herniated stomach in the left thoracic cavity in an 8-year-old boy (arrow).

Table 1: Age, gender, duration between the trauma and hospital admission, type of injury, clinical/radiological findings, site of the trauma, herniated organs into the thorax, associated injuries and outcome of patients.

	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case1: Female Age 5	Roadside accident (blunt). Bruises over left hypochondrium.	Hours.	Respiratory distress with vomiting. Bowel sound auscultated in chest.	Herniation of gut into left haemithorax.
	USG	Ba studies	CT scan	Operative Findings
	Minimal fluid in the abdominal cavity.	-	-	Small gut and stomach herniated through diaphragmatic rent near oesophageal hiatus.
	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case 2: Male Age 12	Roadside accident (blunt). Bruises over right iliac fossa.	Hours.	Respiratory distress with vomiting. Air entry was good on both sides.	Herniation of gut into left haemithorax.
	USG	Ba studies	CT scan	Operative Findings
	-	Herniated small gut.	-	Small gut herniated through posterolateral rent in diaphragm.
	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case 3: Male Age 6	Roadside accident (blunt). Contusion over hypogastrium and bruises over left iliac fossa.	1 year.	Respiratory distress.	Left pyopneumothorax.
	USG	Ba studies	CT scan	Operative Findings
	Tube was in the stomach. Minimal fluid in the abdominal cavity.	-	-	Spleen, small bowel, and colon herniated through posterolateral rent in diaphragm.
	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case 4: Male Age 8	Diving in pond from about 3 m height.	7 days.	Respiratory distress with hydropneumothorax.	Bilious fluid from left-sided drain.
	USG	Ba studies	CT scan	Operative Findings
	-	-	Herniated stomach and small intestine in left haemithorax.	Stomach, large gut, and spleen herniated through posterolateral rent in diaphragm.

	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case 5: Male Age 16	Stab injury spanning lower chest and upper abdomen on left side.	1 year.	Severe abdominal pain and vomiting.	Opacity in left lung base.
	USG	Ba studies	CT scan	Operative Findings
	-	-	-	Small gut and large bowel hernias with gangrene of large bowel; resection anastomosis performed.
	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case 6: Male Age 6	Roadside accident (rolled over) with bruises over whole of abdomen and back.			
	USG	Ba studies	CT scan	Operative Findings
	-	-	-	Small bowel, large bowel, and spleen hernias in left haemithorax.
	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case 7: Female Age 11	Fall from about 2.5 m; had pneumothorax. Intercostal tube was inserted initially; increased air and bilious drainage. Patient was then referred to our institute.	10 days.	Respiratory distress, bilious drainage through intercostal tube.	Pneumothorax; left side.
	USG	Ba studies	CT scan	Operative Findings
	-	-	-	Tube was in the stomach, which was herniating through posterolateral rent.

	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case 8: Female Age 15	Bullet Injury with entry wound in left hypochondrium and exit wound on the lower aspect of chest on back side.	Soon after injury.	Decreased breath sounds on left side, vomiting and respiratory distress on 3rd day of injury.	Haemopneumothorax on left side.
	USG	Ba studies	CT scan	Operative Findings
	-	-	-	Stomach herniated through posterolateral rent in the diaphragm.
	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case 9: Male Age 6	Roadside accident (rolled over) with bruises over whole of abdomen and back.	Hours.	Chest pain, abdominal pain.	Herniation of gut on left side.
	USG	Ba studies	CT scan	Operative Findings
	Haemoperitoneum.	-	-	Stomach and small intestine herniated through posterolateral rent in diaphragm.
	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case 10: Male Age 18	Roadside accident (blunt). No visible bruises over the abdomen.	Soon after injury.	Respiratory distress.	Herniation of gut into left haemithorax.
	USG	Ba studies	CT scan	Operative Findings
	-	-	-	Herniation of small gut through posterolateral rent in diaphragm.
	Mode of Injury	Duration*	Clinical Findings	Plain X-rays
Case 11: Male Age 1	Roadside accident (blunt). Bruises over left lumbar area.	Soon after injury.	Respiratory distress.	Herniation of gut into left haemithorax.
	USG	Ba studies	CT scan	Operative Findings
	-	-	-	Herniation of large gut through posterolateral rent in diaphragm.

Post-operative period was uneventful for all cases except Cases 9 and 11; both patients had bronchopneumonia.

* Duration between occurrence of trauma and hospital admission.

Abbreviations: USG = ultrasonography, Ba = barium, CT = computerised tomography.

normally higher intrathoracic pressure and the elasticity of the contracted diaphragm. The diaphragm most frequently tears at the junction of the muscular and tendinous elements that is called the centrum tendinosum (5). Herniation of the stomach, small and large intestines, kidneys, and spleen may follow the rupture acutely or years after the original injury. Delayed herniation of the abdominal viscera may occur under a number of circumstances. In some cases, the diaphragm weakens secondary to the aggressive inflammatory response at the ruptured site. Patients are also at risk just after tracheal extubation when the intrathoracic pressure quickly becomes negative. Alternatively, visceral content herniation may occur slowly, as the physiologically negative intrathoracic pressures gently pull the abdominal contents through the diaphragmatic defect (6).

Most diaphragmatic ruptures occur on the left side. This is believed to be due to a congenitally weaker left haemidiaphragm and the protective effect of the liver on the right side (5). In our study, all of the cases were left-sided.

While the classic physical signs of diaphragmatic herniation include unilateral breath sounds, a scaphoid abdomen, and bowel sounds over the lung fields, these clues are not consistently present. Many victims simply demonstrate respiratory distress as their sole pulmonary finding. Additionally, 90% to 95% of individuals with diaphragmatic ruptures have other significant injuries. The most common associated pathologies include lacerations of the spleen, liver, and kidney; pelvic fractures; major vessel disruption; long bone fractures; and head trauma (7).

Diaphragmatic injury following blunt trauma remains rare in children and may be more difficult to assess than in adults for both anatomical and physiological reasons. The compliance of the paediatric chest wall may result in internal injury in the absence of the external evidence of major injury (8). As minor injuries have caused ruptures of the diaphragm, the timing of the impact during the respiratory cycle is possibly more important than the severity of the trauma because it creates a significant pressure gradient across the diaphragm. Chest X-ray is one of the most important methods for the detection of diaphragmatic rupture and herniation. The chest X-ray, however, is diagnostic in only 25% to 50% of cases (9). Suggestive findings on the chest X-ray include an interrupted, indistinct, or elevated haemidiaphragm, bowel loops or air-fluid levels in the lung space, and a displaced nasogastric tube into the chest. Rib fractures, pneumothoraces,

haemothoraces, lower lobe collapse, and pleural effusions are associated complications that are usually evident on plain films and may increase the suspicion of a more extensive injury (9).

Other imaging modalities that are available to the emergency physician to assess the integrity of the diaphragm include ultrasonography (USG) and CT scan (7). Diaphragmatic discontinuity, diaphragmatic thickening, segmental nonrecognition of the diaphragm, intrathoracic herniation of the abdominal viscera, elevation of the diaphragm, and both haemothorax and haemoperitoneum are strong predictors of a blunt diaphragmatic rupture (10).

As diaphragmatic tears do not close spontaneously, a diaphragmatic rupture requires surgical closure (5). Laparotomy is the favoured surgical approach to acute diaphragmatic rupture, given that approximately 50% of patients with blunt diaphragmatic injuries have other intra-abdominal pathologies (9). Thoracotomy is commonly employed to repair chronic ruptures.

Successful surgical and bioprosthetic repair of TDR poses a serious challenge for surgeons. With the advent of molecular tools, proteomics, regenerative medicine and systems biology, surgeons, physicians, molecular biologists and bioengineers are now following a common translational path to reach a viable solution for successful tissue and organ reconstruction. To the best of our knowledge, there has not yet been a report of any mechanical strain-induced expression of a tendon-specific protein that leads to a visible phenotypic effect in traumatic rupture that contributes to traumatic diaphragmatic herniation. Neither is there any data available that substantiates gene or protein expression in traumatic diaphragmatic hernia or associated multi-organ complications. New strategies ranging from laparoscopic patch and intestinal sub-mucosa to stem cell and tissue-engineered bioprosthetic construct-based strategies for the repair and reconstruction of diaphragmatic hernias have been attempted in several laboratories around the world (11–15). However, most of the data for these types of studies come from congenital cases, and these remain challenging options because of high complication rates. Ongoing and future experiments in animal models for developing foetal cell-based therapies will open newer avenues towards developing a translational approach to successfully deal with traumatic diaphragm reconstruction in children.

Conclusion

TDR, though uncommon, does occur and should remain a diagnostic possibility in children because these patients do not undergo self-healing and require surgical correction of the defect. It should remain a possibility even in cases of remote trauma, as exemplified by one of our patients who presented after one year of trauma. A CT scan is the best imaging modality because it clearly depicts the anatomy with 3-dimensional reconstructions and highlights other associated organ injuries, which is extremely helpful in treating the patient. Stem cells and tissue-engineered bioprosthesis, which are being tested in several laboratories around the world, can be more relevant in such cases as they can better integrate into growing tissues.

Authors' Contributions

Conception and design, statistical expertise: DBD
Obtaining of funding, provision of study patients: KNR

Collection and assembly of data: RN

Analysis and interpretation of the data, drafting of the article: SR

Critical revision of the article: SR, DBD

Final approval of the article: KNR, SR, DBD

Administrative, technical, or logistic support: KNR, SM

Correspondence

Dr Dhara B Dhaulakhandi
PhD Cell and Molecular Biology, Otolaryngology (All India Institute of Medical Sciences)
Department of Biotechnology & Molecular Medicine
Post Graduate Institute of Medical Sciences
Pt BD Sharma University of Health Sciences
Rohtak 124 001
Haryana, India
Tel: +91-9416312237
Fax: +91-1262-211911
Email: BTMM.submissions@gmail.com

References

- Rodriguez-Morales G, Rodriguez A, Shatney CH. Acute rupture of the diaphragm in blunt trauma: Analysis of 60 patients. *J Trauma*. 1986;**26**(5):438–444.
- Fataar S, Rad FF, Schulman A. Diagnosis of diaphragmatic tears. *Br J Radiol*. 1979;**52**:375–381.
- Sharma LK, Kennedy RF, Heneghan WD. Rupture of the diaphragm resulting from blunt trauma in children. *Can J Surg*. 1977;**20**(6):553–556.
- Shackleton KL, Steward ET, Taylor AJ. Traumatic diaphragmatic injuries: Spectrum of radiographic findings. *Radiographics*. 1998;**18**(1):49–59.
- Schumpelick V, Steinau G, Schluper I, Prescher A. Surgical embryology and anatomy of the diaphragm with surgical applications. *Surg Clin North Am*. 2000;**80**(1):213–239.
- Ramos C, Koplewitz B, Babyn P, Manson PS, Ein SH. What have we learned about traumatic diaphragmatic hernias in children? *J Pediatr Surg*. 2000;**35**(4):601–604.
- Friedlaender E, Tsarouhas N. Traumatic diaphragmatic rupture in a pediatric patient: A case report. *Pediatr Emerg Care*. 2003;**19**(5):340–342.
- Brandt ML, Luks FI, Spigland NA, DiLorenzo M, Laberge JM, Ouimet A. Diaphragmatic injury in children. *J Trauma*. 1992;**32**(3):298–301.
- Rothrock S, Green S, Morgan R. Abdominal trauma in infants and children: Prompt identification and early management of serious and life-threatening injuries. Part II: Specific injuries and ED management. *Paediatr Emerg Care*. 2000;**16**(3):189–195.
- Nchimi A, Szapiro D, Ghaye B, Willems V, Khamis J, Haquet L, et al. Helical CT of blunt diaphragmatic rupture. *AJR Am J Roentgenol*. 2005;**184**(1):24–30.
- Fuchs JR, Kaviani A, Oh JT, LaVan D, Udagawa T, Jennings RW, et al. Diaphragmatic reconstruction with autologous tendon engineered from mesenchymal amniocytes. *J Pediatr Surg*. 2004;**39**(6):834–838.
- Kunisaki SM, Fuchs JR, Kaviani A, Oh JT, LaVan DA, Vacanti JP, et al. Diaphragmatic repair through fetal tissue engineering: A comparison between mesenchymal amniocyte- and myoblast-based constructs. *J Pediatr Surg*. 2006;**41**(1):34–39.
- Fauza DO, Marler JJ, Koka R, R. Forse RA, Mayer JE, Vacanti JP. Fetal tissue engineering: diaphragmatic replacement. *J Pediatr Surg*. 2001;**36**(1):146–151.
- Holcomb GW 3rd, Ostlie DJ, Miller KA. Laparoscopic patch repair of diaphragmatic hernias with Surgisis. *J Pediatr Surg*. 2005;**40**(8):E1–E5.
- Sandoval JA, Lou D, Engum SA, Fisher LM, Bouchard CM, Davis MM, et al. The whole truth: Comparative analysis of diaphragmatic hernia repair using 4-ply vs 8-ply small intestinal submucosa in a growing animal model. *J Pediatr Surg*. 2006;**41**(3):518–523.