

Editorial

'Aqualisation' of neuraxis: Wondrous neuraqua CSF 1

Manu Kothari, Atul Goel

1

View and Review

Organization of neurology services in India: Unmet needs and the way forward

Mandaville Gourie-Devi

4

Original Articles

Endoscopic management of brain abscesses

Yad Ram Yadav, Mallika Sinha, Neha, Vijay Parihar

13

Pattern of cerebellar perfusion on single photon emission computed tomography in subcortical hematoma: A clinical and computed tomography correlation

Jayantee Kalita, Usha K. Misra, Prasen Ranjan, P. K. Pradhan

17

Imaging features in Hirayama disease

Hemant A. Sonwalkar, Rakesh S. Shah, Firosh K. Khan, Arun K. Gupta, Narendra K. Bodhey, Surjith Vottath, Sukalyan Purkayastha

22

Delayed habituation in Behcet's disease

Sefa Gulturk, Melih Akyol, Hulusi Kececi, Sedat Ozcelik, Ziyet Cinar, Ayse Demirkazik

27

Erythrocyte indicators of oxidative changes in patients with graded traumatic head injury

Chandrika D. Nayak, Dinesh M. Nayak, Annaswamy Raja, Anjali Rao

31

Repeat gamma knife radiosurgery for recurrent or refractory trigeminal neuralgia

Liang Wang, Zhen-wei Zhao, Huai-zhou Qin, Wen-tao Li, Hua Zhang, Jian-hai Zong, Jian-Ping Deng, Guo-dong Gao

36

Taste dysfunction in vestibular schwannomas

Rabi Narayan Sahu, Sanjay Behari, Vimal K. Agarwal, Pramod J. Giri, Vijendra K. Jain

42

Surgical management of traumatic intracranial pseudoaneurysms: A report of 12 cases

Xiang Wang, Jin-Xiu Chen, Chao You, Min He

47

Expression of truncated dystrophin cDNAs mediated by a lentiviral vector

Sun Shunchang, Chen Haitao, Chen Weidong, He Jingbo, Peng Yunsheng

52

Gamma knife radiosurgery for glomus jugulare tumors: Therapeutic advantages of minimalism in the skull base

Manish S. Sharma, A. Gupta, S. S. Kale, D. Agrawal, A. K. Mahapatra and B. S. Sharma

57

Case Reports

Subarachnoid hemosiderin deposition after subarachnoid hemorrhage on T2*-weighted MRI correlates with the location of disturbed cerebrospinal fluid flow on computed tomography cisternography	
<i>Yoshifumi Horita, Toshio Imaizumi, Yuji Hashimoto, Jun Niwa</i>	62
Anesthesia management of awake craniotomy performed under asleep-awake-asleep technique using laryngeal mask airway: Report of two cases	
<i>Gadhinglajkar Shrinivas Vitthal, Rupa Sreedhar, Mathew Abraham</i>	65
High cervical C3-4 'disc' compression associated with basilar invagination	
<i>Atul Goel</i>	68
Short-lasting unilateral neuralgiform headache with conjunctival injection and tearing: Response to antiepileptic dual therapy	
<i>Ravi Gupta, Manjeet S. Bhatia</i>	71
Correlation of autism with temporal tubers in tuberous sclerosis complex	
<i>Kavitha Kothur, Munni Ray, Prahbjot Malhi</i>	74
Non-traumatic carotid dissection and stroke associated with anti-phospholipid antibody syndrome: Report of a case and review of the literature	
<i>Benzi M. Kluger, Richard L. Hughes, C. Alan Anderson, Kathryn L. Hassell</i>	77
Osteoma of anterior cranial fossa complicated by intracranial mucocoele with emphasis on its radiological diagnosis	
<i>Jinhu Ye, Hui Sun, Xin Li, Jianping Dai</i>	79
Vasospasm after transsphenoidal pituitary surgery: A case report and review of the literature	
<i>Manish Kumar Kasliwal, Ravinder Srivastava, Sumit Sinha, Shashank S. Kale, Bhawani S. Sharma</i>	81
Chondromyxoid fibroma of the seventh cervical vertebra	
<i>Ashish Jonathan, Vedantam Rajshekhar, Geeta Chacko</i>	84
Acute progressive midbrain hemorrhage after topical ocular cyclopentolate administration	
<i>Tarkan Calisaneller, Ozgur Ozdemir, Erkin Sonmez, Nur Altinors</i>	88

Letters to Editor

Digital subtraction angiography laboratory with inbuilt CT (DynaCT): Application during intracranial aneurysm embolization	90
Concomitant tuberculous and pyogenic cerebellar abscess in a patient with pulmonary tuberculosis	91
Drug compliance after stroke and myocardial infarction: Is complementary medicine an issue?	93

Multiple intracranial developmental venous anomalies associated with complex orbitofacial vascular malformation	93
Nitrofurantoin-induced peripheral neuropathy: A lesson to be re-learned	94
Posterior longitudinal ligament cyst as a rare cause of lumbosacral radiculopathy with positive straight leg raising test	96
Aqueductal stenosis caused by an atypical course of a deep collector vein draining bilateral cerebellar developmental venous anomalies	97
Recovery of increased signal intensity of the cervical cord on magnetic resonance imaging after surgery for spontaneous spinal epidural hematoma causing hemiparesis	98
Simultaneous thalamic and cerebellar hypertensive hemorrhages	100

Neuroimages

MRI and MRA in spontaneous intracranial arterial dissection <i>S. Raghavendra, Sanjeev V. Thomas, Krishnamoorthy Thamburaj, Bejoy Thomas</i>	102
Shunt catheter migration into pulmonary arteries <i>Miikka Korja, Matti K. Karvonen, Arto Haapanen, Reijo J. Marttila</i>	103
Susceptibility weighted imaging in holohemispheric venous angioma with cerebral hemiatrophy <i>Sivaraman Somasundaram, Chandrasekharan Kesavadas, Bejoy Thomas</i>	104

Forthcoming Events	105
--------------------------	-----

Instructions to Authors	106
-------------------------------	-----

Referees List - 2007	000???
----------------------------	--------

The copies of the journal to members of the association are sent by ordinary post. The editorial board, association or publisher will not be responsible for non-receipt of copies. If any of the members wish to receive the copies by registered post or courier, kindly contact the journal's / publisher's office. If a copy returns due to incomplete, incorrect or changed address of a member on two consecutive occasions, the names of such members will be deleted from the mailing list of the journal. Providing complete, correct and up-to-date address is the responsibility of the members. Copies are sent to subscribers and members directly from the publisher's address; it is illegal to acquire copies from any other source. If a copy is received for personal use as a member of the association/society, one cannot resale or give-away the copy for commercial or library use.

Organization of neurology services in India: Unmet needs and the way forward

Mandaville Gourie-Devi

Emeritus Professor of Neurology, Institute of Human Behavior and Allied Sciences, Chairperson of Department of Neurophysiology and Senior Consultant in Neurology, Sir Gangaram Hospital, New Delhi, India

Recognition of the magnitude of the burden and disability and mortality consequent to neurological disorders has led to global initiatives of declaring them as a “global epidemic”, emphasizing public health approach and integration of neurology care with general health care. Epidemiological transition with increase in neurological disorders in India, the gross mismatch between the need and the available trained manpower and infrastructural facilities are posing challenges to health planners and policy makers for providing neurological care to the community. Alternative approaches of optimal utilization of rural, community health, satellite clinic and district models of the health care with close interaction with tertiary centres, nongovernmental agencies and private sector may facilitate achieving the goal of taking neurology care to the ‘unreached’.

Key words: Neurological disorders, burden, prevalence, disability, health care models, neurological services

Global Scenario of burden of Neurological Disorders

The recent report published by the World Health Organization “Neurological disorders, public health challenges” states that about one billion people worldwide suffer from neurological disorders and that 6.8 million people die annually from these disorders.^[1] The spectrum of neurological disorders includes epilepsy, stroke, headache, Alzheimer’s disease and other dementias, Parkinson’s disease, multiple sclerosis, brain injuries, neuroinfections etc. Stroke is the second most common cause of mortality and a major cause of disability.

Neurological disorders accounted for 4.2% of the global burden of disease in 1996^[2] which has significantly increased to 6.29% in 2005 assessed by the disability adjusted life years (DALYs) for common neurological disorders (epilepsy, dementia, Parkinson’s disease,

multiple sclerosis, cerebrovascular disease, poliomyelitis, tetanus, meningitis, Japanese encephalitis).^[1] Further rise in burden is expected and by 2030 it is estimated to be 6.77% [Table 1]. An increase in mortality due to neurological disorders from 11.67% of the total mortality in 2005 to 12.22% in 2030 is also anticipated.^[1] Thus neurological disorders can be considered as a global epidemic calling for a global campaign to promote advocacy, improve awareness, education and prevention and to enhance neurology care.^[3]

The prevalence of neurological disorders globally in the year 2005 was 155.36 per 1000 population, nearly a third of them due to nutritional deficiency and toxins [Table 2].^[1] While dementia is estimated to increase in 2030, the decrease in prevalence rate of all neurological disorders to 143.5 can be attributed to control of infectious diseases and improved nutritional status. Based on the projections of United Nations that by 2025, there will be 1.2 billion elderly people in the world and 71% will be living in the developing countries, it is expected that neurological disorders in the elderly will show a significant rise, particularly in the developing world.

Neurological disorders pose a great challenge to healthcare in developing countries in view of limited resources and manpower that are inadequate to tackle the increasing burden. Recognizing the urgency to meet the challenges, a “Committee on Nervous System Disorders in Developing Countries” formed by the Board on Global Health, Institute of Medicine, National Academy of Sciences, USA (the author was a member of this committee) reviewed the magnitude, pattern, causes, prevention and low-cost treatment of common neurological disorders. The committee strongly recommended integration of neurological care into the public health system and emphasized that this approach is a recognized way of increasing coverage through an affordable and accessible service involving the community.^[4]

Table 1: Burden of neurological disorders, in disability adjusted life years (DALYs) in the World

Year	2005			2030		
Population	6441919466			7917115397		
Total DALYs	1469 610 066			1526745574		
Disorder	DALYs	%	Per 100000 population	DALYs	%	Per 100000 population
Epilepsy	7307975	0.50	113.44	7441536	0.49	93.99
Alzheimer and other dementias	11077525	0.75	171.96	18394267	1.20	232.34
Parkinson's disease	1616523	0.11	25.09	2015065	0.13	25.45
Multiple sclerosis	1509696	0.10	23.44	1648303	0.11	20.82
Migraine	7659687	0.52	118.90	7596089	0.50	95.95
Cerebrovascular disease	50784770	3.46	788.35	60864051	3.99	768.77
Poliomyelitis	115167	0.01	1.79	13261	0.00	0.17
Tetanus	6422611	0.44	99.70	3173636	0.21	40.09
Meningitis	5336882	0.36	82.85	2038968	0.13	25.75
Japanese encephalitis	561038	0.04	8.71	149931	0.01	1.89
Total for all neurological disorders	92391874	6.29	1434.23	103335108	6.77	1305.21

Source^[1]**Table 2: Prevalence of neurological disorders in the world**

Year	2005		2030	
Population	6441919466		7917115397	
Disorder	Number	Per 1000	Number	Per 1000
Epilepsy	39891898	6.19	50503933	6.38
Alzheimer and other dementias	24446651	3.79	44016718	5.56
Parkinson's Disease	5223897	0.81	7236712	0.91
Multiple sclerosis	2492385	0.39	3279199	0.41
Migraine	326196121	50.64	412894420	52.15
Cerebrovascular disease	61537499	9.55	76826249	9.70
Neuroinfections	18169479	2.82	13290180	1.68
Nutritional and neuropathies	352494535	54.72	285369403	36.04
Neurological injuries	170382211	26.45	242728912	30.66
Total	1000834676	155.36	1136145726	143.50

Source^[1]

Emerging Health Scenario in India

India, with a population of more than one billion, is facing new challenges in the health sector due to a paradigm shift in disease burden. Added to the burden of communicable diseases with emerging and re-emerging infections and nutritional deficiency disorders, there is epidemiological transition with increasing incidence of non-communicable disorders (NCD), some attributable to lifestyle and others to increased life expectancy.^[5] Amongst the NCDs, neurological disorders have been recognized to lead to significant mortality, morbidity, disability and socioeconomic loss. Epilepsy, cerebrovascular disorders, migraine, dementia, Parkinson's disease, motor neuron disorder, traumatic injuries, brain damage due to birth trauma, neuromuscular disorders, demyelinating disorders and neurological disorders consequent to nutritional deficiency and exposure to neurotoxic substances contribute to significant burden. Further, widely prevalent infections of the nervous system such as tuberculosis, malaria, cysticercosis and viral infections, particularly Japanese encephalitis and HIV also lead to neurological deficits.

The unique characteristics of neurological disorders such as chronicity, progressive degeneration, limited therapeutic options and lack of specific treatment for many diseases, further contribute to the disease burden and morbidity. The effect of stigma associated with epilepsy, particularly in developing countries, on disease burden cannot be easily estimated since it leads to complexities in diagnosis and management and a large treatment gap.^[6] Redeeming features are recent developments including (i) advances in diagnostic modalities of imaging, immunological and molecular tests, (ii) introduction of new therapeutic strategies such as thrombolysis, immunomodulating agents (immunoglobulins, interferon) and (iii) use of new drugs for treatment of epilepsy, Parkinson's disease and neuroinfections which have ushered a sea change in the clinical practice of neurology, enabling early diagnosis and more effective treatment leading to improved outcome of the disease.

Indian Scenario of Neurological Disorders

Population-based neuroepidemiologic surveys during the last 15 years, using standardized WHO

questionnaire with modifications,^[7-9] in different regions of the country have shown the prevalence rate of neurological disorders to vary from 967 to 4070 per 100000 population [Table 3].^[10-16] The Parsis, a distinct ethnic group, have an unusually high prevalence rate, but they do not represent the general population of the country. In these surveys, infections of the nervous system and traumatic injuries of brain, spinal cord and peripheral nerves have not been included and therefore it would be an underestimate of the total burden of neurological disorders.

All age groups are affected with peaking in the elderly above 60 years of age. In India according to the 2001 census^[17] there are 77 million people above the age of 60 years and it is expected that by 2025 there will be a huge increase to 177 million with consequent significant rise in age-related disorders such as cerebrovascular disorders, Parkinson's disease and dementia. Contradicting the perception of the policy-makers and administrators that neurological disorders are seen mostly in the urban population, neuroepidemiologic surveys have demonstrated that the prevalence in the rural population is significantly higher than in the urban population. In a large community-based survey in Bangalore of a population of 102557 comprising an urban population of 51502 and rural population of 51055, the prevalence rates were 2190 and 4070, respectively, with a ratio of 1:1.85.^[15] This fact is critical for planning infrastructure and trained manpower for providing equitable neurology care in the country.^[18] Prevalence of some common disorders is shown in Table 4. Based on the prevalence studies, it may be estimated that in India there are 20 to 30 million

people with neurological disorders and the common disorders include epilepsy (6 to 8 million), headache (10-12 million), strokes (1 to 2 million). Transient ischemic attacks are not included in surveys and hence the actual burden of cerebrovascular disorders will be higher than these projected figures.

Special mention needs to be made regarding care of people with epilepsy in view of the associated stigma, the myths and misconceptions about the nature of the disorder^[19,20] and the consequent burden to the patients and their family members, factors which are not considered while determining the disability adjusted life years (DALYS). Some of the crucial factors leading to a wide treatment gap of 38 to 80% in the country^[19,21] are stigma associated with epilepsy preventing patients seeking medical advice, lack of awareness that epilepsy is a brain disorder and that it can be treated. It is therefore important to move forward from descriptive neuroepidemiologic studies to interventional strategies for prevention and treatment of neurological disorders.^[22]

Disability Consequent to Neurological Disorders

Neurorehabilitation forms an integral component of neurology services since disability due to neurological disorders, unlike other diseases, is a major issue. The disabilities are categorized as affecting mobility, disturbance of cognition and behavior, causing pain, disturbance of consciousness and function.^[23] Further, a specific neurological disability may result from a combination of different impairments or a single

Table 3: Prevalence of total neurological disorders - Indian scenario

Author	Place	Year	Nature of population	Population surveyed	Prevalence rate/100000
Gourie-Devi ^[10]	Gowribidanur	1987	Rural	57,660	1,382
Bharucha ^[11]	Bombay	1987	Urban	851 (Parsis)	9400
Kapoor ^[12]	Ballabgarh	1989	Rural	48,798	3,487
Razdan ^[13]	Kuthar Valley	1994	Rural	63,645	967
Das ^[14]	Malda	1996	Rural	37,286	2,856
Gourie-Devi ^[9,15]	Bangalore	1996	Total	1,02,557	3,126
		2004	Rural	51,055	4,070
			Urban	51,502	2,190

Table 4: Prevalence of common neurological disorders in India (per 1000 population)

Disorder	Gourie-Devi 1987	Bharucha 1987	Kapoor 1989	Razdan 1994	Das 1996	Gourie-Devi 2004
Epilepsy	4.63	4.7	4.02	2.47	3.05	8.83
Headache	1.73	N.A	16.95	N.A	18.58	11.19
Stroke	0.52	17.6	0.88	1.43	1.26	1.50
Mental retardation and Cerebral palsy	1.63	2.4	1.09	3.3	0.64	1.42
Parkinson's disease	0.07	7.1	N.A	1.4	0.16	0.33
Peripheral neuropathy	0.52	15.2	N.A	2.99	0.75	1.28
Post poliomyelites sequelae	0.99	N.A	4.95	2.18	0.55	1.1

NA-data not available

impairment may lead to multiple disabilities. There are only a few published reports on the magnitude and severity of disability consequent to neurological disorders. There are two hospital-based studies and one community-based study. Taly and Chaudhuri^[24] in the hospital-based study at the National Institute of Mental Health and Neurosciences, Bangalore, observed that at discharge from the hospital 92% of 1093 patients had disability with varying degree of severity. Follow-up after six months showed that the disability persisted in 79% but there was a slight favorable change with increase in the proportion of those with mild disability. However, there was significant impact on the quality of life since 80% of them required assistance for daily activities. In the other hospital-based study at the Postgraduate Institute of Medical Education and Research, Chandigarh, Prabhakar *et al.*,^[25] reported that 12 months after discharge from the hospital, 60% of 258 patients had either partial or no improvement or even worsening of the disability. Assessment using Dysfunction Analysis Questionnaire showed that there was significant dysfunction in social, vocational, personal, family and cognitive domains.^[26] The dysfunction varied across different neurological disorders and was more in the elderly. In the isolated report on community survey as a component of Bangalore Urban Rural Neuroepidemiological survey (BURN), disability assessment done using modified Barthel's index showed that 20% of 3128 persons identified with neurological disorders, had significant disability and it was twice more common in the rural region compared to the urban.^[27] These studies draw attention to the need for a comprehensive rehabilitation program for care of neurologically disabled people in the 'post-acute phase' so that they can function as useful citizens as stated in the policy statement by the World Health Organization.^[28] Apart from rehabilitation focused on motor, sensory and cognitive deficits there is a need for psychosocial rehabilitation and effective utilization of community resources.^[29,30]

Economic Burden of Neurological Disorders

The economic aspects of the burden of neurological disorders include direct cost of outpatient care, drugs, emergency admissions, in-patient care and expenses for transport, particularly from remote areas to centers of healthcare and hospitals. Indirect costs are unemployment, under-employment, income lost by family members, disability-associated issues and excess mortality. Intangible costs are attributable to pain, disability and suffering. Economic burden of neurological disorders, with the exception of epilepsy, has not been determined in India. In a well-designed study by Thomas *et al.*,^[31] conducted in six medical centers, one each from six states including Andhra

Pradesh, Gujarat, Kerala, Karnataka, Maharashtra and Tamil Nadu, the annual cost of epilepsy per patient was INR 13755. Based on this data and the number of people with epilepsy in the country, the economic burden was estimated to be INR 68.75 billion. An important issue emanating from this study was that the indirect costs of travel expenses to the hospital situated at a considerable distance and the consequent loss of productivity amounted to 14% of the total annual cost. In the absence of data for other neurological disorders, it may be assumed that the indirect cost would be almost similar. Providing neurology care at the level of the community at the "doorstep" will mitigate this problem to a considerable extent.

Public Health Perspective

Conventionally, the emphasis in neurological practice has been on meticulous clinical examination, logical interpretation of the clinical profile and systematic and thorough investigations followed by complete synthesis. Focus has been on research to understand the disease process, to develop new and more precise diagnostic techniques and novel therapeutic modalities. These approaches have indeed been highly rewarding and need to be vigorously pursued to find answers to research questions. Additionally, in view of the large burden of neurological disorders, a public health approach to provide primary healthcare at the community level and develop a robust system for referral of complex problems to specialists and centers of excellence has to be put in place.^[32] Close interaction of the health sector with politicians, administrators, economists and social scientists is the backbone of a public health model. The World Health Organization (WHO) had launched the initiative of "Neurology and Public Health" in 1993 with the goals: i) to draw attention to neurological illnesses and emphasize the frequency and severity of these disorders, ii) to acknowledge neurological disorders as public health problems, iii) to emphasize the possibilities for their prevention and iv) to ensure care and treatment for people with neurological illnesses at all levels, particularly at the primary level. To achieve these objectives, the approach was to promote a national and international policy for improving neurology care, to convince the governments to consider neurological disorders as a public health problem and to increase public awareness of neurological disorders. It was reiterated that (i) prevention of stroke through national intervention programs to reduce the risk factors of hypertension, diabetes mellitus, smoking, obesity, lifestyle modification, (ii) control of infestations such as cysticercosis and (ii) reduction of birth injuries which contribute to significant proportion of epilepsy, will be possible through public health approach.^[32]

Through the collaborative effort of WHO and the World

Federation of Neurology, the status of neurology care and the lacunae in manpower and facilities across 109 countries from the six WHO regions (Africa, Americas, Eastern Mediterranean, Europe, South-East Asia and Western Pacific) was published in the Neurology Atlas.^[33] Valuable data for neurological services at the primary level, availability of specialists, beds, specialized neurology services etc. obtained through questionnaire from one key person in each country, has been included (author was the key person for India). This crucial information will be very useful for rational formulation of policies and health planning to develop the required neurological services.^[34]

Organization and Delivery of Neurology Services

The current scenario in the country is far from satisfactory with a gross mismatch of the burden of neurological disorders and the availability of trained neurologists, other essential health professionals and infrastructural facilities. The ratio of one neurologist for population varies widely among developed countries from one to two neurologists per 20000 to 100000; in developing countries the ratio is much higher at one neurologist for 3 to 5 million people and in many regions of the world there are no neurologists.^[35] In the US there is one neurologist per 26200 and in Africa it is one per 5,009,908 [Table 4].^[36,37] The scenario is grim in India with one neurologist for 3,200,000 in 1998, which has slightly improved to the current ratio of one neurologist for 1,250,000 and 800 neurologists for a population of more than one billion people [Table 5].^[38,39]

A high-powered committee was set up by Government of India, the Technology Information Forecasting and Assessment Council (TIFAC) to assess the technology and manpower requirements in various fields including health (author was a member of the subcommittee to assess the needs in neurology).^[40] Even with a modest expectation of one neurologist for 200,000, the requirement would be more than 5000 neurologists and since only 80 to 90 neurologists are trained annually in the country, half a century or more is likely to elapse before this goal is achieved.^[39] A third or more of the

neurologists are located in the metropolitan cities of Bangalore, Chennai, Delhi, Hyderabad, Kolkata and Mumbai and others in big towns but hardly any specialist care is available to the people in the rural areas. The exception to this bleak national scene is Kerala with 126 neurologists for a population of 31 million with a ratio of approximately 1:250,000 and reasonable distribution across 12 of 14 districts, with the sole exception of clustering of neurologists in the Trivandrum district (personal communication Dr. Abdul Salam Naha, President of Calicut Neurological Society). Keeping in view the constraints of specialist manpower, TIFAC recommended that there is a need to conceptualize and consider alternative strategies for organizing services at the peripheral, regional and tertiary levels [Table 6]. The crucial recommendations were to improve neurology care at all levels of the hierarchy of the health care pyramid, enhance manpower and meet the growing technology requirements. There is also an urgent need to upscale capacity-building by increasing the number of training institutes and the annual intake of candidates without compromising on the standard of training. It goes without saying that the place of neurology in the medical curriculum has to be expanded.

Models for Delivery of Neurology Care

The existing structure of the healthcare system in the country would influence the pattern of medical practice and consequently the care of people with neurological disorders. For developing models useful guidelines comprise, 1) policy statement to consider neurological disorders as a public health problem; 2) decentralization of care and development of resources and skills at the primary level; 3) close interaction between general practitioners, physician and specialist neurologist; 4) training in diagnosis and management of common and treatable neurological disorders (epilepsy, stroke, migraine, peripheral neuropathy, Parkinson's disease etc) for health providers; 5) ensuring continuity of care and adequate documentation of medical records; 6) adequate and regular supply of essential drugs; 7) establishing systems/network for referral of patients to tertiary centers for intensive and advanced diagnostic techniques and treatment; 8) partnership between governmental and nongovernmental agencies with focused goals of providing comprehensive neurology care and 9) spreading awareness and public education about neurological disorders with specific emphasis on

Table 5: Average population per neurologist

Country	Population
United States	26200
Canada	53000
United Kingdom	164000
Latin America	202000
Middle East	613000
Asia	902000
Sub-Saharan India	1540000 (711856-5009908)
1998	3200000
2002	2180000
Present position	1250000

Table 6: Neurological Services for India by the year 2020

Services	Population	Existing	Projected
Peripheral center	2,000,000	Nil	500
Regional center	10,000,000	55	100
Apex center	50,000,000	10	20

preventive aspects, particularly with reference to stroke and epilepsy.

Various models shown in Figure 1 are complementary to each other with 'bottoms up' approach and hierarchal level of expertise and facilities.^[41] A strong system of two-way communication needs to be in place with the care of the patient entrusted to the local physician after diagnostic tests and acute management has been completed and further line of treatment decided at the tertiary center.

Rural model

The prevalence of neurological disorders in the rural areas, as mentioned earlier, is almost twice that in the urban area and since 70% of India's population lives in the villages, of the estimated 20-30 million people with neurological disorders 15 to 20 million of them will be in the rural regions.^[15] Availability of neurology care at the village level would avoid the costs involved in travel, loss of wages during the period and most importantly delay in treatment. Rural models for providing neurology care with reference to epilepsy, implemented in Karnataka, West Bengal, Malawi, Kenya and Ecuador have been found to be feasible, practicable and successful in achieving the desired goals.^[42-46] This concept can be expanded to include other common neurological disorders.

Community health center model

Most of the medical colleges and medical institutions have their peripheral Community Health Centers (CHC) which provide general healthcare and undertake immunization programs covering generally a population of 50000 to 100000 and some of them closely interact with primary health centers. Many of the CHCs also have mobile health services reaching out to interior villages. Resident doctors who provide healthcare can be trained to diagnose and treat common neurological disorders and the patients requiring more detailed investigations and management can be provided care at the parent institution.^[47]

Satellite clinic model

Through a joint venture and active commitment of a tertiary institute, in this case the National Institute of

Mental Health and Neurosciences, the local government agencies at the taluk/district level and nongovernmental agencies, neurology services were provided in five peripheral centers located within 50 to 100 km of the apex center through satellite outreach clinics. The clinical expertise of consultants combined with the facilities in the peripheral hospital and the contribution of free drugs by the NGOs for common neurological disorders during monthly camps at a fixed place and on a fixed day of the week, was found to be a viable model.^[48] It was observed that common neurological disorders such as epilepsy, migraine, peripheral neuropathy, movement disorders etc could be easily managed. Duplication of this model by other institutions will help in reaching out to smaller towns and villages.

District model

There are 593 districts in the country with population ranging from 1.5 to 2 million and it is estimated that there will be approximately 45000 to 60000 people with neurological disorders in a district. A district model for healthcare delivery has the advantages that it is an independent administrative unit and the administrative head, the district health officer, has powers for planning health-related activities, implementation and monitoring health programs at the district level. Intersectorial coordination is possible and all health programs of the primary health centers and taluks are under the jurisdiction of the district. The Bellary district model for mental healthcare has been shown to be feasible for the integration of mental healthcare with general health services and became the cornerstone of the National Mental Health Plan with implementation in 25 districts and currently plans are afoot to extend the program to cover 100 districts. Based on this experience, a district model of epilepsy care was developed by the author and colleagues with the support of WHO, essentially focusing on training of district medical officers in identification, diagnosis and treatment of epilepsy.^[41,49,50] Over a two-year period during six workshops 148 district medical officers from 11 states were trained and evaluation of the training program by comparison of pre-test to post-test questionnaire showed that there was significant improvement of knowledge, skills and confidence in providing care to people with epilepsy. Based on this road map, the experience emerging from mental health and epilepsy services can be translated to provide comprehensive neurology care by suitable modifications of the training modules.

Tertiary center

Facilities for neurology care are available in about 60 medical colleges, which may be considered as regional centers. It was planned in the TIFAC^[39,40] that by 2020 there should be at least double the number of centers. Presently there are 262 medical colleges^[51] and if




Figure 1: Models for delivery of neurological care to the community

facilities for neurology care are created in at least 50% of the medical colleges, the defined goal of expanding neurology care and training of medical professionals can be achieved. Centers with advanced state-of-the-art facilities are now available in 10 apex centers. There is a need to increase these centers to at least 20 by the year 2020. All these figures pertain to the government sector only and since affordability is a major concern, the emphasis is on this sector. An increasing number of hospitals in the private sector are offering excellent care but at a prohibitive cost.

Integration and Interaction of Models of Neurology Care

It must be reiterated that the various models of neurology care suggested are not mutually exclusive but are complementary as shown in Figure 2. While common neurological disorders like migraine and epilepsy can be easily managed at the peripheral centers, cerebrovascular disorders after the initial emergency care can be referred to the tertiary center which is functionally linked and in proximity to the peripheral centers. The respective role of medical professionals and specialist neurologists may be defined as shown in Figure 2. It is envisaged that nodal neurologists at the tertiary centers, be it at the national or state level, would have a crucial responsibility of training medical professionals in addition to providing neurology care of a high order. The necessary infrastructure for providing

neurological services along with regular supply of drugs at different levels of the healthcare pyramid ranging from basic essential facilities at the primary care level to advanced state-of-the-art facilities at the tertiary centers along with appropriate manpower has to be established. Enlisting the support of nongovernmental agencies and the private sector is also an essential component of providing neurology care. This network if operationally effective and efficient, can lead to a national program for neurology care, as envisaged and enunciated by WHO and the World Federation of Neurology.^[33] On the anvil is the agenda to harness telemedicine, which currently is restricted to a few institutes in the country and exploit the full potential to enhance the clinical network between the tertiary and primary care centers.

Conclusions

It has been brought into focus that neurological disorders need to be recognized as a public health problem and advocacy be promoted to draw the attention of health planners and administrators in view of the magnitude of the problem of 30 million or more people with neurological disorders in the country. The grossly inadequate number and misdistribution of neurologists with the major proportion concentrated in metropolis and bigger towns and lack of infrastructure has deprived a large proportion of the patients, particularly the economically challenged and those in rural and remote areas, of neurology care. Alternative models of neurology care drawing upon the rich experience of healthcare delivery through rural community health centers, satellite clinics and district models with close interaction with the specialist neurologists in tertiary institutes, having a pivotal role in training the medical professionals in the peripheral centers, form the backbone of national program for neurology care in the country. Concentrated action in this direction is an urgent necessity.

Foot note

The author is currently Member of the World Health Organization Expert Advisory panel on Neurosciences. Earlier served as (i) Convener, Sub-group on Neurological, Neurosurgical, Psychiatric Disorders and Addictions, Technological Information, Forecasting and Assessment Council (TIFAC) for 2020, Department of Science and Technology, New Delhi; (ii) Member of Committee on “Neurology in Public Health”, World Health Organization; (iii) Member of Committee on Nervous System Disorders in Developing Countries” Board on Global Health, Institute of Medicine, National Academy of Sciences, USA and (iv) Resource person for India for development of “Atlas: Country resources for neurological disorders” World Health Organization and World Federation of Neurology.




Figure 2: Integration of neurology care at various levels of health care

References

- World Health Organisation. Neurological disorders: public health challenges. Geneva; 2006.
- Murray CJ, Lopez AD, editors. The global burden of disease: A comprehensive assessment of mortality and disability from diseases, injuries and risk factors in 1990 and projected to 2020. Global burden of disease and injury series, vol.1. Cambridge (MA): Harvard School of Public Health on behalf of the World Health Organization and the World Bank; 1996.
- Bergen DC, Silberberg D. Nervous system disorders. Arch Neurol 2002;59:1194-6.
- Jablensky A, Johnson R, Bunney W Jr, Cruz M, Durkin M, Familusi J, et al. Neurological, psychiatry and developmental disorders: Committee on nervous system disorders in developing countries (Board on Global Health, Institute of Medicine). National Academy Press: Washington DC; 2001.
- Reddy KS, Shah B, Verghese C, Ramdoss A. Responding to the threat of chronic diseases in India. Lancet 2005;366:1744-9.
- Meinardi H, Scott RA, Reis R, Sander JW; ILAE Commission on the developing world. The treatment gap in epilepsy: The current situation and ways forward. Epilepsia 2001;42:136-49.
- Gourie-Devi M, Rao VN, Prakash R. A protocol to detect neurological disorders in the community. Indian J Med Res 1988;88:443-9.
- Gourie-Devi M, Gururaj G, Satishchandra P. Neuroepidemiology in developing countries. Manual for descriptive studies. Bangalore: National Institute of Mental Health and Neuro Sciences (NIMHANS Publication No.33); 1994.
- Gourie-Devi M, Gururaj G, Satishchandra P, Subbakrishna DK. Neuroepidemiological pilot survey of an urban population in a developing country: A study in Bangalore, South India. Neuroepidemiology 1996;15:313-20.
- Gourie-Devi M, Rao VN, Prakash R. Neuroepidemiological study in semi-urban and rural areas in South India: Pattern of neurological disorders including motor neurone disease. In: Gourie-Devi M, editor. Motor neurone disease: Global clinical patterns and international research. Oxford and IBH: New Delhi; 1987. p. 11-21.
- Bharucha NE, Bharucha EP, Dastur HD, Schoenberg BS. Pilot survey of the prevalence of neurological disorders in the Parsi community of Bombay. Am J Prev Med 1987;3:293-9.
- Kapoor SK, Banerjee AK. Prevalence of common neurological diseases in a rural community of India. Indian J Com Med 1989;14:171-6.
- Razdan S, Kaul RH, Motta A, Kaul S, Bhatt RK. Prevalence and pattern of major neurological disorders in rural Kashmir (India) in 1986. Neuroepidemiology 1994;13:113-9.
- Das SK, Sanyal K. Neuroepidemiology of major neurologic disorders in rural Bengal. Neurol India 1996;44:47-58.
- Gourie-Devi M, Gururaj G, Satishchandra P, Subbakrishna DK. Prevalence of neurological disorders in Bangalore, India: A community-based study with a comparison between urban and rural areas. Neuroepidemiology 2004;23:261-8.
- Gourie-Devi M, Gururaj G, Satishchandra P. Neuroepidemiology in India: Development during three decades. NIMHANS J 1999;17:379-87.
- Census of India- Census data; National summary data. 2001. Available from: <http://www.censusindia.gov.in>.
- Gourie-Devi M, Gururaj G, Satishchandra P. Neuroepidemiology: Present insights and future prospects. NIMHANS J 1999;17:423-37.
- Radhakrishnan K, Pandian JD, Santoshkumar T, Thomas SV, Deetha TD, Sarma PS, et al. Prevalence, knowledge, attitude and practice of epilepsy in Kerala, South India. Epilepsia 2000;41:1027-35.
- Gourie-Devi M, Anderson DW, Rao VN. Neuroepidemiologic survey in a developing country: Some questions and answers. In: Anderson DW, editor. Neuroepidemiology: A tribute to Bruce Schoenberg. CRC Press: Boston; 1991. p. 13-23.
- Gururaj G, Reddy GN, Shariff IA, Subbakrishna DK. Factors associated with utilization of services by epilepsy patients in rural areas: A descriptive study. NIMHANS J 1994;12:135-40.
- Gourie-Devi M, Gururaj G, Satishchandra P. Interventional neuroepidemiology: From descriptive studies to interventions. In: Venkataraman S, editor. Progress in Clinical Neurosciences. Neurological Society of India: New Delhi; 1999. p. 225-41.
- Hewer RL. The epidemiology of disabling neurological conditions. In: Greenwood R, Barnes MP, McMillan RM, Ward CD, editors. Neurological rehabilitation. Churchill Livingstone: New York; 1993. p. 3-12.
- Taly AB, Chaudhuri JR. Neurorehabilitation: An emerging concept. In: Venkataraman S, editor. Progress in clinical neurosciences. Neurological Society of India: New Delhi; 1997. p. 427-37.
- Prabhakar S, Chopra JS, Pershad D. Physical disability and psychosocial functioning in neurological disorders: A follow up study. Neurol India 1991;39:63-6.
- Pershad D, Verma SK, Malhotra S, Malhotra S, Das K, Khan HA. Further validation of the dysfunction analysis questionnaire. Indian J Clin Psychol 1983;10:445-50.
- Gururaj G, Gourie-Devi M, Taly AB, Satishchandra. Epidemiology of neurological disorders and consequent disabilities-Indian Scenario: Implications for rehabilitation policies and programmes. In: Taly AB, Nair KPS, Murali T, editors. Neurorehabilitation principles and practice. National Institute of Mental Health and Neuro Sciences: Bangalore; 1998. p. 1-35.
- World Health Organisation Technical Report Series, 744. Hospital and Health for all. Geneva; 1987. p. 87-8.
- Prakash R, Gourie-Devi M. Scope of social work services in neurological setting: Initial observations. J Med Psy Social Work 1987;1:15-20.
- Prakash R, Gourie-Devi M. Rehabilitation of neurologically disabled persons, Implications for community participation. Rehabilitation Asia 1988;29:8-12.
- Thomas SV, Sarma PS, Alexander M, Pandit L, Shekhar L, Trivedi C, et al. Economic burden of epilepsy in India. Epilepsia 2001;42:1052-60.
- Leonardi M. Neurology and public health in developing countries. In: Clifford RF, editor. Recent advances in tropical neurology. Elsevier Science Publishers BV: Amsterdam, the Netherlands; 1995. p. 1-6.
- World Health Organization, World Federation of Neurology. Atlas: Country resources for neurological disorders. Geneva; 2004.
- Janca A, Aarli JA, Prilpko L, Dua T, Saxena S, Saraceno B. WHO/WFN survey of neurological services: A worldwide perspective. J Neuro Sci 2006;247:29-34.
- Aarli JA. Neurology and WHO: Challenges and issues-A report. World Neurol 2002;17:10-1.
- Bergen DC. World Federation of Neurology Task Force on Neurological Services: Training and distribution of neurologists worldwide. J Neuro Sci 2002;198:3-7.
- Bower JH, Zeneba G. Neurologic services in the nations of Africa. Neurology 2005;64:412-5.
- Singhal BS. Neurology in developing countries: A population perspective. Arch Neurol 1998;55:1019-21.
- Gourie-Devi M. Medical technologies for neurological sciences. In: Mathew L, editor. Technology management in health care. Defence bioengineering and electromedical laboratory (DEBEL): Bangalore; 1998. p. 62-7.
- Technology Information Forecasting and Assessment Council (TIFAC) report on Health care, technology vision 2020. Department of Science and Technology: New Delhi; 1996.
- Gourie-Devi M, Satishchandra P, Gururaj G. Delivery of epilepsy care to the community: Towards the national epilepsy control programme. In: Agarwal SP, Goel DS, Ichhpujani RL, Salhan RN, Shrivastava S, editors. Mental health. An Indian perspective 1946-2003. Directorate General of Health Services Ministry of Health and Family Welfare: New Delhi; 2004. p. 295-305.
- Mani KS, Rangan G, Srinivas HV, Sridharan VS, Subbakrishna DK. Epilepsy control with phenobarbital or phenytoin in rural south India: The Yelandur study. Lancet 2001;357:1316-20.
- Pal DK, Das T, Chaudhury G, Johnson AL, Neville BG. Randomised controlled trial to assess acceptability of phenobarbital for childhood epilepsy in rural India. Lancet 1998;351:19-23.
- Watts AE. A model for managing epilepsy in rural community in Africa. BMJ 1989;298:805-7.
- Feksi AT, Kaamugisha J, Sanders JW, Gatiti S, Shorvon SD. Comprehensive primary healthcare: Anti-epileptic drug treatment programme in rural and semi urban Kenya. Lancet 1991;337:406-9.
- Placencia M, Sander JW, Shorvon SD, Paredes V, Suarez J, Cascante SM. Anti-epileptic drug treatment in a community healthcare setting in Northern Ecuador: A prospective 12-month assessment. Epilepsy Res

- 1993;14:237-44.
47. Sriram TG, Chandrasekhar CR, Moily S. Epilepsy in primary care: A study of follow-up profile and response to treatment. *Nimhans Jr* 1998;8:133-7.
48. Reddy GN, Channabasavanna SM, Gourie-Devi M, Das BS, Prabhu GG, Shariff IA, *et al.* Extension of the mental health services by satellite clinics as a model. *NIMHANS Jr* 1986;4:71-5.
49. Gourie-Devi M, Satishchandra P, Gururaj G. National workshop on public health aspects of epilepsy for senior personnel of state health departments in India. *Ann Indian Acad Neurol* 1999;2:43-8.
50. Gourie-Devi M, Satishchandra P, Gururaj G. Epilepsy in primary care: Epilepsy control program in India: A district model. *Epilepsia* 2003;44:58-62.
51. Central Bureau of health intelligence, 2007. Available from: <http://www.cbhidghs.nic.in>.

Accepted on 12-01-2008

Source of Support: Nil, **Conflict of Interest:** None declared.