

Detection of *Wolbachia pipientis*, including a new strain containing the *wsp* gene, in two sister species of *Paraphlebotomus* sandflies, potential vectors of zoonotic cutaneous leishmaniasis

Parviz Parvizi^{1/+}, Ali Bordbar¹, Narmin Najafzadeh^{1,2}

¹Molecular Systematics Laboratory, Parasitology Department, Pasteur Institute of Iran, Tehran, Iran

²Department of Microbiology, Science and Research Branch of Fars, Islamic Azad University, Tehran, Iran

Individual, naturally occurring Phlebotomus mongolensis and Phlebotomus caucasicus from Iran were screened for infections with the maternally inherited intracellular Rickettsia-like bacterium Wolbachia pipientis via targeting a major surface protein gene (wsp). The main objective of this study was to determine if W. pipientis could be detected in these species. The sandflies were screened using polymerase chain reaction to amplify a fragment of the Wolbachia surface protein gene. The obtained sequences were edited and aligned with database sequences to identify W. pipientis haplotypes. Two strains of Wolbachia were found. Strain Turk 54 (accession EU780683) is widespread and has previously been reported in Phlebotomus papatasi and other insects. Strain Turk 07 (accession KC576916) is a novel strain, found for first time in the two sister species. A-group strains of W. pipientis occur throughout much of the habitat of these sandflies. It is possible that Wolbachia is transferred via horizontal transmission. Horizontal transfer could shed light on sandfly control because Wolbachia is believed to drive a deleterious gene into sandflies that reduces their natural population density. With regard to our findings in this study, we can conclude that one species of sandfly can be infected with different Wolbachia strains and that different species of sandflies can be infected with a common strain.

Key words: *Wolbachia pipientis* - *wsp* gene - *Paraphlebotomus* - *L. major* - Iran

The prevalence of zoonotic cutaneous leishmaniasis (ZCL), caused by *Leishmania major*, is increasing in many parts of Iran. ZCL originates as a disease of gerbils and *Leishmania* parasites are transmitted by sandflies that live and breed in gerbil burrows (Mohebali et al. 2004, Mirzaei et al. 2011). This form of leishmaniasis is of great public health importance in rural areas of 15 out of 32 provinces of Iran (Yaghoobi-Ershadi et al. 2005, Parvizi et al. 2010a). The disease is generally restricted to areas that are heavily infested by sandflies, which are the vectors of ZCL. The Turkmen Sahara region is one of the known focal areas of ZCL in Iran and includes two important ZCL foci (Gonbad Kavous and Maraveh Tapeh) in rural areas. *Phlebotomus papatasi* (Diptera: Psychodidae) is the main vector and *Paraphlebotomus* species (*Paraphlebotomus mongolensis* and *Paraphlebotomus caucasicus*) are secondary principal vectors of ZCL in this district (Nadim & Faghih 1968, Yaghoobi-Ershadi et al. 1996, Parvizi & Ready 2008).

Wolbachia pipientis is a maternally inherited endoparasitic bacterium belonging to the α -proteobacteria family that infects 20-75% of all insect species, including sandflies (Werren et al. 1995, 2008, West et al. 1998,

Jeyaprasakash & Hoy 2000, Werren & Windsor 2000, Hilgenboecker et al. 2008, Azpurua et al. 2010).

This bacterium has attracted attention because it induces a number of intriguing abnormalities in the host's reproductive system (O'Neill et al. 1997, Stouthamer et al. 1999). These intracellular microorganisms affect the biology of their invertebrate hosts in many ways, ranging from mutualistic effects to the establishment of reproductive isolation and thus, speciation (Werren 1997, Bordenstein et al. 2001, Telschow et al. 2005, Werren et al. 2008).

A wide range of phenotypes are expressed under *W. pipientis* infection, associated with processes ranging from evolution to speciation (Breeuwer & Werren 1990, Hurst & Schilthuizen 1998) to mechanisms involved in the genetic modification and biological control of pest arthropods (Sinkins et al. 1997, Bourtzis & O'Neill 1998), but these bacteria are commonly considered to be reproductive parasites (Moran et al. 2008, Werren et al. 2008).

It has been noted that most *Wolbachia*-induced phenotypes increase the proportion of females among the offspring of infected females and pose unique threats to host populations (Werren et al. 2008).

The phenotypes associated with these threats include (i) induction of parthenogenesis in the host, (ii) transformation of males into functional daughters (feminisation), (iii) provoking the selective abortion of sons of infected mothers (male killing) and (iv) cytoplasmic incompatibility (CI), which is the most common phenotype occurring in the mosquito (Hoffmann & Turelli 1997, Werren et al. 2008). The reproductive distortions caused by *Wolbachia* infection are known to result in embryonic death and subsequent egg hatching failure due to disruptions during

doi: 10.1590/0074-0276108042013004

Financial support: Pasteur Institute of Iran (367) (to PP)

+ Corresponding author: parp@pasteur.ac.ir

Received 6 September 2012

Accepted 24 January 2013

the early events of fertilisation. Thus, a *Wolbachia* strain can promote its rapid vertical transmission throughout a population of the *Phlebotomus* genus (Benlarbi & Ready 2003). *Wolbachia* are vertically transmitted from mothers to daughters, but can occasionally be horizontally transmitted between individuals, including individuals belonging to different species (Pascal et al. 2006).

Wolbachia display tropism for the reproductive tissue of their hosts and are transmitted vertically from insect to insect through the ovules. Interspecific transmission appears to occur horizontally, with the possible aid of parasitoids (Stouthamer et al. 1999, Charlat et al. 2002, Kondo et al. 2002, Ahrens & Shoemaker 2005, Baldo et al. 2005, Espino et al. 2009).

The *Wolbachia* surface protein (*wsp*) is commonly used as a marker for strain typing (Schaub et al. 1989, Van Meer et al. 1999, Pintureau et al. 2000, Shoemaker et al. 2002, Kyei-Poku et al. 2005) (Fig. 1) and a strain typing system utilising *wsp* has been developed (Zhou et al. 1998).

The *wsp* gene is analogous to the antigens employed for serotyping pathogenic bacteria (Stouthamer et al. 1999, Baldo et al. 2006). It is approximately 10 times more variable in its DNA sequence than 16S rRNA (Zhou et al. 1998).

Traditionally, *Wolbachia* spp detected in arthropods have been divided in two groups (A and B) based on the sequences of their 16S rRNA, *ftsZ* and *wsp* genes (Werren et al. 1995, Zhou et al. 1998). Both groups contain *Wolbachia* spp that have been detected in several genera of sandflies. Group A includes the *Wolbachia* spp detected in *Sergentomyia* and from *Phlebotomus*. Group B contains the *Wolbachia* spp detected in sandflies belonging to the *Phlebotomus* and *Lutzomyia* genera (Werren et al. 1995, Zhou et al. 1998, Ono et al. 2001).

These bacteria have been detected using molecular tools and reported as “*Wolbachia* species” from both arthropod and nematode hosts and only a single species has thus far been properly isolated and bears a valid name, *W. pipientis* (Lo et al. 2002).

Wolbachia may have an influential effect on the control of disease vectors based on its potential role in reducing the ability of the host to reproduce, thus decreasing its population size (Beard et al. 1993). Therefore, it is worthwhile to investigate the occurrence and distribution of *Wolbachia* in sandflies in nature.

The first objective of this study was to determine if *W. pipientis* could be detected in two sister species of *Paraphlebotomus* sandflies in which the bacterium has not been previously reported. In these two sandfly spe-

cies, *P. mongolensis* and *P. caucasicus*, the external genitalia of males provide the only diagnostic morphological characteristics. The females of these sister species cannot be separated morphologically or molecularly based on the structure of the spermathecae, the weakly developed pharyngeal armature or the mitochondrial cytochrome *b* gene (Theodor & Mesghali 1964, Killick-Kendrick 1999, Essegir et al. 2000, Parvizi et al. 2010a, b). The second objective of this study was to improve our knowledge regarding the detection of one or more strains of *W. pipientis* in one or both sister species of sandfly. Its third objective was to determine if sandflies can be simultaneously infected by two different strains of *W. pipientis*. Finally, the fourth objective of this study was to demonstrate the presence of *Wolbachia* in *Paraphlebotomus* sandflies, which would have a significant impact in assessing the diversity of *Wolbachia* strains, particularly in *L. major* vectors.

MATERIALS AND METHODS

Origin and identification of sandflies - The Turkmen Sahara villages in northeastern Iran are a focus of ZCL, where *L. major* parasites are the major causative agent of ZCL. Two sister *Paraphlebotomus* species were collected regularly from 18 villages of Gonbad Kavous and Maraveh Tapeh in late June and the middle of September in 2008 and 2009. There were also collections performed in September 2010 during peak sandfly activity in the transmission seasons of two successive years (Table I). Additionally, sticky paper and CDC traps were used to sample sandflies in the ruins of mud-walled outhouses adjacent to animal shelters, around houses, at the entrance of gerbil burrows and in houses surrounded by gerbil burrows at the edge of rural areas.

The specimens were identified as *P. mongolensis* males, *P. caucasicus* males or *P. mongolensis/P. caucasicus* females, based on morphological characters of the head and the abdominal terminalia (Lewis 1982).

DNA extraction, gene amplification and sequencing - DNA was extracted from the thorax and attached anterior abdomen of each sandfly. The samples were screened for the presence of *W. pipientis* using the general (or non-strain specific) primer pair *wsp* 81F and 691R (Zhou et al. 1998) (Fig. 1). Polymerase chain reaction (PCR) amplification was carried out according to the protocol of Benlarbi and Ready (2003).

Each 20 µL PCR mixture consisted of 1X Promega buffer B, 25 mM MgCl₂, 200 µM each dNTP, 0.4 µM each primer, one unit of Taq DNA polymerase (Promega) and 2 µL of sandfly genomic DNA. PCR amplification was carried out as follows: 2 min denaturation at 94°C, followed by 35 cycles of denaturation at 94°C for 30 sec, annealing at 50°C for 30 sec and extension at 72°C for 1 min 30 sec, with a final extension at 72°C for 10 min.

Following amplification, the samples were fractionated via horizontal submerged gel electrophoresis using 1.5% agarose gels with DNA size markers (Promega PCR markers G316A or Bioline Hyperladder IV). The obtained DNA fragments were visualised by ethidium bromide staining and then excised and purified using




Fig. 1: schematic presentation of the *Wolbachia* surface protein (*wsp*) gene, an appropriate candidate as an antigenic serotyping for strain typing of pathogenic bacteria of *Wolbachia*.

the GeneClean II Kit (BIO 101 Inc), prior to sequencing each strand (Testa et al. 2002). The sequences were then edited and aligned with database sequences using Sequencher™ version 4.4 software (Gene Codes Corp) to identify unique sequences (= haplotypes), which were analysed phylogenetically using MEGA software.

RESULTS

P. mongolensis and *P. causicus* were identified based on morphological characteristics of their external genitalia. A total of 203 *P. mongolensis* and *P. causicus* sandflies were trapped at two sites in Turkmen Sahara in the province of Golestan.

The wild sandflies were identified as belonging to two sister species of the *Paraphlebotomus* subgenus. In total, 74 out of the 203 sandflies examined were infected with *Wolbachia*. The *Wolbachia wsp* gene was detected for the first time in all three groups and was distributed as follows: in 33/96 males of *P. mongolensis*, 10/15 males of *P. causicus* and 31/92 females of *P. mongolensis/P. causicus* (31/92), corresponding to infection rates of 34.37%, 66.66% and 33.69%, respectively (Table I).

We attempted sequencing in all 74 sandflies with *Wolbachia* infections. However, only 25 of the sandflies presented sufficient DNA for sequencing and/or showed readable sequences demonstrating the presence of *Wolbachia* strains.

Two strains of *Wolbachia* were found in *P. mongolensis* and *P. causicus* sandflies (Fig. 2). The common strain, Turk 54 (accession EU780683) and a novel strain, Turk 07 (accession KC576916), were found for first time in *P. mongolensis* and *P. causicus* sandflies from

Turkmen Sahara. Following phylogenetic analyses, these strains appeared to belong to the A-group. A total of 19 of the *P. mongolensis* and *P. causicus* sandflies exhibited the Turk 54 strain haplotype, while six of the sandflies were infected with strain Turk 07 (Fig. 3, Table II).

```
Turk 07  AGCTACTACA TTGTTTGA ATACAGGT GAATTTTAC CAGAGTAT TAGATCC:A TTAAC:CA TCTT:TATA 80
Turk 54  AGCTACTAGC TTGTTTGA ATACAGGT GAATTTTAC GAGACTGA GATAGCTCA CAATCTGA CTTTATAA
      *
Turk 07  GCTGTGTGTT GTGCTTTGG T:TACAAA::A::T::G::GAGACAT CAGGTGTAT GTTGAAGAG TTTATTCATA 160
Turk 54  GCTTCTTTTA TGCTGTGTTG TGCTGCTATT GTTTATAAAA GAGAGACAT CAGGTGTAT GTTGAAGAGC TTTATTCATA
      * * * * *
Turk 07  CCTAAACAAA AATGATGTTA AGATGTAGT ATTACCCCA GCAGATACTA TTGCGAAGC TTTACAGCA ATTCAGGAC 240
Turk 54  GCTAAGCAAG GATGACTTT::GCTGTAGC TCTACTGCA GCAATGCA::G::ACAG TTTACAGCA ATTCAGGAC
      * * * * *
Turk 07  TAGTTAAGTT TTATTAGAT ATAGCAATTG AGATATGCT TATCACTCA TATGTTGTTG TTGTTGTTG TGCAGGATAT 320
Turk 54  TAGTTAAGTT TTATTAGAT ATAGCAATTG AGATATGCT TATCACTCA TATGTTGTTG TTGTTGTTG TGCAGGATAT
      *
Turk 07  ATTAGCACAC CTTTGCAAC GCTG:TG:A ATAG:TCAAA ATGTAATTT TCTTTTCTT GGTCAAGTAA AGCTGATGT 400
Turk 54  ATTAGCACAC CTTTGCAAC TCTG:TG:A GTAG:TCAAA ATGTAATTT TCTTTTCTT GGTCAAGTAA AGCTGATGT
      *
Turk 07  TAGCTATGAT GTAACCTGAG AATCAAACT CTATGCGGTT GCTGTTATT TTGTTCTTA TGTTGCTCAT TTGAT::A: 480
Turk 54  TAGTTATGAT GTAACCTGAG AATCAAACT CTATGCGGTT GCTGTTATT TTGTTCTTA TGTTGCTCAT TTGAT::A:
      *
Turk 07  GGA::GA:AA A:A::GTAGA TGAACCAAA AGGTTGCTG ATAAAGGCGC ATATAAG 538
Turk 54  AACTGACAAA AGATGGCAAA GGAACCTCA AGGTT::CT: TTACA:GCAC :TGT:TGG
      * * * * *
```

Fig. 2: alignments of the two *Wolbachia* surface protein gene sequence of *Wolbachia pipientis* isolated from Iranian sandflies strain Turk 54 isolated first from *Phlebotomus papatasi* and then most different sandfly species with the GenBank sequence EU780683, Turk 07 isolated from *Paraphlebotomus mongolensis* and *Paraphlebotomus causicus*. Nucleotide differences are marked by a star.

TABLE I
Paraphlebotomus causicus and *Paraphlebotomus mongolensis*
detected *Wolbachia* surface protein (*wsp*) gene in different villages in two locations, Turkmen Sahara, Iran

Genus Subgenus	Location	Gonbad Kavous							Maraveh Tapeh							Total <i>wsp</i> gene Ve per species n (%)	Total <i>wsp</i> gene +Ve n (%)
	Species	Villages	Dashlibroun	Shourdegesh	Kheyrikhajeolia	Inchebroun	Dozalum Fadavi	Daneshmand	Okhitapeh	W-Gharagol	E-Gharagol	Makhtumgholi	Souzes	Khajekaldi	Jafarbay		
<i>Phlebotomus</i> <i>Paraphlebotomus</i>	<i>P. mongolensis</i> (M)	4	3	1	2	1	15	4	0	0	1	1	0	1	33/96 (34.37)	33/203 (16.26)	
	<i>P. causicus</i> (M)	2	0	0	1	1	2	2	0	0	1	0	1	0	10/15 (66.66)	10/203 (4.92)	
	<i>P. mongolensis</i> and <i>P. causicus</i> (F)	6	2	2	5	1	2	1	2	1	2	3	1	3	31/92 (33.69)	31/203 (15.27)	
Total		12	5	3	8	3	19	7	2	1	4	4	2	4	74/203		
(%)		5.91	2.46	1.48	3.94	1.48	9.35	3.45	0.99	0.49	1.97	1.97	0.99	1.97	(36.45)		
		28.08							8.37								

F: female; M: male; +ve: *Wolbachia* positive.

All of the *wsp* gene fragments amplified from *P. mongolensis* and *P. caucasicus* were sequenced directly. The 564-bp Turk 54 haplotype (minus primers) (GenBank accession EU780683) was indistinguishable from that of the A-group strain of *W. pipientis* (wPap), which was previously isolated from *P. papatasi* originating from Israel/West Bank (AF237883) (Ono et al. 2001), India (GenBank accession AF237882) (Ono et al. 2001), Spain and Iran (Benlarbi & Ready 2003, Parvizi et al. 2003). All of these sandflies from Spain and Iran were wild caught, as was the pool of sandflies possessing the AF237883 sequence. The other sequences were isolated from *P. papatasi* bred in laboratory colonies. Using the same *wsp* gene primers, Cui et al. (1999) amplified a fragment of approximately 600 bp from a colony of *P. papatasi* originating from Israel/West Bank, North Sinai in Egypt and Saudi Arabia. However, these authors did not report any sequence data. Haplotype wPap was originally reported as presenting an ambiguous base (C/T) at nucleotide position 102 (GenBank accession AF020082) (Zhou et al. 1998).

The new Turk 07 haplotype differed in a number of point mutations and insertion-deletion events from the previously described haplotypes (Fig 2, Table II). The new sequence was aligned with the available *wsp* sequences of *Wolbachia* from sandflies and other insects and phylogenetic relationships were inferred. The

phylogram generated via neighbour-joining analysis placed the sequences from *P. papatasi* on the same terminal branch within the A-group of strains of *W. pipiens* (Fig. 3).

DISCUSSION

Here, we report the detection of *Wolbachia* in two potential vectors of ZCL (*P. mongolensis* and *P. caucasicus*) for the first time. *Wolbachia* had previously been isolated from various insects including sandflies, but infection of *P. mongolensis* and *P. caucasicus* had not been detected. A new *W. pipientis* strain (Turk 07) was found in both *P. mongolensis* and *P. caucasicus*. Additionally, the widespread common strain (Turk 54) was found in these two sister species.

The infection of the two *Paraphlebotomus* species with the same strain of *Wolbachia* may have been due to recent horizontal transmission of *Wolbachia* across host species. Alternatively, in the case of the closely related species *P. mongolensis* and *P. caucasicus*, infection may be due to co-divergence and co-evolution of *Wolbachia* strains and their hosts.

Another intriguing finding of this study was that one *Paraphlebotomus* species can be infected with different *Wolbachia* strains.

The *wsp* gene was used for phylogenetic analysis of the *Wolbachia* strains from which the sequences originated based on a previous study that detected *Wolbachia* in sandflies with the *wsp* gene (Werren et al. 1995, Zhou et al. 1998, Ono et al. 2001, Wu & Hov 2012).

The *wsp* sequences obtained from the *P. mongolensis* and *P. caucasicus* Turk 54 strains showed a high similarity to sequences deposited in GenBank as “incompatibility symbiont of *P. papatasi*” (Fig. 3, Table II).

However, the Turk 54 haplotype *wsp* sequences isolated from *Paraphlebotomus* were identical to a sequence previously reported from *P. papatasi* from Tunisia and Iran (Benlarbi & Ready 2003). In contrast, the Turk 07 sequences obtained from *P. mongolensis* and *P. caucasicus* were unique.

Phylogenetic studies using 16S rDNA have established that strains of *W. pipientis* from highly divergent hosts form a monophyletic clade related to the *Ehrlichia* assemblage within the α -subdivision of proteobacteria (O'Neill et al. 1992, Rousset & Solignac 1995). The examination of 16S rDNA sequences is a conservative approach that has allowed not only the phylogenetic placement of bacterial species, but also the phylogenetic resolution of *W. pipientis* into two strain groups: A and B (O'Neill et al. 1992). More recently, the more rapidly evolving *wsp* gene has been used to improve the phylogenetic resolution within the *W. pipientis* species clade, which was divided into four groups (A-D) and 12 subgroups (Zhou et al. 1998, Ono et al. 2001). The A and B groups are consistent with those identified via 16S rDNA for *W. pipientis* strains from insects, mites and crustaceans, whereas groups C and D are consistent with the strains from filarial nematodes.

Wolbachia is normally maintained in nature through vertical transmission (Werren 1997). Although horizontal transfer appears to be infrequent within host populations, it has previously occurred between two host sisters




Fig. 3: unrooted neighbour-joining tree showing the relationships of the haplotypes of the *Wolbachia* surface protein gene fragment for the isolates of *Wolbachia* listed in Table II.

species. This demonstrates that the phylogeny of *Wolbachia* strains differs noticeably from that of their hosts (O'Neill et al. 1992, Werren et al. 1995). This conclusion is supported by experiments revealing the transfer of symbionts between host individuals (Huigens et al. 2000).

The demonstrated horizontal transmission of *Wolbachia* could shed light on sandfly control because *Wolbachia* is believed to drive a deleterious gene into mosquitoes that reduce their natural population density (Beard et al. 1993, Sinkins et al. 1997).

The results of a detailed phylogenetic analysis including characterisation of the limitations of such an approach could serve as a foundation for understanding the evolution of *Wolbachia* bacteria.

Because genetically similar strains are often found in similar insect hosts, ecological interactions among hosts may mediate horizontal transfers. In particular, it appears that the horizontal transmission of *Wolbachia* has occurred quite recently and frequently within the genus *Phlebotomus*.

TABLE II
Wolbachia strains isolated from different hosts in various geographical origins and Iran

Host	Strain	Geographic origin	GenBank accession	<i>Wolbachia</i> surface protein gene fragment (bp)
<i>Drosophilidae</i> sp.	<i>A_PanPNM_Droso12b</i>	Panama	EU126468	513
<i>Drosophila melanogaster</i>	<i>wMel</i>	China	EU395833	632
<i>Phlebotomus papatasi</i>	<i>wPhleb3lf</i>	France	JN315984	568
<i>D. melanogaster</i>	<i>wMel</i>	Wuhan	FJ403330	638
<i>D. melanogaster</i>	<i>wMel</i>	Yunnan	FJ403332	637
<i>D. melanogaster</i>	-	Ukraine	HM775088	600
<i>D. melanogaster</i>	-	India	JF505383	590
<i>Aphis fabae</i>	<i>wsp_GRA4</i>	Greece	JN384045	620
<i>Phlebotomus duboscqi</i>	<i>wPhlebM4</i>	France	JN315983	524
<i>Paraphlebotomus mongolensis</i>	(Turk 07)	Iran	KC576916	551
<i>Paraphlebotomus caucasicus</i>				
<i>Calyptrate muscoid</i> fly	<i>A_MexSon_Calyp150671</i>	Mexico	EU126482	486
<i>P. papatasi</i>	<i>Wpap</i>	Israel	AF020082	564
<i>P. papatasi</i>	<i>wPap</i>	India	AF237882	563
<i>P. papatasi</i>	<i>wPap</i>	Israel	AF237883	563
<i>P. papatasi</i>	-	Egypt	AY288297	561
<i>P. papatasi</i>	<i>papa01</i>	Tunisia/Iran	EU780683	563
<i>P. mongolensis</i>	(Turk 54)			
<i>P. caucasicus</i>				
<i>Phasgonophora sulcata</i>	-	Canada	HM355820	596
<i>P. sulcata</i>	-	Canada	HM355823	596
<i>P. sulcata</i>	-	Canada	HM355821	596
<i>Hasgonophora sulcata</i>	-	Canada	HM355824	596
<i>Culex pipiens quinquefasciatus</i>	<i>Wpip</i>	United States of America	AF301012	558
<i>Cinara cedri</i>	<i>wsp_BS_Salamanca(CCeS)</i>	Spain	JN384046	585
<i>Bemisia tabaci</i>		Bangladesh	JQ305095	490
<i>Phlebotomus longicuspis</i>	<i>wPhleb54d</i>	France	JN315985	517
<i>P. perniciosus</i>	<i>wPrn</i>	Italy	AF237884	554
<i>P. perniciosus</i>	-	France	DQ402519	510
<i>Cimex lectularius</i>	<i>TIH</i>	Japan	AB475129	562
<i>Calyptrate muscoid</i> fly	<i>A_NY_Calyp150743b</i>	United States of America	EU126486	501
<i>Simulium damnosum</i>	-	Ghana	FN563975	565
<i>Bemisia tabaci</i>	<i>wBtab ch2</i>	China	FJ404651	601
<i>B. tabaci</i>	<i>wBtab ch4</i>	China	FJ404653	607

Wolbachia is also interesting in an evolutionary context due to the diversity and elegance of its interactions with its hosts and because the barriers to cross cross-mating between populations may reinforce the genetic divergence between them or speciation (Laven 1967, Bordenstein 2003).

The presence of *W. pipientis* is not strictly associated with ZCL endemicity, as this bacterium has been found in populations of *P. papatasi* from non-endemic areas of Iran (Parvizi et al. 2003).

Independent of their mitochondrial haplotype, geographical origin and habitat, wild *P. papatasi* have been found to be either uninfected with *W. pipientis* or infected with one common widespread strain. This raises the possibility of using a genetically modified strain of *W. pipientis* to drive transgenes through wild sandfly populations to intervene in the transmission of *Leishmania*.

CI is a recognised phenotype caused by natural *Wolbachia* infection in sandflies. It is believed that *Wolbachia* might prevent the transmission of virus and parasite infections (Dobson et al. 2002).

Until now, it was believed that individual strains of *Wolbachia* could only be assigned to a particular species of sandfly. However, in the present study, it has been shown that one species of sandfly can be infected with different *Wolbachia* strains and that different species of sandflies can be infected either with a common haplotype or common strain. While *Wolbachia* strains were also found in two species of the *Paraphlebotomus* subgenus for the first time in the present study, the first two observations are more significant. The *wsp* gene is a very useful tool for typing different *Wolbachia* strains. The frequent incidence of *Wolbachia* strains and their ability to manipulate host reproduction has led to *Wolbachia* being proposed as important agents in the evolution of arthropod hosts. *Wolbachia* provides a starting point for inducing changes in host sex or sexuality. By manipulating *Wolbachia* via transgenes, it is hoped that these bacteria may be able to be used as a system to decrease vector-borne-diseases and to reduce the transmission of diseases in the future. As a main vector of *L. major*, these bacteria can be employed as a gene-driving system for spreading anti-pathogen transgenes through wild populations of sandflies.

The infection rates of *Wolbachia* among the observed Iranian populations of *P. mongolensis* and *P. caucasicus* were low and the reason for such low infection rates could be that there are different densities of the infecting bacteria in different locations. The low detection rates of *Wolbachia* in *P. caucasicus* males in comparison with *P. mongolensis* males and the females of both species can be explained by the fact that most of the *P. mongolensis* males and the females of both species were caught in gerbil burrows. Therefore, it is possible that *Wolbachia* might be spread by transgenes in gerbil burrows.

Further studies examining *Wolbachia* species in sandflies are needed to reveal the relationship between *Wolbachia* endosymbionts and phlebotomine hosts.

ACKNOWLEDGEMENTS

To Mehdi Baghban, for helping with the field working, and to Elnaz Alaei Novin, for helping in Molecular Systematics Laboratory.

REFERENCES

- Ahrens M, Shoemaker D 2005. Evolutionary history of *Wolbachia* infections in the fire ant *Solenopsis invicta*. *BMC Evol Biol* 5: 35.
- Azpurua J, de la Cruz D, Valderama A, Windsor D 2010. *Lutzomyia* sand fly diversity and rates of infection by *Wolbachia* and an Exotic *Leishmania* species on Barro Colorado Island, Panama. *PLoS Negl Trop Dis* 4: e627.
- Baldo L, Dunning HJC, Jolley KA, Bordenstein SR, Biber SA, Choudhury RR, Hayashi C, Maiden MC, Tettelin H, Werren JH 2006. Multilocus sequence typing system for the endosymbiont *Wolbachia pipientis*. *Appl Environ Microbiol* 72: 7098-7110.
- Baldo L, Lo N, Werren J 2005. Mosaic nature of the *Wolbachia* surface protein. *J Bacteriol* 187: 5406-5418.
- Beard CB, O'Neill SL, Tesh RB, Richards FF, Aksoy S 1993. Modification of arthropod vector competence via symbiotic bacteria. *Parasitol Today* 9: 179-183.
- Benlarbi M, Ready PD 2003. Host-specific *Wolbachia* strains in widespread populations of *Phlebotomus perniciosus* and *P. papatasi* (Diptera: Psychodidae) and prospects for driving genes into these vectors of *Leishmania*. *Bull Entomol Res* 93: 383-391.
- Bordenstein SR 2003. Symbiosis and the origin of species. In K Bourtzis, T Miller, *Insect symbiosis*, CRC Press, New York, 320 pp.
- Bordenstein SR, O'Hara FP, Werren JH 2001. *Wolbachia*-induced incompatibility precedes other hybrid incompatibilities in *Nasonia*. *Nature* 409: 707-710.
- Bourtzis K, O'Neill SL 1998. *Wolbachia* infection and arthropod reproduction. *Bioscience* 48: 287-293.
- Breeuwer JA, Werren JH 1990. Microorganisms associated with chromosome destruction and reproductive isolation between two insect species. *Nature* 346: 558-560.
- Charlat S, Nirgianaki A, Bourtzis K, Mercot H 2002. Evolution of *Wolbachia*-induced cytoplasmic incompatibility in *Drosophila simulans* and *D. sechellia*. *Evolution* 56: 1735-1742.
- Cui W, Zhang SN, Chen W, Morgan EH 1999. Strong aperiodic x-ray variability and quasi-periodic oscillation in x-ray Nova XTE J1550-564. *Astro J Lett* 512: 1-12.
- Dobson SL, Fox CW, Jiggins FM 2002. The effect of *Wolbachia*-induced cytoplasmic incompatibility on host population size in natural and manipulated systems. *Proc Biol Sci* 269: 437-445.
- Espino CI, Gómez T, González G, Brazil do Santos MF, Solano J, Sousa O, Moreno N, Windsor D, Ying A, Vilchez S, Osuna A 2009. Detection of *Wolbachia* bacteria in multiple organs and feces of the triatomine insect *Rhodnius pallescens* (Hemiptera, Reduviidae). *Appl Environ Microbiol* 75: 547-550.
- Esseghir S, Ready PD, Ben-Ismail R 2000. Speciation of *Phlebotomus* sandflies of the subgenus *Larrousius* coincided with the late Miocene-Pliocene aridification of the Mediterranean subregion. *Biol J Linn Soc Lond* 70: 189-219.
- Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH 2008. How many species are infected with *Wolbachia*? A statistical analysis of current data. *FEMS Microbiol Lett* 281: 215-220.
- Hoffmann AA, Turelli M 1997. Cytoplasmic incompatibility in insects. In SL O'Neill, AA Hoffman, JH Werren (eds.), *Influential passengers: microbes and arthropod reproduction*, Oxford University Press, Oxford, p. 42-80.
- Huigens ME, Luck RF, Klaassen RHG, Maas MFP, Timmermans MJTN, Stouthamer R 2000. Infectious parthenogenesis. *Nature* 672: 178-179.
- Hurst GDD, Schilthuizen M 1998. Selfish genetic elements and speciation. *Heredity* 80: 2-8.

- Jeyaprakash A, Hoy M 2000. Long PCR improves *Wolbachia* DNA amplification *wsp* sequences found in 76% of sixty-three arthropod species. *Insect Mol Biol* 9: 393-405.
- Killick-Kendrick R 1999. The biology and control of phlebotomine sand flies. *Clin Dermatol* 17: 279-289.
- Kondo N, Nikoh N, Ijichi N, Shimada M, Fukatsu T 2002. Genome fragment of *Wolbachia* endosymbiont transferred to X chromosome of host insect. *Proc Natl Acad Sci USA* 99: 14280-14285.
- Kyei-Poku GK, Colwell DD, Coghlin P, Benkel B, Floate KD 2005. On the ubiquity and phylogeny of *Wolbachia* in lice. *Mol Ecol* 14: 285-294.
- Laven H 1967. Speciation and evolution in *Culex pipiens*. In R Wright, R Pal (eds.), *Genetic of insect vectors of disease*, Elsevier, Amsterdam, p. 251-275.
- Lewis DJ 1982. A taxonomic review of the genus *Phlebotomus*. *Bull Brit Nat Hist (Entomol)* 45: 121-209.
- Lo N, Casiraghi M, Salati E, Bazzocchi C, Bandi C 2002. How many *Wolbachia* supergroups exist? *Mol Biol Evol* 19: 341-346.
- Mirzaei A, Roushani S, Taherkhani H, Farahmand M, Kazemi B, Hedayati M, Baghaei A, Davari B, Parvizi P 2011. Isolation and detection of *Leishmania* species among naturally infected *Rhombomys opimus*, a reservoir host of zoonotic cutaneous leishmaniasis in Turkmen Sahara, North East of Iran. *Exp Parasitol* 129: 375-380.
- Mohebbi M, Khamesipour A, Mobedi I, Zarei Z, Hashemi-Fesharki R 2004. Double-blind randomized efficacy field trial of alum precipitated autoclaved *Leishmania major* vaccine mixed with BCG against canine visceral leishmaniasis in Meshkin-Shahr district, I.R. Iran. *Vaccine* 22: 4097-4100.
- Moran NA, McCutcheon JP, Nakabachi A 2008. Genomics and evolution of heritable bacterial symbionts. *Annu Rev Genet* 42: 165-190.
- Nadim A, Faghih M 1968. The epidemiology of cutaneous leishmaniasis in the Isfahan province of Iran. I. The reservoir. II. The human disease. *Trans R Soc Trop Med Hyg* 61: 534-542.
- O'Neill HW, Hoffmann AA, Werren JH 1997. *Influential passengers: inherited microorganisms and arthropod reproduction*, Oxford University Press, Oxford, 320 pp.
- O'Neill SL, Giordano R, Colbert AME, Karr TL, Robertson HM 1992. 16S rRNA phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic incompatibility in insects. *Proc Natl Acad Sci USA* 89: 2699-2702.
- Ono M, Braig HR, Munstermann LE, Ferro C, O'Neill SL 2001. *Wolbachia* infections of phlebotomine sand flies (Diptera: Psychodidae). *J Med Entomol* 38: 237-241.
- Parvizi P, Benlarbi M, Ready PD 2003. Mitochondrial and *Wolbachia* markers for the sandfly *Phlebotomus papatasi*: little population differentiation between peridomestic sites and gerbil burrows in Isfahan province, Iran. *Med Vet Entomol* 17: 351-362.
- Parvizi P, Naddaf SR, Alaei-Novin E 2010a. Molecular typing and phylogenetic analysis of some species belonging to *Phlebotomus* (*Larrousius*) and *Phlebotomus* (*Adlerius*) subgenera (Diptera: Psychodidae) from two locations in Iran. *Iran J Arthropod Borne Dis* 4: 1-10.
- Parvizi P, Ready PD 2008. Nested PCRs of nuclear ITS-rDNA fragments detect three *Leishmania* species of gerbils in sandflies from Iranian foci of zoonotic cutaneous leishmaniasis. *Trop Med Int Health* 13: 1159-1171.
- Parvizi P, Taherkhani H, Ready PD 2010b. *Phlebotomus caucasicus* and *Phlebotomus mongolensis* (Diptera: Psychodidae): indistinguishable by the mitochondrial cytochrome *b* gene in Iran. *Bull Entomol Res* 100: 415-420.
- Pascal C, Pintureau B, Katchadourian C, Grenier S, Bolland P, Robin C, Vallier A 2006. Comparison of *Wolbachia* bacterial density in females of four thelytokous strains of *Trichogramma cordubensis* and *T. evanescens* (Hymenoptera, Trichogrammatidae). *Vestn Zool* 40: 417-425.
- Pintureau B, Chaudier S, Lassabliere F, Charles H, Grenier S 2000. Addition of *wsp* sequences to the *Wolbachia* phylogenetic tree and stability of the classification. *J Mol Evol* 51: 374-377.
- Rousset F, Solignac M 1995. Evolution of single and double *Wolbachia* symbioses during speciation in the *Drosophila simulans* complex. *Proc Natl Acad Sci USA* 92: 6389-6393.
- Schaub GA, Boker CA, Jensen C, Reduth D 1989. Cannibalism and coprophagy are modes of transmission of *Blastocrithidia triatomae* (Trypanosomatidae) between triatomines. *J Protozool* 36: 171-175.
- Shoemaker DD, Machado CA, Molbo D, Werren JH, Windsor DM, Herre EA 2002. The distribution of *Wolbachia* in fig wasps: correlations with host phylogeny, ecology and population structure. *Proc R Soc Lond B Biol Sci* 269: 2257-2267.
- Sinkins SP, Curtis CF, O'Neill SL 1997. The potential application of inherited symbiont systems to pest control. In SL O'Neill, A Hoffman, J Werren (eds.), *Influential passengers*, Oxford University Press, Oxford, p. 155-175.
- Stouthamer R, Breeuwer JAJ, Hurst GDD 1999. *Wolbachia pipientis*: microbial manipulator of arthropod reproduction. *Annu Rev Microbiol* 53: 71-102.
- Telschow A, Hammerstein P, Werren JH 2005. The effect of *Wolbachia* versus genetic incompatibilities on reinforcement and speciation. *Evolution* 59: 1607-1619.
- Testa JM, Montoya-Lerma J, Cadena H, Oviedo M, Ready PD 2002. Molecular identification of vectors of *Leishmania* in Colombia: mitochondrial introgression in the *Lutzomyia townsendi* series. *Acta Trop* 84: 205-218.
- Theodor O, Mesghali A 1964. On the Phlebotominae of Iran. *J Med Entomol* 1: 285-300.
- Van Meer MMM, Witteveldt J, Stouthamer R 1999. Phylogeny of the arthropod endosymbiont *Wolbachia* based on *wsp* gene. *Insect Mol Biol* 8: 399-408.
- Werren JH 1997. Biology of *Wolbachia*. *Annu Rev Entomol* 42: 587-609.
- Werren JH, Baldo L, Clark ME 2008. *Wolbachia*: master manipulators of invertebrate biology. *Nat Rev Microbiol* 6: 741-751.
- Werren JH, Guo LR, Zhang W 1995. Evolution and phylogeny of *Wolbachia*: reproductive parasites of arthropods. *Proc R Soc Lond B Biol Sci* 251: 55-63.
- Werren JH, Windsor DM 2000. *Wolbachia* infection frequencies in insects: evidence of a global equilibrium? *Proc R Soc Lond B Biol Sci* 267: 1277-1286.
- West SA, Cook JM, Werren JH, Godfray HC 1998. *Wolbachia* in two insect host-parasitoid communities. *Mol Ecol* 7: 1457-1465.
- Wu K, Hoy MA 2012. *Cardinium* is associated with reproductive incompatibility in the predatory mite *Metaseiulus occidentalis* (Acari: Phytoseiidae). *J Invertebr Pathol* 110: 359-365.
- Yaghoobi-Ershadi MR, Akhavan AA, Mohebbi M 1996. *Meriones libycus* and *Rhombomys opimus* (Rodentia: Gerbillidae) are the main reservoir hosts in a new focus of zoonotic cutaneous leishmaniasis in Iran. *Trans R Soc Trop Med Hyg* 90: 503-504.
- Yaghoobi-Ershadi MR, Zahraei-Ramazani AR, Akhavan AA, Jalali-Zand AR, Abdoli H, Nadim A 2005. Rodent control operations against zoonotic cutaneous leishmaniasis in rural Iran. *Ann Saudi Med* 25: 309-312.
- Zhou W, Rousset F, O'Neill S 1998. Phylogeny and PCR-based classification of *Wolbachia* strains using *wsp* gene sequences. *Proc R Soc Lond B Biol Sci* 265: 509-515.