

Relationship between orofacial pain and absenteeism among workers in Southern Brazil

Anderson Nardi¹, Edgard Michel-Crosato², Maria Gabriela Haye Biazevic³,
Edgard Crosato⁴, Eduardo Pizzatto⁵, Dagmar de Paula Queluz⁶

¹MS, Universidade do Oeste de Santa Catarina (Unoesc), Joaçaba (SC), Brazil

²DDS, MS, PhD, Assistant Professor, Department of Community Dentistry, Faculdade de Odontologia, Universidade de São Paulo (USP), São Paulo (SP), Brazil

³DDS, MS, PhD, Post-doctoral student, Department of Community Dentistry, Faculdade de Odontologia, USP, São Paulo (SP), Brazil

⁴DDS, MS, PhD, Professor, Department of Community Dentistry, Faculdade de Odontologia, USP, São Paulo (SP), Brazil

⁵DDS, MSPH, PhD, Professor, Faculdade de Odontologia, Universidade Positivo, Curitiba (PR), Brazil

⁶DDS, MSPH, PhD, Professor, Department of Community Dentistry, Faculdade de Odontologia de Piracicaba, Universidade Estadual de Campinas (Unicamp), Campinas (SP), Brazil.

Abstract

Aim: To verify the relationship between orofacial pain and absenteeism in workers of slaughter and meat processing industries in the Southern region of Brazil. **Methods:** A cross-sectional study, with the random sample of 401 workers of slaughter and meat processing industries in the Southern region of Brazil, was carried out. A questionnaire referred to the situation of absenteeism caused by nine different types of orofacial pain and also the amount of time the employee was kept from work. **Results:** Only 60 workers (15%) reported having missed work due to orofacial pain in the six months prior to the study. The prevalence of absenteeism resulting from orofacial pain was of 15%. The types of orofacial pain that resulted in absenteeism were: spontaneous toothache (9.7 %); toothache caused by cold or hot liquids or by sweet foodstuff (6.5%) and pain around and behind the eyes (3.2%). There was a predominance of absenteeism in half and full work shifts for the types of orofacial pain experienced. Associations between absenteeism from induced toothache and gender ($p < 0.05$), absenteeism and spontaneous toothache and family income ($p = 0.011$), and between absenteeism and the self awareness of their oral health condition, as well as the nine types of orofacial pain ($p < 0.001$) were observed. **Conclusions:** The prevalence of absenteeism as a result of orofacial pain was low.

Keywords: absenteeism, facial pain, facial pain/epidemiology, oral health.

Introduction

Absenteeism is an issue of growing interest as a result of the economic importance of competitiveness, driving companies to seek means to reduce its occurrence and, consequently, to increase the profitability and achieve sustained growth^{1,2}.

Various epidemiological studies have demonstrated that the prevalence of absenteeism resulting from dental reasons varies from 10 to 35%, and the average number of working hours lost varies from 1.24 to 6.20 working hours/workers/years³⁻⁹.

The pain is a private percept that arises in a conscious brain, typically in response to a noxious provoking stimulus, but, sometimes, in the absence of a stimulus. The relation of the percept to the stimulus is variable, and depends on the individual's prior expectations and beliefs, and on his/her cognitive and emotional state – not just on the nature of the stimulus itself. While acute pain is, by definition, a brief and self-limiting process, chronic pain comes to dominate the life and concerns of the patient, and often also family, friends and other caregivers. In addition to the severe erosion in quality of life of the pain sufferer and those around

Received for publication: July, 17, 2008

Accepted: February 04, 2009

Correspondence to:

Edgard Michel Crosato

Departamento de Odontologia Social
da Faculdade de Odontologia da USP
Avenida Professor Lineu Prestes, 2.227 –
Cidade Universitária
CEP 05508-000 – São Paulo (SP), Brazil

E-mail: michelcrosato@usp.br

him/her, chronic pain imposes severe financial burdens on many levels. These include: costs of healthcare services and medication, job absenteeism and disruption in the workplace, loss of income, non-productivity in the economy and in the home, financial burden on family, friends and employers, worker compensation costs and welfare payments. The workers can reduce absenteeism and healthcare utilization from the risks from dental disease by incorporating dental education into workplace wellness.

The aim of this study was to verify the relationship between orofacial pain and absenteeism in workers of slaughter and meat processing industries in the Southern region of Brazil.

Material and methods

The study population was composed of a sample of 401 workers of slaughter and meat processing industries in the Southern region of Brazil. The company workforce was composed of 1,187 employees. Sample size calculation was done using the following criteria: confidence level 95%, sample mistake 4% and unknown prevalence, and estimated prevalence 50%.

The selection process of the samples was realized through simple random drawings. An observational and cross-sectional study design was used.

Data collected referred to absenteeism resulting from orofacial pain, as well as social and economic characteristics of the study population (gender, age, marital status, area of work, address, educational level and family income).

A questionnaire developed by Locker & Grushka⁶⁻⁷ was used as an instrument for the survey, which was validated in Brazil in the Bambuí Health and Ageing Study (BHSA)¹⁰.

The questions in the survey instrument referred to the situation of absenteeism caused by nine different types of orofacial pain and also the amount of time the employee was kept from work. The questionnaire was prepared to gather information about orofacial pain and absenteeism at present or in the recent past. All questions referred to the period encompassing the six months prior to the survey. This relatively short period of time is normally used to minimize bias due to possible lapse of memory of the surveyed study population^{11,12}.

About the family income, low means until two minimum wage, and high means two or more minimum wages. The oral health, it was self related good or poor.

The questionnaire was first applied to 15 employees. Thereafter, it was applied by duly trained surveyors to the employees that agreed to be enrolled as volunteers after granting authorization from the company management and after the participants had signed an informed consent form to take part in the study. The research project was approved by the Research Ethics Committee of the Universidade de Passo Fundo (UPF) under the protocol number 194/03.

Data were analyzed by the chi-square test using the Stata 8.0 software and presented in tables, according to the frequency distribution. A 5% significance level was adopted for all analyses.

Results

The social and economic status of the studied population is shown on **Table 1**.

Among the participants in the study, 60 employees declared having lost working hours due to orofacial pain in the six months prior to the survey. Consequently, the prevalence of absenteeism due to orofacial pain in this study was of 15%. The types of orofacial pain that cause more absenteeism were: spontaneous toothache (9.7%); toothache caused by cold or hot liquids or sweet foodstuff (6.5%) and pain around and behind the eyes (3.2%).

Table 2 shows the percentage of loss of working hours for each of the nine types of orofacial pain described. The predominance of loss of half and full work shifts was observed for all types of orofacial pain experienced. A half work shift was considered as a period of four hours, in other words, half a day's work. The types of orofacial pain that caused two and three days or more of absence from work were the spontaneous toothache, toothache caused by hot or cold liquids or sweet foodstuff, pain in front of the ears and pain around and behind the eyes.

Considering the prevalence of absenteeism according to the number of types of pain experienced by workers, 26 employees informed having been absent from work due to one type of orofacial pain (6.5%), 18 employees informed having been absent as a result of two types of orofacial pain (4.5%), nine employees informed having missed work as a result of feeling more than three types of orofacial pain and only one worker informed having missed working days as a result of feeling the nine types of orofacial pain (0.2%) in the last six months (**Table 2**).

The association between absenteeism and gender indicated that the prevalence of absenteeism resulting from toothache caused by hot or cold liquids or sweet foodstuff ($p = 0.043$) was higher for male workers (**Table 3**).

Table 1. Percentage of gender, age, marital status, education level, family income among workers of a meat processing industry. Joaçaba (SC), Brazil, 2003

Variable	Category	n	%
Gender	Male	285	71.1
	Female	116	28.9
Age	18 to 21	91	22.7
	22 to 31	191	47.6
	32 to 41	99	24.7
	42 to 51	20	5.0
Marital status	Single	178	44.4
	Married	223	55.6
Education level	Cannot read or write	1	0.2
	Incomplete Elementary School	75	18.7
	Complete Elementary School	78	19.5
	Incomplete High School	73	18.2
	Complete High School	142	35.4
Family income	College Education	32	8.0
	From 1 to 2 minimum wages	180	44.9
	From 3 to 5 minimum wages	182	45.4
	From 6 to 10 minimum wages	33	8.2
	From 11 to 20 minimum wages	6	1.5

Table 4 shows that the prevalence of absenteeism resulting from spontaneous toothache was higher among workers with the worst level of family income ($p = 0.011$).

An association was also observed between absenteeism resulting from orofacial pain and self awareness of oral health ($p < 0.001$). Employees who declared having poor oral health conditions presented a higher prevalence of absenteeism resulting from orofacial pain, in the nine types of orofacial pain studied, than those who declared having a good oral health status (**Table 5**).

Discussion

The prevalence of absenteeism due to orofacial pain for the workers of the meat processing industry was of 15%, and the average number of lost working hours as a result of dental causes was of 0.88 working hours/employee/year.

When comparing the results of this study to those of various epidemiological studies³⁻⁹, it was noted that the prevalence of absenteeism for dental reasons in the study population matched the previously found values. However, the average number of working hours lost

by the employees of the meat processing industry was below those results previously presented.

The orofacial pains that caused the highest levels of absenteeism were: spontaneous toothache (9.7%); toothache caused by cold or hot liquids or sweet foodstuff (6.5%) and pain around and behind the eyes (3.2%). These pains have the characteristic of acute cases with a relatively short duration and that can be rapidly treated, reestablishing the workers' capabilities to perform their functions and enabling their prompt return to their jobs. Considering the amount of time absent from work as a result of orofacial pain, there was a predominance for the loss of half or a full work shift as shown in **Table 2**. The average number of leave-of-absence days due to dentist certificates was of 1.5 days/employee/year. These results are consistent with those presented in previous studies¹¹⁻¹⁴.

Absenteeism caused by dental reasons in this meat processing industry was smaller than absenteeism resulting from medical causes in both average length and total number of leaves of absence, being these data similar to the survey by Reisine¹⁵.

Employees who declared having poor oral health status presented a higher prevalence of absenteeism due to orofacial pain than those who declared having a good oral health status. In this sense,

Table 2. Percentage of working hours lost for each type of orofacial pain among workers of a meat processing industry. Joaçaba (SC), Brazil, 2003

Type of orofacial pain	No absence (%)	Absent on half a shift (%)	Absent for a full shift (%)	Absent for 1 day (%)	Absent for 2 days (%)	Total (%)
Spontaneous toothache	90.7	5.2	1.4	2.2	0.5	100.0
Induced toothache	93.5	3.7	1.4	1.2	0.2	100.0
Burning sensation on the tongue	98.8	0.5	0.2	0.5	0.0	100.0
TMJ pain	97.5	1.2	0.8	0.5	0.0	100.0
Pain on chewing	98.0	1.0	0.5	0.5	0.0	100.0
Pain on opening the mouth	99.0	0.5	0.3	0.2	0.0	100.0
Pain in front of the ears	97.5	1.5	0.0	0.5	0.5	100.0
Pain in the face or cheeks	97.3	1.5	1.0	0.2	0.0	100.0
Pain around or behind the eyes	97.0	1.3	1.4	0.3	0.0	100.0

Table 3. Association between absenteeism resulting from facial pain and gender among the workers of a meat processing industry. Joaçaba (SC), Brazil, 2003

Causes of absenteeism		Gender		Odds Ratio	Confidence interval (95%)	p
		Male	Female			
Spontaneous toothache	No	253	109	1.970	0.843 - 4.599	0.111
	Yes	32	7			
Induced toothache	No	262	113	3.307	0.973 - 11.236	0.043
	Yes	23	3			
Burning sensation on the tongue	No	281	115	1.637	0.181 - 14.804	0.658
	Yes	4	1			
TMJ pain	No	278	113	0.948	0.241 - 3.733	0.940
	Yes	7	3			
Pain on chewing	No	279	114	1.226	0.244 - 6.164	0.805
	Yes	6	2			
Pain on opening the mouth	No	283	114	0.403	0.056 - 2.894	0.350
	Yes	2	2			
Pain in front of the ears	No	280	111	0.396	0.113 - 1.396	0.137
	Yes	5	5			
Pain in the face or cheeks	No	277	113	1.088	0.283 - 4.175	0.902
	Yes	8	3			
Pain around or behind the eyes	No	275	113	1.370	0.370 - 5.070	0.636
	Yes	10	3			

Table 4. Association between absenteeism from orofacial pain and family income among the workers of a meat processing industry. Joaçaba (SC), Brazil, 2003

Causes of absenteeism		Income		Odds Ratio	Confidence interval (95%)	p
		Low	High			
Spontaneous toothache	No	155	207	2.385	1.200 - 4.738	0.011
	Yes	25	14			
Induced toothache	No	166	209	1.469	0.662 - 3.261	0.342
	Yes	14	12			
Burning sensation on the tongue	No	177	219	1.856	0.307 - 11.229	0.494
	Yes	3	2			
TMJ pain	No	174	217	1.871	0.520 - 6.733	0.331
	Yes	6	4			
Pain on chewing	No	176	217	1.233	0.304 - 5.000	0.769
	Yes	4	4			
Pain on opening the mouth	No	178	219	1.230	0.172 - 8.822	0.836
	Yes	2	2			
Pain in front of the ears	No	174	217	1.871	0.520 - 6.733	0.331
	Yes	6	4			
Pain in the face or cheeks	No	174	216	1.490	0.447 - 4.963	0.514
	Yes	6	5			
Pain around or behind the eyes	No	175	213	0.761	0.244 - 2.367	0.636
	Yes	5	8			

Tabela 5. Association between absenteeism from orofacial pain and self-awareness of oral health among workers of a meat processing industry. Joaçaba (SC), Brazil, 2003

Causes of absenteeism		Oral health		Odds Ratio	Confidence interval (95%)	p
		Good	Poor			
Spontaneous toothache	No	37	325	6.787	3.309 - 13.923	0.000
	Yes	17	22			
Induced toothache	No	39	336	11.748	5.043 - 27.370	0.000
	Yes	15	11			
Burning sensation on the tongue	No	50	346	27.680	3.033 - 252.646	0.000
	Yes	4	1			
TMJ pain	No	47	344	17.078	4.269 - 68.321	0.000
	Yes	7	3			
Pain on chewing	No	48	345	21.563	4.231 - 109.892	0.000
	Yes	6	2			
Pain on opening the mouth	No	50	347	7.940	6.127 - 10.289	0.000
	Yes	4	0			
Pain in front of the ears	No	49	342	6.980	1.950 - 24.984	0.001
	Yes	5	5			
Pain in the face or cheeks	No	49	341	5.799	1.705 - 19.722	0.002
	Yes	5	6			
Pain around or behind the eyes	No	45	343	17.150	5.073 - 57.980	0.000
	Yes	9	4			

self-awareness of the oral health status coincided with the impact observed through the application of the instruments: work-related activities are more severely affected among employees who presented the perception of their poor oral health condition.

This study did not demonstrate any associations between prevalence of absenteeism due to orofacial pain and age, marital status, schooling, geographical area (rural or urban), company sector or work shift.

There are two types of absenteeism: absenteeism through the absence of work and physically present absenteeism. The first type can be measured and its costs can be calculated through the absence.

The second type cannot be measured, since it represents the worker who cannot perform his/her normal working activities due to pain, despite being physically present to the workspace¹³.

Labor is increasingly becoming effective and instrumental in the social-economical advancements in our society¹⁴. The attention of the authorities responsible for the implementation of the directives of a country that has its economy based on labor should be oriented towards the health and welfare. Consequently, workers should be the objective of measures and policies to preserve their physical, mental and social well-being. Therefore, maximum productive capacity is achieved when the worker is satisfied in his basic health needs¹¹⁻¹⁶.

Studies have shown that oral problems caused difficulties or incapacity to perform normal working activities, study or sleep in a percentage that ranges from 8 to 60%¹⁶⁻¹⁸.

In the present study, associations between absenteeism from induced toothache and gender ($p = 0.05$); absenteeism, spontaneous toothache and family income ($p = 0.011$); and absenteeism, self-awareness of the oral health condition and the nine types of orofacial pain ($p = 0.00$) were observed. The prevalence of absenteeism as a result of orofacial pain was low.

It is important to point out that the methodological aspects of future epidemiological studies about orofacial pain and absenteeism must be standardized, in such a way that the results of the various studies could be compared with greater reliability.

Since absenteeism for dental reasons was not pronounced, the company did not present losses in productivity because their work force contemplates a surplus percentage of employees to compensate for absences.

References

1. Gift HC, Reisine ST, Larach DC. The social impact of dental problems and visits. *Am J Public Health*. 1992;82:1663-8.
2. Berndt ER, Bailit HL, Keller MB, Verner JC, Finkelstein SN. Health care use and at-work productivity among employees with mental disorders. *Health Aff*. 2000;19:244-56.
3. Midorikawa ET. Odontology in worker's health as a new professional specialty: definition of the activity field and functions of the surgeon dentist in worker's health team. [Doctoral Tesis]. São Paulo: Faculdade de Odontologia of Universidade de São Paulo; 2000. 337p.
4. Hollister MC, Weintraub JA. The association of oral status with systemic health, quality of life, and economic productivity. *J Dent Educ*. 1993;57:901-12.
5. Schou L. Oral health promotion at worksites. *Int Dent J*. 1989;39:122-8.
6. Locker D, Grushka M. The impact of dental and facial pain. *J Dent Res*. 1987;66:1414-7.
7. Locker D, Grushka M. Prevalence of oral and facial pain and discomfort: preliminary results of a mail survey. *Community Dent Oral Epidemiol*. 1987;15:169-72.
8. Hooper HA. Dental services in industry: observations on their effects in the reduction on absenteeism. *Industrial Medicine*. 1942;11:157-62.
9. Bailit H, Beazoglou T, Hoffman W. Work loss and dental disease. Report to the Robert Wood Johnson Foundation. University of Connecticut Health Center; 1982.
10. Matos DL, Lima Costa MF, Guerra HL, Marcenes W. Projeto Bambuí: avaliação de serviços odontológicos privados, públicos e de sindicato. *Rev Saúde Pública*. 2002;36:237-43.
11. Macfarlane TV, Blinkhorn AS, Davies RM, Kinsey J, Worthington HV. Orofacial pain in the community: prevalence and associated impact. *Community Dent Oral Epidemiol*. 2002;30:52-60.
12. Reisine ST, Miller J. A longitudinal study of work loss related to dental diseases. *Soc Sci Med*. 1995;21(12):1309-14.
13. Jaafar N, Razak IA, Zain RB. The social impact of oral and facial pain in an industrial population. *Ann Acad Med*. 1989;18:553-5.
14. Naito M, Yuasa H, Nomura Y, Nakayama T, Hamajima N, Hanada N. Oral health status and health-related quality of life: a systematic review. *J Oral Sci*. 2006;48:1-7.
15. Reisine ST. Dental disease and work loss. *J Dent Res*. 1984;63:1158-61.
16. Reisine ST. Dental health and public policy: the social impact of dental disease. *Am J Public Health*. 1985;75:27-30.
17. Reisine ST. The impact of dental conditions on social functioning and the quality of life. *Annu Rev Public Health*. 1988;9:1-19.
18. Johnson NW, Glick M, Mbuguye TN. Oral health and general health. *Adv Dent Res*. 2006;19:118-21.