

their association with nerves and vessels and are particularly useful, when surgical management of the lesion is contemplated^[2]

MRI of lymphangioma usually shows a multiloculated heterogeneous mass with low signal intensity on T1-weighted images, and high signal intensity on T2-weighted images because of its content. The authors' experience suggests that most lymphangiomas have a characteristic appearance on MRI. The information obtained with MRI can help in providing a preoperative diagnosis, in planning surgical resection, and in defining recurrence^[5].

A total surgical excision, if feasible without a major sacrifice of adjacent organs, can be the best therapeutic option^[3,6]. Sclerotherapy presents an interesting therapeutic method in invasive forms^[1]. In our patient complete resection was performed. The anatomopathologic study confirmed the diagnosis. Postoperative course was simple.

The presentation of this case draws attention to the possibility of occurrence of such an etiologic lesion in rectus abdominis muscle. We stress the necessity of imaging as a diagnostic tool for this location.

Key words: Lymphangioma; Cysts; Rectus Abdominis Muscle

References

1. Mirza B, Ijaz L, Saleem M, et al. Cystic hygroma: an overview. *J Cutan Aesthet Surg* 2010; 3(3):139-44.
 2. Wiegand S, Eivazi B, Barth PJ, et al. Pathogenesis of lymphangiomas. *Virchows Arch* 2008; 453(1):1-8.
 3. Erol O, Ozçakar L, Inanici F. Cystic hygroma in the quadriceps muscle: a sanguine diagnosis for knee pain. *Joint Bone Spine* 2005; 72(3):267-9.
 4. Wever DJ, Heeg M, Mooyaart EL. Cystic hygroma of the shoulder region. A case report. *Clin Orthop Relat Res* 1997; 338:215-8.
 5. Siegel MJ, Glazer HS, St Amour TE, et al. Lymphangiomas in children: MR imaging. *Radiology* 1989; 170(2):467-70.
- Bezzola T, Bühler L, Chardot C, et al. Surgical therapy of abdominal cystic lymphangioma in adults and children. *J Chir (Paris)* 2008; 145(3):238-43.

Cardiac Tamponade; A rare Presentation of Childhood Hypothyroidism

Sudhir Mehta

Department of Pediatrics, Sri Aurobindo Institute of Medical Sciences and PG Institute, Indore, India

Received: Aug 12, 2013; Accepted: Oct 06, 2013;
First Online Available: Oct 30, 2014

Myxedema heart disease is a well known entity. Pericardial effusions have been reported in 50-73% of pediatric patients with hypothyroidism in various series^[1,2], but none of these describes the symptomatic pericardial effusions or cardiac tamponade.

5 year female child presented with complaints of periorbital puffiness and mild abdominal distension intermittently for last 6 months. She was symptomatically treated with diuretics off and on without much benefit. This time she reported to pediatric emergency with progressively increasing respiratory distress for last 5 days. There was no history of any fever, decreased urine output, orthopnea, or Koch's contact. She was third live issue of non consanguineous marriage born by normal vaginal delivery following uncomplicated pregnancy. She was immunized for age with no significant family history. She had history of global development delay, development age corresponded to 3 years. On examination she was afebrile with HR of 70/min, RR of 44/min and BP of 80/60 mmHg with pulsus paradoxus. She had coarse facies with dilated neck veins. Her skin was dry and coarse. She had short stature with height less than 3rd centile and normal weight. Her cardiovascular examination revealed silent precordium. Her cardiac border was extending 1.5 cm beyond right sternal margin and 2 cm beyond apex beat on left side on percussion. Her heart sounds were muffled with no murmur. Her abdominal examination revealed free fluid in abdomen. Rest of systemic examination was normal. Chest X ray revealed massive cardiomegaly with CT ratio of

678%. Immediate ECHO done on urgent basis, revealed large pericardial effusion with collapse of right atrium during diastole with ejection fraction of 65% and intrinsically normal heart. ECHO guided pericardiocentesis drained 150 ml of fluid. The fluid was clear with 3.3 mg/dl of proteins, 110 mg/dl cholesterol, 20 RBC/ μ L, 14 lymphocytes/ μ L, 7

* **Corresponding Author; Address:** Sri Aurobindo Institute of Medical Sciences and PG Institute, Indore, India
E-mail: samsudhir123@yahoo.co.in

neutrophils/ μ L and rest of pericardial examination was normal. Rest of hematological and urinary examination was normal. Searching for tuberculosis was also negative. In view of features suggested of hypothyroidism on clinical examination and non infectious nature of pericardial fluid, her thyroid profile was done which was suggested of hypothyroidism with $T_3=0.92$ nmol/L (1.39-3.7nmol/L), $T_4=3.2\mu$ g/dl (5.5-12.8 μ g/dl), $TSH=40$ mIU/L(0.7-6.4mIU/L). Her anti TPO antibodies were negative and USG neck revealed mild enlargement of thyroid gland. She was started on L-thyroxine with dosage of 4 μ g/kg/d. She responded to treatment with thyroid replacement and her pericardial effusion disappeared after 2 months on repeat ECHO. She is on regular follow up and showing improvement in each regards.

Although mild to moderate pericardial effusions of no clinical significance are frequently reported in pediatric myxedema heart disease, but cardiac tamponade is rare. Literature reports only 2 adolescent girls with cardiac tamponade in association with hypothyroidism, but none has been reported in young children. Author reports a rare case of cardiac tamponade in a young child with previously unrecognized hypothyroidism. Besides bradycardia, minimal pericardial effusions are most common cardiac manifestations of clinical hypothyroidism in adults. Most of pediatric age group patients presenting with this belonged to neonatal period or to Down's syndrome^[1,2]. Myxedema heart disease was first described by Zondek in 1918 and was completely defined by Fahr in 1925^[3]. The pathophysiological derangements responsible for pericardial effusions are increased egress of proteins from blood, decreased lymphatic clearance of proteins and abnormal electrolyte metabolism^[4]. At molecular level, commonly observed signs or symptoms in hypothyroidism are attributed to changes in expression of various gene products, which includes alpha myosin heavy chain, beta-1 adrenergic receptors, voltage gated potassium channels and sarcoplasmic reticulum calcium ATPase^[5]. The rarity of cardiac tamponade in hypothyroidism is attributed to slow accumulation of fluid and marked distensibility of pericardium^[6].

Alexander first described 'Gold paint effusion' to describe the golden brown appearance of pericardial fluid due shimmering satin cholesterol crystals^[7]. High cholesterol content of pericardial fluid have been attributed to disturbance of lipid metabolism; probably churning action of heart causing precipitation of cholesterol from pericardial fluid. Treatment of cardiac tamponade due to hypothyroidism involves urgent pericardiocentesis followed by thyroxine replacement. Treatment of pericardial effusion with thyroxine replacement leads to resolution of effusion in 2-12 months^[5]. In conclusion myxedema heart disease should be suspected in any child with symptoms suggestive of hypothyroidism, presenting with even hemodynamically significant pericardial effusion or cardiac tamponade. Patients with hypothyroidism-associated pericardial effusion should be monitored for development of cardiac tamponade.

Key words: Cardiac Tamponade; Hypothyroidism, Childhood

References

1. Bhupathi R, Kothari SS, Gupta AK, et al. Cardiac function in hypothyroidism: effect of replacement therapy. *Indian Pediatr* 1999; 36(8):779-84.
2. Ferooki ZQ, Hoffman WH, Perry BL et al. Myocardial dysfunction in hypothyroid children. *Am J Dis Child* 1983; 137(1):65-8.
3. Fahr G. Myxedema Heart. *J Amer Med Assoc* 1925; 84:345-49.
4. Parving HH, Hansen JM, Nielsen SL, et al. Mechanism of edema formation in myxedema - increased protein extravasation and relatively slow lymphatic drainage. *N Engl J Med* 1979; 301(9): 460-5.
5. Ladenson PW. Recognition and management of cardiovascular disease relative to hypothyroid dysfunction. *Am J Med* 1990; 88(6):638-41.
6. Lange K. Capillary permeability in myxedema. *Am J Med Sci* 1944; 205:5-15.
7. Alexander JS. Pericardial effusion of 'Gold Paint' appearance due to presence of cholesterol. *Br Med J* 1919; 2(3067):463.