

## Original Article

# Thermal injury in pregnancy: predicting maternal and fetal outcome

**P. Agarwal**

Plastic surgery unit, Department of Surgery, N.S.C.B. Government Medical College, Jabalpur-482003, M. P., India.

**Address for correspondence:** Pawan Agarwal, 292/293, Napier Town, Jabalpur- 482001, M.P., India. E-mail: [drpawanagarwal@yahoo.com](mailto:drpawanagarwal@yahoo.com)

## ABSTRACT

This study was undertaken to assess the parameters that may predict maternal and foetal outcome in 49 thermally injured pregnant women in a Plastic surgery unit in tertiary referral center in the last five years. There were 33 maternal deaths and 34 fetal deaths. In general, pregnancy as such does not influence maternal outcome after thermal injury. In first and second trimester best chance for foetal survival is to ensure maternal survival and in the last trimester fetal survival depends upon fetal maturity. Maternal survival is less likely if the burn wound exceeds 50% total body surface area. Thermal injury does increase the risk of spontaneous abortion and premature labour. Early obstetric intervention is indicated in patients with fatal burn and complications.

## KEY WORDS

Pregnancy, Thermal injury, Outcome

## INTRODUCTION

The literature on burns in pregnancy is limited but this condition is not uncommon in India. The incidence of female burn patients of reproductive age varies from 0.6%-15% in different series with highest incidence reported from India.<sup>1-7</sup> Thermal injury sustained during pregnancy presents special management problems for the gravid woman and her foetus. The paucity of published data on the specific problems of burnt pregnant women has made it difficult to determine the incidence, morbidity, mortality rate or most effective management program for them.<sup>8</sup> More information is required to provide guidelines for plastic surgeons and obstetricians, counseling the patients and her family for any additional risk of abortion, stillbirth, premature delivery, impaired foetal growth or malformations. In this study, an attempt was made to determine the factors responsible for poor maternal and foetal outcome in order to plan interventions

to lower the morbidity and mortality. The present study describes the experience of 49 burn patients who were managed in a modest burn and plastic surgery setup with limited obstetric facilities.

## MATERIALS AND METHODS

This is a retrospective study from 1998 to 2003 including 49 patients admitted in the burn and plastic surgery unit in tertiary referral center. In all patients the extent of burns was calculated by using the rule of 9's. A burn less than 10% in total body surface area (TBSA) was considered as a minor burn. A major burn was defined as a partial or full thickness burn affecting more than 10% TBSA with 10-19% being graded as moderately severe, 20-50 % as severe and more than 50% as critical. All patients received routine thermal injury care, including fluid resuscitation, wound care and nutritional support. Patients were treated with 1% topical silver sulphadiazine and systemic antibiotics

when indicated according to culture /sensitivity report. Wound care included hydrotherapy, desloughing, escharectomies, debridement and skin grafting. None of the patients underwent early excision and skin grafting. Nutritional support included daily intake of protein and calories according to the basal energy expenditure. Obstetric consultation was sought promptly on admission for each pregnant patient. Confirmation of pregnancy was done by history, physical examination, pregnancy test and ultrasonography of abdomen. State of foetus was recorded at admission and monitored. Maternal assessment included cause of burn, site, depth and extent of burn, burn related complications and maternal outcome. Foetal outcome was assessed in relation to gestational age, maternal TBSA involvement and complications.

## RESULTS

49 patients age ranging from 18 years to 35 years (mean age 23.08 years) were assessed. All were in good health with normal pregnancies at the time of injury. The percentage of TBSA varied from 8 % to 100% (mean 71.47%). Four patients had minor and 45 patients had major burns. Resuscitation was successful in all patients except 13 who sustained 100% burn and died within 48 hours of admission to hospital. The injuries were predominantly domestic and fire and hot liquids were major causative factors. Gestational age at the time of injury varied from 8 weeks to 34 weeks with 11 cases in first trimester, 21 in second trimester and 17 patients in last trimester [Table 1]. Thirty three mothers died after sustaining severe burns and foetal deaths occurred in 34 of the 49 cases. With increasing TBSA burn complication rate and maternal and foetal loss

increased. (Maternal mortality 21.42 % in < 50% burn to 88.23% in > 50%; Foetal mortality 28.57 % in < 50% burn to 88.23% in > 50%)

Maternal outcome in relation to TBSA and maternal complications are shown in Table 2. Maternal death was directly attributable to irreversible burn shock in 15 cases, Septicemia in 8 cases and respiratory complications including smoke inhalation pulmonary edema and embolism in 10 cases. In TBSA <50% only 3 out of 15 mothers died due to septicemia. In contrast when TBSA was >50% 30 out of 34 mothers died. Fetal death occurred in 34 of 49 (69.3%) cases and was related to the extent of maternal burn. Fetal loss during first trimester was high (8 out of 11 mothers aborted). Twenty one mothers were seen in second trimester and 6 fetuses survived and the 17 fetuses 6 survived burn injury to mothers in the last trimester. Only one mother delivered a premature baby during initial hospital admission. The remaining pregnancies were continued normally but no information was available on the ultimate outcome of pregnancies carried to the term when patients were discharged before delivery.

## DISCUSSION

The possibility of pregnancy must be considered when any woman of reproductive age sustains a burn injury. Although rare, an extensive burn during pregnancy is a serious complication. In this study, 12.29% of all women of reproductive age admitted with burns were pregnant. Because pregnancy tests were not done routinely the true incidence of pregnancy associated with burn injuries, especially in the first trimester, remains unknown. Therefore

**Table 1: Relationship between gestational age and foetal outcome**

	TBSA 1 - 10%			TBSA 10 - 19%			TBSA 20 - 49%			TBSA > 50%		
	1trim	2trim.	3trim.	1 trim	2 trim.	3 trim.	1 trim	2 trim.	3 trim.	1trim	2trim.	3trim.
Mother	1		3	1	1		3	5	1	7	15	12
Alive	1		3	1	1		1	4	1	1	2	1
Dead							2	1		6	13	11
Normal pregnancy	1		3	1	1		1	3	1	1	2	
Premature delivery												1
Abortion							2	1		6	13	3
Stillbirth								1				8

**Table 2: Maternal burn size, complications and outcome**

TBSA %	Number	Irreversible burn shock	Respiratory complications	Septicemia	Mother		Foetus	
					Alive	died	Alive	died
< 10%	4	Nil	Nil	Nil	4		4	
10 - 19%	2	Nil	Nil	Nil	2		2	
20 - 49%	9	Nil	Nil	3	6	3	5	4
> 50%	34	13	10	5	4	30	4	30
<b>Total</b>	<b>49</b>	<b>15</b>	<b>10</b>	<b>8</b>	<b>16</b>	<b>33</b>	<b>15</b>	<b>34</b>

all female burn patients of childbearing age should be tested for pregnancy unless the pregnancy is obvious.<sup>8</sup> There are specific physiological changes that occur during pregnancy that may have an impact after thermal injury on maternal and foetal well-being. Burns cause many maternal physiological changes, and places additional stress on systems that are already highly modified. Pregnancy is associated with hyperdynamic cardiovascular state. After burns there is increased capillary permeability and third space loss leading to hypovolemia, which may in turn lead to hypotension if the patient is inadequately resuscitated. This may lead to placental insufficiency, fetal ischemia, hypoxia and acidosis, as the placenta is a less preferred site. All these events lead to premature labor. Thus aggressive fluid resuscitation, upright posture and oxygen supplementation should be provided to the mothers even in the absence of smoke inhalation.<sup>7,9</sup>

There is a linear co-relationship between the maternal total percentage of burns and the probability of death.<sup>6,10</sup> In the present study, the fatality rate among patients with TBSA of 50% or more was 3.33 times the fatality rate among women with smaller burns. Percentage TBSA affected was therefore continues to be the best predictor of survival. Pregnancy as such does not pose any additional burden on the patients and their prognosis is same as non-burnt female.<sup>11,12</sup>

Fetal survival depends upon the gestational age, extent of maternal injury and maternal survival. In our study fetal survival during first trimester was 27.2 % in comparison with 28.5 % in second and 35.2 % in third trimester. There was a significant correlation between fetal complications and the maternal percentage of TBSA burn.<sup>7</sup> In this respect, fetal risk has been shown to increase with increasing

maternal TBSA burns. Most fetuses survive when the mother survives and remain free of severe complications such as sepsis, hypotension and hypoxia.<sup>12,13</sup> In the first trimester fetal loss is very high since 8 of the 11 mothers aborted. But out of these 11 mothers 7 were having burn >50% TBSA. Therefore it is directly related with the extent of maternal thermal injury. In the second trimester fetal loss largely depended on maternal survival and majority of the fetuses survived if the mother survived the thermal injury. In the third trimester fetal survival is dependent more on gestational age and less on maternal survival. It is important to establish the gestational period precisely by the menstrual history and foetal ultrasound examination at the time of the burn accident that in turn determine the choice of obstetric procedures and ultimate foetal outcome. Fetuses delivered before 24 weeks generally will not survive, while those delivered after 32 weeks will do well with modern neonatal intensive care if born without hypoxia or birth trauma. Ex-utero survival of fetuses between 24 weeks to 32 weeks is difficult to predict therefore when preterm labour occurs, pharmacological inhibition of labour should be considered. Tocolytic therapy may be of value in preventing premature delivery or as a temporary method of arresting labour while maternal homeostasis is restored.<sup>14</sup> The general scheme for the obstetric management of a pregnant burns patients is given in Table 3.

None of the patients in our series received tocolytic therapy and it should not be considered until the maternal injury has been stabilized and is contraindicated in patients with existing cardio-vascular or pulmonary disease.<sup>15</sup> Parenteral magnesium sulfate is a better choice because it has less vasodilator and metabolic effects.<sup>16</sup> Prevention of hypovolaemia and hypoxia is crucial since even a short

**Table 3: The general scheme for the management of pregnancy in burns patients**

<b>Total % burn</b>	<b>Age of gestation</b>		<b>Management</b>
< 30	First trimester		No obstetric interference
	Second trimester		No obstetric interference
	Third trimester	More than 36 wks	Induce labour / caesarian section
		Less than 36 wks	Conservative approach and monitoring of heart rate
30-50	First trimester		Foetal monitoring by ultrasound 3-4 wks
	Second trimester		Foetal monitoring every 3-4 wks. Tocolytic therapy
	Third trimester	More than 32 wks	Deliver foetus within 48 h
		Less than 36 wks	Careful foetal monitoring
50-70	First trimester		Terminate pregnancy
	Second trimester		Terminate pregnancy
	Third trimester	If baby is viable	Induce labour / caesarian section within 24h
		Intrauterine death	No active intervention up to 4 wks / monitoring of foetus of haemocoagulation factors
> 70	First trimester		No treatment
	Second trimester		No treatment
	Third trimester		Caesarian section as an emergency procedure at the earliest

period of maternal hypoxia can lead to precipitation of labour or fetal death. An additional factor is invasive burn wound sepsis, which may result in premature labour and adverse vasoactive effects. Thermally injured tissue produces prostaglandins and increases the synthesis of free arachidonic acid. Organisms in the burn wound contribute to these effects. This increases uterine contractility thus initiating premature labour and abortion.<sup>4, 17</sup> Obstetric considerations affect the choice of route and the timing of the delivery; spontaneous vaginal delivery is generally preferred even in presence of perineal burns. When there are obstetric indications for a caesarian section, this can be performed safely even when the lower abdominal wall is burnt. In a stable patient with less than 50% burn if the foetus is immature, premature labour could be suppressed and pregnancy allowed to continue until term. If burn wound exceeds 50% TBSA and foetus is at risk, obstetric intervention is indicated within the first 24 hours for high-risk third trimester pregnancies.<sup>18</sup> Other indications for obstetric intervention include progressive foetal distress or significant maternal complications.

Principles of management of a pregnant burns patient is not different from non pregnant burns patient.<sup>19</sup> It includes early and adequate resuscitation, combating sepsis by limited choice of antibiotics and local antibacterial agents, early surgical intervention, adequate nutrition and close co-operation with obstetric team.<sup>2, 6, 7, 12</sup> Prevention of hypovolaemic shock by adequate early fluid therapy is required to maintain the uterine blood flow, which in turn maintains foetus tissue  $pO_2$  levels within the normal range. It is recommended to maintain the mother's blood pressure within the normal range and a urine output of 30-60 ml/h. In cases of inhalation injury, ventilatory support may be required. Ventilatory support should be initiated when maternal  $pO_2$  is less than 60 mm Hg as inhaled carbon monoxide can cross the placental barrier to compete for binding sites on foetal hemoglobin, provoking foetal cardiac edema, and affecting cardiac development. Burns complicated by pregnancy considerably reduces use of the commonly used drugs like Chloramphenicol, Aminoglycoside Tetracycline Ciprofloxacin Vancomycin, Rifampicin, Sulpha drugs, Salicylates and Povidone iodine for systemic and topical treatment of burns.<sup>19</sup> The only antibiotics that can be considered safe in pregnancy are the penicillin and cephalosporins, other commonly used antibiotics (Imipenem, teicoplanin) should be avoided during pregnancy unless they are necessary and their advantages outweigh any possible risk. Under anaesthesia

it is important to avoid the use of ketamine, which increases the excitability of the myometrium and may trigger effective uterine contractions; also, when the pregnancy is near term and delivery is imminent, ketamine may cause respiratory depression in the neonate.<sup>19</sup> Similarly, depolarizing agents like succinylcholine should not be used as it causes hyperkalemia, instead a non-depolarizing muscle relaxant like curare or pancuronium is preferred.<sup>20</sup> Povidone-iodine is widely used for burns cleansing but if used for > 20% of burns large amounts of iodine can be absorbed through the burn wound and pass through the placenta to affect thyroid functions and cause metabolic acidosis. Sulpha drugs (silver sulphadiazine) are also potentially teratogenic if administered before the 14th week of pregnancy and lead to growth retardation and low birth weight if subsequently administered. Certain other uncommon topical antibiotics like Chloramphenicol, either in powder form or as an ointment in association with collagenase, are teratogenic in first trimester and responsible for gray baby syndrome in the third trimester. Gentamicin is capable of passing through the placenta after absorption through burns wounds and can lead to ototoxicity and nephrotoxicity if used after the 14th week of pregnancy. Rifamycin for local use is absolutely contraindicated in the last period of pregnancy because its interference with bilirubin metabolism can cause indirect hyperbilirubinaemia, with the risk of kernicterus. Salicylates exert an anti-prostaglandin action and may therefore have a protective effect on pregnancy. For the same reason they should not be used in the final period of pregnancy because they prolong gestation and delay spontaneous delivery.

Early coverage of the burns by tangential excision and split-thickness skin grafting facilitates healing of the wounds and minimizes septic complications thus improving maternal and foetal outcome.<sup>21, 22</sup> Wounds over the abdomen and breast should be treated first. Early healing of the abdominal wound favors pain-free stretching of the abdominal skin during the developing pregnancy, facilitates obstetric supervision of the growing foetus and performance of caesarian section if required. Early surgery and healing of the breast wound prevents infection and sloughing of nipples and permits subsequent breast-feeding.<sup>9, 19</sup>

Pregnant women with self-inflicted burns have more fatality rate as compared to unintentional burns. This difference in mortality may be attributable to the higher percentage of TBSA burned in suicidal burn patients compared with

women with accidental burns. Burns sustained by pregnant women occur in similar circumstances as non-pregnant women and most of them are in good health. Like all burn wound management, once the resuscitation is complete the goal is to combat sepsis, achieve early wound coverage and provide adequate nutrition during convalescence. Since the patient is recumbent for long period whenever possible left lateral position is recommended to prevent uterine compression of inferior vena cava.

Long-term follow-up of the patients with severe abdominal wall scarring showed that pregnancies could be accommodated normally without increasing the risk of abortion, premature labour or difficulty during labour. Scarred abdomen does not pose any difficulty for caesarean section.<sup>23, 24</sup> The best chance for pregnant burned patients lies in prevention or decreasing the extent of burn by counseling, prohibiting the use of explosive kerosene stoves and proper primary aid to burn victims. Plastic surgeons and obstetricians should work in close association for management of this difficult problem.<sup>25</sup> Emphasis should be on burn prevention.

## CONCLUSION

In general, pregnancy as such does not influence maternal outcome after thermal injury and best chance for foetal survival is to ensure maternal survival. Maternal survival is less likely if the burn wound exceeds 50% total body surface area. Thermal injury does increase the risk of spontaneous abortion and premature labour and foetal survival depends upon foetal maturity.

## REFERENCES

1. Mulla N, Albuquerque MD, Mex N. Labour following severe thermal burns. *Am J Obstet Gynecol* 1958;76:1338-41.
2. Matthews RN. Obstetric implications of burns in pregnancy. *Br J Obstet Gynaecol* 1982;89:603-9.
3. Schmitz JT. Pregnant patients with burns. *Am J Obstet Gynecol* 1971;110:57.
4. Stage HA. Severe burns in the pregnant patients. *Obstet Gynecol* 1973 42:259-62.
5. Stilwell JH. A major burn in early pregnancy with maternal survival and pregnancy progressing to term. *Br J Plast Surg* 1982;35:33-5.
6. Rayburn W, Smith B, Feller I, Varner M, Cruikshank D. Major burns during pregnancy: Effects on foetal well-being. *Obstet Gynecol* 1984;63:392-5.
7. Taylor JW, Plunkett GD, McManus WF, Pruitt BA. Thermal injury during pregnancy. *Obstet Gynecol* 1976;47:434-8.
8. Guo SS, Greenspoon JS, Kahn AM. Management of burn injuries during pregnancy. *Burns* 2001;27:394-7.
9. Bartle EJ, Sun JH, Wang XW. Burns in pregnancy. *J. Burn Care Rehabil* 1988;9:485-7.
10. Cheah SH, Sivanesaratnam V. Burns in pregnancy-maternal and foetal prognosis. *Aust. N. Z. J. Obstet. Gynaecol* 1989;29:143-5.
11. Akhtar MA., Mulawkar PM, Kulkarni HR. Burns in pregnancy: Effect on maternal and foetal outcomes. *Burns* 1994;20:351-5.
12. Amy BW, McManus WF, Goodwin CW, Mason A, Pruitt BA. Thermal injury in the pregnant patient. *Surg Gynecol Obstet* 1985;161:209-12.
13. Rode H, Millar AJ, Cywes S, Bloch CE, Boes EG, Theron EJ, Lodder JV, van der Merwe AE, de Kock M. Thermal injury in pregnancy-the neglected tragedy. *S Afr Med J* 1990;77:346-8.
14. Deitch EA, Rightmire DA, Clothier J, Blass N. Management of burns in pregnant women. *Surg. Gynecol Obstet* 1985;161:1-4.
15. Benedetti TJ. Maternal complications of parenteral beta sympathomimetic therapy for premature labor. *Am J Obstet Gynaecol* 1983;145:1-6.
16. Barden TP. Premature labour. In: RT Bolognese, RH Schwarz, Schneider J, editors. *Perinatal medicine, Management of high-risk fetus and neonate*. 2<sup>nd</sup> edn. Baltimore: Williams and Wilkins, co; 1982. p. 240-303.
17. Herndon DN, Abston S, Stein MD. Increased thromboxane B<sub>2</sub> levels in the plasma of burned and septic burned patients. *Surg Gynaecol Obstet* 1984;159:210-3.
18. Ullmann Y, Blumenfeld Z, Hakim M, Mahoul I, Sujov P, Peled IJ. Urgent delivery, the treatment of choice in term pregnant women with extended burn injury. *Burns* 1997;23:157-9.
19. Napoli B, D'Arpa N, Masellis M, Graziano R. Burns in pregnancy *Annals of Burns and Fire Disasters* – vol XIII-n° 1 - March 2000
20. Gronert GA, Theye RA. Pathophysiology of hyperkalemia induced by succinylcholine. *Anesthesiology* 1975;43:89-99.
21. Mabrouk AR, El-Feky AEH. Burns during pregnancy: A gloomy outcome. *Burns* 1997;23:596-60.
22. Prasanna M, Singh K. Early burn wound excision in 'major' burns with 'pregnancy': a preliminary report. *Burns* 1996;3:234-37.
23. Kitzmiller WJ, Neale HW, Warden GD, Smith D. The effect of full-thickness abdominal wall burns sustained during childhood on subsequent childbearing ability. *Ann Plast Surg* 1998;40:111-3.
24. McCauley RL, Stenberg BA, Phillips LG, Blackwell SJ, Robson MC. Long-term assessment of the effects of circumferential truncal burns in pediatric patients on subsequent pregnancies. *J Burn Care Rehabil* 1991;12:51-3.
25. El-Gallal ARS., Yousef SM. burns in pregnancy: a ten-year review of admitted patients *Annals of Burns and Fire Disasters*-vol. XV-n. 3 September 2002.