

Antisteroidogenic Activity of Ethanol Extract of *Ammania baccifera* (L.) Whole Plant in Female Albino Mice Ovaries

RAMAIYAN DHANAPAL, SUBRAMANI KAVIMANI, VRUSHABENDRA SWAMY BHYRAPUR MATHA, MALAYA GUPTA and SANATH KUMAR BASU

Department of Pharmaceutics (R.D.); Department of Pharmacology (V.S.B.M.), Rural College of Pharmacy, Devanahalli, Bangalore, Karnataka; Mother Theresa Research Institute of Health Sciences, Pondicherry (S.K.); Division of Pharmacology (M.G.); Division of Pharmaceutics (S.K.B.), Javadpur University, Kolkata, West Bengal, India.

Received March 22, 2005; Revised June 9, 2005; Accepted June 11, 2005

This paper is available online at <http://ijpt.iuims.ac.ir>

ABSTRACT

Ethanol (90%) extract of *Ammania baccifera* (L.) whole plant (EEAB) was evaluated for possible antisteroidogenic activity in mature female mice ovaries. The ethanol extract at the doses of 100, 200 and 400 mg/kg body weight (i.p) arrested the normal estrus cycle at dioestrus phase and significantly decreased weight of ovaries. The cholesterol and ascorbic acid content in ovaries were significantly elevated in treated mice. The extract also significantly inhibited the activity of Δ^5 - 3β -hydroxy steroid dehydrogenase (Δ^5 - 3β -HSD) and Glucose-6-phosphate dehydrogenase (G-6-PD), the two key enzymes involved in ovarian steroidogenesis. Results of this study suggested that the ethanol extract of whole plant of *Ammania baccifera* (L.) acts as an antisteroidogenic agent.

Keywords: *Ammania baccifera*, Ovarian steroidogenesis, Δ^5 - 3β -HSD, G-6-PD, Medicinal plant

Ammonia baccifera Linn (Lythraceae) is a glabrous, erect branching herb, found as weed in rice-fields and marshy localities throughout India. The leaves are acrid and used in the treatment of rheumatic pain, as laxative, rubifacient and external remedy for ring worm [1]. This plant was found to possess hypothermic, hypertensive, antiurolithiasis, antibacterial and CNS depressant activities [2-4]. Steroids triterpenes, coumarins, flavonol and tannins were previously isolated from various parts of the plant [5, 6]. It has come to our notice that the rural people of Tamilnadu use this plant to produce sterility in animals. In the present study, we evaluated the antisteroidogenic activity of the ethanol extract of *Ammania baccifera* (L.) whole plant in female mice ovaries.

MATERIALS AND METHODS

Plant Material

The whole plant of *Ammonia baccifera* (L.) was collected from Trichy, Tamilnadu, India and was identified and authenticated by Prof. Sri. Ganesh, Botanist Madura College, Madurai, Tamilnadu. A voucher specimen MG-3 has been kept in our laboratory for future reference. The whole plant was dried under shade, powdered by a mechanical grinder and was passed through 40-

mesh sieve and stored in airtight container for further use.

Preparation of Extract

About 1 kg of the powdered plant material was successively extracted using petroleum ether (40-60°C), chloroform, and then ethanol (90%) in a Soxhlet extraction apparatus. The various extracts were concentrated and the traces of the solvent were completely removed under reduced pressure and were stored in a vacuum desiccator for further use. The yield was found to be petroleum ether extract (0.9%); chloroform extract (1.7%) and ethanol extract (3.6%) w/w with respect to dried powder. Preliminary qualitative chemical tests indicated the presence of steroids, triterpenoids, flavonoids and tannins. Further investigation was carried out using the ethanol extract.

Animals

Adult female albino mice of Swiss strain 20 ± 2 g were acclimatized to normal environmental condition in the laboratory for one week and given a standard pellet diet (Hindustan Lever) and water *ad libitum*.

The experiment was performed under the guidance of the Ethical Committee, Javadpur University, Kolkata-32.




Fig 1. Effect of ethanol extract of whole plant of *Ammania baccifera* (L.) (EEAB) on content of ascorbic acid and cholesterol in female mice ovaries.




Fig 2. Effect of ethanol extract of whole plant of *Ammania baccifera* (L.) (EEAB) on the activities of Δ^5 -3 β -HSD in matured female mice ovarian tissues.

Experimental Design

Treatment of Animals. The mice showing four consecutive normal oestrus cycle were then divided into five groups (n=10). In their proestrus phase the first group received normal saline (5 ml/kg/day) whereas group 2, group 3, group 4, and group 5, received vehicle (Propylene glycol 5 ml/kg body weight), ethanol extract of whole plant of *Ammania baccifera* (L.) (EEAB) (100, 200 and 400 mg/kg/body weight) intraperitoneally respectively for every alternate days for 18 days. Oestrus cycle was observed everyday by microscopic examination of vaginal smear. On the 19th day the mice were killed by cervical location, 24 hours after the final does and after 18 hours fasting. Ovaries were dissected out, weighed and kept on ice for biochemical estimations.

Biochemical Estimation. Ovarian tissues about 3 mg weight, were carefully homogenized in Potter Elvehjem homogenizer using chloroform: ethanol mixture (2:1) and non-polar part was extracted out and total cholesterol content was estimated according to the method of Kingsley and Roscoe [7].

About 5 mg of tissue was homogenized in Potter Elvehjem homogenizer using 2.5 ml ice cold 5% metaphosphoric acid and centrifuged for 20 min at 355 rpm then ascorbic acid content was calculated [8].

About 2 mg of ovarian tissue was homogenized in Potter Elvehjem homogenizer using 1 ml of normal saline and 1 ml of 0.1 M phosphate buffer (pH 7.4) and

centrifuged. The activity of Δ^5 -3 β -HSD was estimated as described by Rabin et al [9].

About 3 mg of ovarian tissue was again homogenized in Potter Elvehjem homogenizer using 0.5M Tris-HCL (pH 8.3) and centrifuged. The activity of G-6-PD was estimated as described Iohr and Waller [10].

Protein was estimated with Folin's phenol reagent and the activities of enzymes were expressed in unit per mg of protein as described by Lower et al [11].

Statistical Analysis

Statistical analysis was done by using Student' t-test.

RESULTS

Ethanol extract of whole plant of *Ammania baccifera* (L.) (EEAB) arrested normal spelling cycle at dioestrus phase at doses of 100, 200 and 400 mg/kg body weight after 13,10 and 6 days of treatment respectively. It was found that the EEAB significantly reduced the wet weight of ovaries in a dose dependent manner ($p < 0.05$ by 100mg and $p < 0.01$ by both 200 and 400 mg). The ethanol extract of whole plant of *Ammania baccifera* (L.) (EEAB) at 100, 200 and 400 mg/kg body weight significantly increased the level of total cholesterol and ascorbic acid contents of ovaries in treated mice. The activities of Δ^5 -3 β -HSD were inhibited significantly ($p < 0.05$ by 100 mg and $p < 0.001$ by both

Table 1. Effect of ethanol extract of whole plant of *Ammania baccifera* (L.) (EEAB) on weight of ovaries, content of ascorbic acid, cholesterol and the activities of G-6-PD and Δ^5 -3 β -HSD in matured female mice ovarian tissues.

Design of Treatment	No. of days	Weight of ovary (mg)	Ascorbic acid (g/mg of ovary)	Cholesterol (g/mg of ovary)	G-6-PD (U/mg of protein)	Δ^5 -3 β -HSD (U/mg of protein)
Saline (5 ml/kg b.w, i.p)	18	14 ± 1.2	83 ± 2.2	50 ± 1.7	4.1 ± 0.05	1.08 ± 0.04
Vehicle (PG) (5 ml/kg b.w, i.p)	18	13 ± 1.8	80 ± 3.8	53 ± 0.4	4.2 ± 0.02	1.15 ± 0.06
EEAB (100 mg b.w, i.p)	18	10.0 ± 0.4*	109 ± 2.6*	72 ± 5.7*	3.5 ± 0.06**	0.9 ± 0.03*
EEAB (200 mg b.w, i.p)	18	9.3 ± 0.3**	123 ± 3.4**	93 ± 8.3***	3.3 ± 0.03***	0.8 ± 0.04***
EEAB (400 mg b.w, i.p)	18	7.3 ± 0.3**	169 ± 5.4**	148 ± 6.6***	3.0 ± 0.04***	0.6 ± 0.02***

PG=Propylene glycol, b.w = body weight, i.p = intraperitoneal, EEAB=Ethanol extract of *Ammania baccifera* (L.) whole plant.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ significantly different from vehicle control.

200 and 400 mg). Similarly, the activities of G-6-PD were inhibited significantly ($p < 0.01$ by 100 mg and $p < 0.001$ by both 200 and 400 mg) by all the doses of whole plant of *Ammania baccifera* (L.) (EEAB). (Table 1, Fig 1 and Fig 2).

DISCUSSION AND CONCLUSION

The minimum activity of steroid hormones has been reported to occur in the dioestrus stage. [12-14]. This was associated with an elevation in the level of cholesterol as well as ascorbic acid content which serve as precursor for the synthesis of steroid hormones in ovaries, suggesting, that cholesterol and ascorbic acid were not utilized [15-**Error! Reference source not found.**]. The steroidogenesis is under the physiological control of two enzymes [19, 20].

The ethanol extract of whole plant of *Ammania baccifera* (L.) (EEAB) at treated doses arrested normal oestrus cycle at dioestrus phase and also elevated the ascorbic acid and cholesterol content and significantly reduced the activity of Δ^5 -3 β -hydroxy steroid dehydrogenase (Δ^5 -3 β -HSD) and Glucose-6-phosphate dehydrogenase (G-6-PD) in a dose dependent manner. These results revealed that EEAB produced ovarian malfunction by altering substrate and enzyme activities. Preliminary phytochemical tests indicated the presence of flavonoids in the ethanol extract of whole plant of *Ammania baccifera* (L.) Since various flavonoids have been reported [21, 22] to possess antifertility activity, the antisteroidogenic property of the ethanol extract of whole plant of *Ammania baccifera* (L.) might be due to the presence of such compounds.

From the present investigation it may be concluded that the ethanol extract of whole plant of *Ammania baccifera* (L.) may be considered as an antisteroidogenic agent.

ACKNOWLEDGEMENTS

Authors are sincerely thankful to Prof. Dr. UK Mazumdar, Division of Pharmaceutical Chemistry, Jadavpur University, Kolkata, for providing computer facilities for the help in the preparation of manuscript.

REFERENCES

- Kirtikar K R and Basu BD. Indian Medicinal Plants. Vol II, 2nd ed. Dehradun, India Bishen Mahendra Pal Singh, New Delhi. 1972. p. 1071.
- Dhar MN, Dhavan BN, Mehrotra BN, Srimal RC, Tandon JS. Screening of Indian Medicinal Plants for Biological Activities. *Indian J Exp Biol* 1973;**11**:43-54.
- Bharathi K, Srinivasan KK. Evaluation of *Ammania Baccifera* Linn. for antiurolithic activity in albino rats. *Indian J Exp Boil* 1994;**32**(5):311-3.
- Al-Sharma A, Mitschar LA. Comprehensive surgery of indigenous Iraqi plants for potential economic value. 1. screening results of 327 species for alkaloids. *J Nat Prod* 1979;**42**(6):633-42.
- Atal CK, Srivastava JB, Wali BK, Chakravarty RB, Dhawan BN, Rastogi RP. Screening of Indian Medicinal Plant for Biological activities. *Indian J Experimental Biology* 1978;**16**:330-49.
- Thakkar SM, Deshmukh VK, Saoji AN, Duragkar NJ. Herbal drugs for urinary stones. *J Indian Chem Soc* 1986;**64**(6):619-20.
- Kingsley GR, Roscoe RS. Determination of free and total cholesterol by direct chloroform extraction. *J Biol Chem* 1949;**180**:315-28.
- Omye ST, Turnbull JD, Souberlich HE. Selected methods for the determination of ascorbic acid in animal cells, tissues and fluids. In: DB McCormick and WrightLD. (Eds) *Methods Enzymology*. Academic press. New York. 1979. p. 623-47.
- Rabin BL, Leipsner G, Deane HW. A rapid sensitive assay procedure for adrenal steroid 3 OL Dehydrogenase activities. *Endocrinology* 1961;**68**:619-25.
- Lohr LC, Waller HD. Glucose-6-Phosphate dehydrogenase In: Berg Meyer H.U. *Methods of enzymatic analysis Verlag Chemie*, Florida. 1974. p. 636-46.
- Lowry OH, Rosenbrough MJ, Farm AL, Randell RJ. Protein Measurement with folin phenol reagent. *J Biol Chem* 1951;**193**:365-74.
- Wiest WG, Wilcox RB, Kirschbuan HT. Rat ovarian 20-Hydroxy steroid dehydrogenase: effect of estrogens and pituitary gonadotrophins. *Endocrinology* 1963;**74**:588-90.
- Ramirez BD, McCamm SM. Fluctuations on plasma luteinizing hormone concentrations during estrous cycle of the rat. *Endocrinology* 1964;**74**:814-16.
- Andreson RR, MsShan WH. Leutinizing Hormone in levels in pig, cow and rat blood plasma during estrous cycle. *Endocrinology* 1966;**76**:976-82.
- Tamooki B, Pincus G. Biogenesis of progesterone in ovarian tissues. *Endocrinology* 1961;**69**:527-33.
- Guillemin R, Sakiz E. Quantitative study of responses to LH after hypophysectomy in the ovarian ascorbic acid depletion test: effect of prolactin. *Endocrinology* 1963;**72**:813-816.
- Krum AA, Morris MD, Bennet LL. Role of cholesterol in the in-vivo biosynthesis of adrenal steroids by dog. *Endocrinology* 1964;**4**:543-7.
- Baille AH, Ferguson MM and Hardt MC. In: *Developments in Steroid Biochemistry*, New York, *Academic Press*. 1960. p. 10-25.
- Armstrong, DG. 3-B-Hydroxy A-5-steroid dehydrogenase activity in the rapidly growing ovarian follicles of the domestic fowl (*Gallus domesticus*). *J Endocrinology* 1982;**93**:415-21.
- Suzuki S, Endo Y, Tanaka S, Iizuka R. Indirect immunofluorescence studies on the steroid producing activity of hamster ova. *Am J Obstet Gynecol* 1984;**148**:76-85.
- Hiremath SR, Rao SH. Antifertility efficacy of the plant *Striga lutea* (scrophulariaceae). *Contraception* 1990;**42**:467.
- Khanna U, Choudhury RR. Antifertility screening of plants, part 1. Investigation of *Butea Monosperma* (Lam) Kutze. *Indian J Med Res* 1968;**56**:1575.

Address correspondence to: Dr. R. Dhanapal, Department of Pharmaceutics, Devanahalli, Bangalore, Karnataka, India.
E-mail: dhanapal2k2@sify.com
