

Original Article

Identification of messenger RNA of fetoplacental source in maternal plasma of women with normal pregnancies and pregnancies with intrauterine growth restriction

Ayala Ramírez, Paola^{*b}; García Robles, Reggie^b; Rojas, Juan Diego^{ca}; Bermúdez, Martha^b; Bernal, Jaime^b

^a Department of Obstetrics and Gynecology Pontificia Universidad Javeriana-Hospital Universitario San Ignacio, Bogotá, D.C., Colombia.

^b Instituto de Genética Humana, Pontificia Universidad Javeriana, Bogotá, D.C., Colombia.

^c Gynecologist physician. Unit of Feto-maternal Medicine

Article info

Article history:

Received 12 september 2011

Received in revised form 25 november 2011

Accepted 17 february 2012

Available online 30 september 2012

Keywords:

RCIU, Messenger RNA, plasma, pregnancy complications, human placental lactogen, Chain Reaction Reverse Transcriptase Polymerase.

ABSTRACT

Objective: to quantify placenta-specific RNA in plasma of women carrying foetuses with intrauterine growth restriction and pregnant women with normal pregnancies.

Materials and methods: 8 pregnant women with foetuses with intrauterine growth restriction were studied as well as 18 women with uncomplicated pregnancies in the third pregnancy trimester. Total free RNA was quantified in maternal plasma by spectrophotometry and the gene expression of hPL (Human Placental Lactogen) at the messenger RNA level through technical Real Time-Chain Reaction Polymerase.

Results: plasma RNA of fetoplacental origin was successfully detected in 100% of pregnant women. There were no statistically significant differences between the values of total RNA extracted from plasma ($p = 0.5975$) nor in the messenger RNA expression of hPL gene ($p = 0.5785$) between cases and controls.

Conclusion: messenger RNA of fetoplacental origin can be detected in maternal plasma during pregnancy.

INTRODUCTION

Intrauterine Growth Restriction (IUGR) is defined as the clinical diagnosis wherein the foetuses do not reach its full growth potential and the final outcome is a decrease in body weight, being below the 10th percentile for gestational age and sex according to growth charts¹. The infant may have a small complexion or because of factors in the mother, in the placenta or foetuses². IUGR is a major burden on perinatal and neonatal morbidity and mortality¹. In addition, children with IUGR subsequently may evolve with changes in growth and neurocognitive development³. The association of low birth weight (LBW) with subsequent development of adult diseases such as insulin resistance and cardiovascular disease

has also been studied, this finding has been termed “fetal programming”⁴. Biomarkers in maternal serum have been shown to have predictive power and sub-optimal sensitivity for fetal growth assessment⁵. Due to the limitations of current techniques, it is necessary to develop more accurate methods for monitoring fetal growth⁶. In 1997 Lo et al., managed to isolate free DNA (deoxyribonucleic acid) circulating in plasma of pregnant women⁷. These studies opened a new possibility of noninvasive prenatal diagnosis where neither the foetuses nor the mother are in any danger⁸. Because it is often not possible to determine the source of DNA found in plasma of pregnant women and can be confused with the maternal DNA, the question of whether RNA could be isolated from fetal-placental origin in maternal plasma and thus determine the expression of specific genes in the placenta was formulated. Poon et al., were the first to report the presence of fetal RNA in the plasma of pregnant woman⁹. Numerous studies have analyzed the gene expression at level of messenger RNA (mRNA) extracted from maternal plasma in fetal pathologies such as trisomy^{10,11}, congenital cardiopathies¹² and maternal complications such as: pree-

*Corresponding Author.

E-mail Address : payala@javeriana.edu.co (Ayala P), rgarcia@javeriana.edu.co (García R), jdiegorojas@yahoo.com (Rojas JD), martha.bermudez@javeriana.edu.co (Bermudez M), jebernal@javeriana.edu.co (Bernal J)

clampsia^{13,14}, RCIU⁶ and previous percreta placenta¹⁵, stating that the use of genetic markers at the level of mRNA of fetal-placental origin circulating in maternal plasma opens up new possibilities for strategies in early detection of fetal pathological conditions, noninvasive prenatal diagnosis, determination of risk for developing complications of pregnancy and pathology severity. Human placental lactogen (hPL) is produced in the placenta and acts as an immunosuppressant inducing tolerance and gestational fetal growth factor². Previous studies have linked low levels of mRNA in maternal plasma of hPL gene in pregnancies complicated by previous placenta¹⁵ and preeclampsia^{14,16}. These studies conclude that real time-PCR is a sensitive method to monitor changes in mRNA levels resulting from apoptotic effects in the placenta and to evaluate the conditions of villous trophoblast invasion¹⁵. The aim of this study was to quantify total RNA by spectrophotometry and hPL gene expression at the level of free mRNA in maternal plasma by RT-qPCR technique in pregnancies that progressed with IUGR and pregnant women with uncomplicated pregnancies and normal foetuses.

MATERIALS AND METHODS

An analytical study was conducted in pregnant women in third trimester of pregnancy whose pregnancies progressed with IUGR and in pregnant women with uncomplicated pregnancy healthy foetuses, assisted by the Department of Gynecology and Obstetrics, of the Hospital Universitario San Ignacio in Bogotá between November 2009 and December 2010. Inclusion criteria were patients with diagnosis by conventional ultrasound or color Doppler ultrasound in fetal, placental and both uterine arteries of IUGR (weight less than 10 percentile for gestational age) and with no apparent etiology, we excluded patients with ultrasonographic diagnosis of IUGR and small newborn for gestational age with known cause, foetuses / newborn with major fetal malformations or diagnostic impression of genetic disease, maternal prothrombotic conditions and pregnant women who expressed their wish not to participate in the study.

The study was approved by the Committee for Research and Bioethics of the Medical Faculty of Pontificia Javeriana University and fulfilled the guidelines established in the Helsinki declaration of 1975 as amended in 2004. All participants filled out a voluntary

	Temperature	Time
Denaturation	95°C	10 minutes
45 Cycles		
Denaturation	95°C	10 seconds
Annealing	62°C	10 seconds
Elongation	72°C	10 seconds
Lecture	72°C	single
Melting curve	95°C	0 seconds
	60°C	1 minute
	95°C	0 seconds
Cooling	40°C	30 seconds

Table 1. PCR amplification conditions of hPL gene.

informed consent.

Sample taking and processing: Two samples of peripheral blood by venipuncture (5 cc) in tubes containing EDTA anticoagulant were taken. The samples were immediately centrifuged at 3000 rpm for 15 minutes and the plasma (supernatant) was transferred to a 15 ml Falcon tube. 1.6 ml of plasma was mixed with 2 ml of Trizol LS (Invitrogen) and stored at -40 ° C until processed.

RNA extraction: extraction was performed in plasma using the protocol recommended by Ng et al.¹⁷ adding 5U of DNase (Epicentre). Total RNA was stored at -40 ° C. The total RNA concentration was determined at 260 nm using GeneQuant Pro (Amersham Bioscience).

Reverse transcription and PCR in Real time (qPCR-RT): reverse transcription of total RNA to cDNA was performed using the SuperScriptIII kit (Invitrogen, Carlsbad, CA, USA) in a reaction with 20 ul final volume containing 1X buffer, 0.5 mM dNTPs, 50 ng of random primer, 0005 M DTT, 40U of RNase OUT, 200U

	Cases (n=8)		Controls (n=21)		p value
	Median	Min-max range	Median	Min-max range	
Maternal age (years)	26,5	20-34	28	20-38	0,3649*
Gestational age at delivery (weeks)	37,5	31-40	38	37-42	0,1052*
Infant birth weight (grams)	2230	895-2775	2855	2470-3460	0,0003*
Placental weight (grams)	460	290-475	470	390-610	0,3731*
Birth weight/placenta weight ratio	4,65	3,09-5,78	6,12	5,49-8,09	0,002*
	n	%	n	%	
Primiparous					
Yes	3	37,5	3	14,29	0,0934**
No	5	62,5	18	85,71	
Socioeconomic status[§]					
1 y 2	5	62,5	11	52,38	0,253***
3 y 4	2	25	9	42,86	
5 y 6	1	12,5	0	0,00	
1 y 2 y 3	6	75	19	90,48	0,057**
4 y 5 y 6	2	25	1	4,76	
Sex					
Female	6	75	14	66,67	1,000**
Male	2	25	7	33,33	

Table 2. Demographic characteristics of the population, * Mann-Withney test. ** Binomial test. *** Fisher test. § A pregnant woman in the control group did not know her socioeconomic information

of SuperScript and 7 ul of RNA, with subsequent digestion with RNaseH 2U, following the manufacturer's recommendations. The cDNA was stored at -20°C until amplification. LigthCycler (Roche) thermal cycler was used using SYBR Green chemistry (LigthCycler FastStart DNA Master SYBR Green I kit, Roche) and the Lightcycler 4.1 software for absolute quantification (Roche Diagnostics). PCR was performed in a reaction volume of 20 ul with 7 ul of sample and 0.25 uM primer, following the manufacturer's recommendations. The program conditions of the PCR are shown in Table 1. The sequences of the primers were: 5' CATGACTCC-CAGACCTCCTTC3' (sense) and 5'TGCGGAGCAGCTCTA-GATTG3' (antisense); it confirmed that these primers do not amplify DNA with the Primer3 software¹⁸ and performing DNA amplification which did not show any fragment. The standard curve was performed as of the PCR product of purified hPL with the Wizard® SV Gel kit and PCR Clean-Up System (Promega), the number of copies was calculated according to Overbergh et al.¹⁹. Serial dilutions were performed with a range from 1X107 to 1X101 copies. Each sample was analyzed in duplicate and each analysis was mounted in a negative control in which cDNA instead of water was placed.

Statistical Analysis: analysis was performed by calculating median and ranges. We performed the hypothesis testing approach to determine whether there is association between the levels of total RNA and mRNA of hPL in plasma of normal controls and patients with IUGR using the Mann-Whitney test with Stata 9.1.

RESULTS

In total, the studied collected data from 12 cases and 22 controls, but excluded 4 cases and 1 control because the newborn had some exclusion criteria. The case group was comprised of 8 women pregnant in the third trimester of pregnancy with a diagnosis of idiopathic IUGR. Table 2 shows the demographic characteristics of the population. The variables of birth weight and birth weight / placental weight index showed statistically significant differences ($p = 0.0003$ and $p = 0.002$, respectively). With respect to the classification of IUGR, 50% of cases had an asymmetric IUGR and 50% had symmetric IUGR. Additionally, 3 patients had mild IUGR, 5 moderate to severe and 2 patients had abnormal Doppler.

RNA extract was achieved from 100% of plasma samples. The results of the quantification of total RNA from plasma and hPL mRNA expression are described in Table 3. There were no statistically significant differences in the concentration of total RNA or mRNA expression of hPL ($p = 0.5975$ and 0.5785 , respectively).

DISCUSSION

In this study, RNA was extracted from plasma of 100% of the samples and it was possible to quantify mRNA expression of the gene hPL by real time-PCR. These findings reproduce the results previously reported in the literature which shows that the placenta is an important organ that releases fetal RNA in maternal plasma^{17,20,21}. The amount of total RNA obtained from plasma in controls is lower than that reported by Maron et al., who obtained plasma RNA levels between 3 and 780 mg of 3 individuals in the third quarter²², however it should be noted that Maron et al study took 10 cc of whole blood and in this study we worked with 5 cc. The median levels of hPL gene mRNA in controls are higher than those reported by Tsui et al.²³ (n=5), Chiu et al.²⁰ (n=6), Okazaki et al.²¹ (n=75) and Ng et al.¹⁷ (n=10), who reported levels in copies /ml of 14.707, 9.900, 3.698 and 122.225 (data inferred from the article) respectively and less than the data reported by Farina et al.¹⁴ (n=30) who reported a median of 2,754,862. No statistically significant differences in levels of total RNA and mRNA hPL gene between cases and controls were found.

This is the first study which evaluated the mRNA levels of hPL gene in patients with a diagnosis of idiopathic IUGR. In patients with preeclampsia have been found that mRNA levels of hPL gene are significantly lower in both pregnant in the first quarter¹⁴ as in the second and third trimesters¹⁶, these studies show that it is possible to monitor placental status in maternal plasma. The absence of statistically significant differences in our study could be due to sample size, so it is recommended expanding it. Another possibility could be that the absence of differences in placental weight between the groups and because hPL is a marker of placental weight, were not significant differences. The results of hPL protein expression in placentas with IUGR show no statistically significant differences with healthy placentas. Although it has been reported a decrease in protein levels of hPL in the plasma of women with significant IUGR differences⁵, we were unable to find these differences at the level of mRNA, which could be due to the slow release of mRNA compared to protein-fetal placental origin. Moreover, development of placental mRNA markers that can be detected in plasma represents a breakthrough in the use of molecules that could be used in all pregnant women regardless of fetal sex using DNA limiting factor. Furthermore Zong et. al., hypothesizes that the quantitative assessment of fetal DNA in maternal plasma together with the measurement of mRNA from different transcripts could help improve identification of women at risk of developing preeclampsia²⁴, which may also be proposed for IUGR.

	Cases (n=8)		Controls (n=21)		p value
	Median	Min-max range	Median	Min-max range	
Total RNA in plasma (µg)	1.75	0,54-9,48	1.86	0,3-11,22	0,5975
mRNA of hPL gene (copies/ml of plasma)	311.500	890-12.700.000	419.500	7.970-20.600.000	0,5785

Table 3. Quantification of total RNA and absolute quantification of hPL gene in pregnant women with foetuses with IUGR and normal pregnancy.

CONCLUSIONS

mRNA of fetoplacental origin was detected in plasma of Colombian pregnant women in third trimester of pregnancy. The data presented demonstrate that it is possible to profile gene expression in the placenta by analyzing maternal plasma. The origin and release mechanisms of these molecules have not been completely elucidated and their understanding is likely to generate knowledge about the pathophysiology of diseases such as preeclampsia and IUGR. Additional studies are needed measuring mRNA levels in plasma of genes involved in the pathophysiology of the disease evaluated in normal and abnormal pregnancies to establish the feasibility of this proposal for screening and diagnosis of pregnancy complications in the clinical routine.

REFERENCES

1. Molina LM, Barbosa RH. Retardo de crecimiento intrauterino (RCIU) y sus alteraciones bioquímicas. *Nova*. 2005(003):88-94.
2. Kjell Haram ES, Ole Myking. Groeth Restriction: Etiology, Maternal and Neonatal Outcome. *Cirr Wom Health Rev*. 2007;3:145-60.
3. Kramer MS. The epidemiology of adverse pregnancy outcomes: an overview. *J Nutr*. 2003;133(5 Suppl 2):1592S-6S.
4. Cikot RJ, Steegers-Theunissen RP, Thomas CM, de Boo TM, Merkus HM, Steegers EA. Longitudinal vitamin and homocysteine levels in normal pregnancy. *Br J Nutr*. 2001:49-58.
5. Markestad T, Bergsjø P, Aakvaag A, Lie RT, Jacobsen G, Hoffman HJ, et al. Prediction of fetal growth based on maternal serum concentrations of human chorionic gonadotropin, human placental lactogen and estradiol. *Acta Obstet Gynecol Scand Suppl*. 1997;165:50-5.
6. Pang WW, Tsui MH, Sahota D, Leung TY, Lau TK, Lo YM, et al. A strategy for identifying circulating placental RNA markers for fetal growth assessment. *Prenat Diagn*. 2009;29(5):495-504.
7. Lo YM, Corbetta N, Chamberlain PF, Rai V, Sargent IL, Redman CW, et al. Presence of fetal DNA in maternal plasma and serum. *Lancet*. 1997:485-7.
8. Bianchi DW. Fetal cells in the maternal circulation: feasibility for prenatal diagnosis. *Br J Haematol*. 1999:574-83.
9. Poon LL, Leung TN, Lau TK, Lo YM. Presence of fetal RNA in maternal plasma. *Clin Chem*. 2000;46(11):1832-4.
10. Tsui NB, Akolekar R, Chiu RW, Chow KC, Leung TY, Lau TK, et al. Synergy of total PLAC4 RNA concentration and measurement of the RNA single-nucleotide polymorphism allelic ratio for the noninvasive prenatal detection of trisomy 21. *Clin Chem*. 2010:73-81.
11. Tsui NB, Wong BC, Leung TY, Lau TK, Chiu RW, Lo YM. Non-invasive prenatal detection of fetal trisomy 18 by RNA-SNP allelic ratio analysis using maternal plasma SERPINB2 mRNA: a feasibility study. *Prenat Diagn*. 2009;29(11):1031-7.
12. Arcelli D, Farina A, Cappuzzello C, Bresin A, De Sanctis P, Perolo A, et al. Identification of circulating placental mRNA in maternal blood of pregnancies affected with fetal congenital heart diseases at the second trimester of pregnancy: implications for early molecular screening. *Prenat Diagn*. 2010;30(3):229-34.
13. Shimizu H, Sekizawa A, Purwosunu Y, Nakamura M, Farina A, Rizzo N, et al. PP13 mRNA expression in the cellular component of maternal blood as a marker for preeclampsia. *Prenat Diagn*. 2009;29(13):1231-6.
14. Farina A, Sekizawa A, Purwosunu Y, Rizzo N, Banzola I, Concu M, et al. Quantitative distribution of a panel of circulating mRNA in preeclampsia versus controls. *Prenat Diagn*. 2006;26(12):1115-20.
15. Masuzaki H, Miura K, Yoshiura K, Yamasaki K, Miura S, Yoshimura S, et al. Placental mRNA in maternal plasma and its clinical application to the evaluation of placental status in a pregnant woman with placenta previa-percreta. *Clin Chem*. 2005:923-5.
16. Schmidt M, Hoffmann B, Kimmig R, Kasimir-Bauer S. mRNA of placental origin in maternal serum of women with normal and preeclamptic pregnancies. *Fetal Diagn Ther*. 2009:269-76.
17. Ng EK, Tsui NB, Lau TK, Leung TN, Chiu RW, Panesar NS, et al. mRNA of placental origin is readily detectable in maternal plasma. *Proc Natl Acad Sci* 2003:4748-53.
18. Rozen S, Skaletsky H. Primer3 on the WWW for general users and for biologist programmers. *Methods Mol Biol*. 2000;132:365-86.
19. Overbergh L, Valckx D, Waer M, Mathieu C. Quantification of murine cytokine mRNAs using real time quantitative reverse transcriptase PCR. *Cytokine*. 1999 Academic Press.; 1999:305-12.
20. Chiu RW, Lui WB, Cheung MC, Kumta N, Farina A, Banzola I, et al. Time profile of appearance and disappearance of circulating placenta-derived mRNA in maternal plasma. *Clin Chem*. 2006:313-6.
21. Okazaki S, Sekizawa A, Purwosunu Y, Iwasaki M, Farina A, Okai T. Measurement of mRNA of trophoblast-specific genes in cellular and plasma components of maternal blood. *J Med Genet*. England; 2006:e47.
22. Maron JL, Johnson KL, Slonim D, Lai CQ, Ramoni M, Alterovitz G, et al. Gene expression analysis in pregnant women and their infants identifies unique fetal biomarkers that circulate in maternal blood. *J Clin Invest*. 2007;117(10):3007-19.
23. Tsui NB, Chim SS, Chiu RW, Lau TK, Ng EK, Leung TN, et al. Systematic micro-array based identification of placental mRNA in maternal plasma: towards non-invasive prenatal gene expression profiling. *J Med Genet*. 2004;41(6):461-7.
24. Zhong XY, Gebhardt S, Hillermann R, Tofa KC, Holzgreve W, Hahn S. Parallel assessment of circulatory fetal DNA and corticotropin-releasing hormone mRNA in early- and late-onset preeclampsia. *Clin Chem*. 2005:1730-3.

Identificación de RNA mensajero de origen fetoplacentario en plasma materno de mujeres con embarazos normales y gestantes con restricción de crecimiento intrauterino

RESUMEN

Objetivo: cuantificar RNA específico de placenta en el plasma de mujeres con embarazos con fetos con Restricción de Crecimiento Intrauterino y gestantes con embarazos normales.

Materiales y métodos: se estudiaron 8 mujeres con embarazos con fetos con Restricción de Crecimiento Intrauterino y 18 mujeres con embarazos sin complicaciones, en el tercer trimestre de embarazo. Se cuantificó el RNA total libre en plasma materno por espectrofotometría y la expresión del gen hPL (Lactógeno Placentario Humano) a nivel de RNA mensajero por medio de la técnica Reacción en Cadena de la Polimerasa en Tiempo Real.

Resultados: se logró detectar RNA en plasma de origen fetoplacentario en el 100% de las gestantes. No se encontraron diferencias estadísticamente significativas entre los valores de RNA total extraído de plasma ($p=0,5975$) ni en la expresión del RNA mensajero del gen hPL ($p=0,5785$) entre casos y controles.

Conclusión: es posible detectar RNA mensajero de origen fetoplacentario en plasma materno durante el embarazo.

Palabras claves: RCIU, RNA mensajero, plasma, complicaciones del embarazo, lactógeno placentario humano, Reacción en Cadena de la Polimerasa de Transcriptasa Inversa.

Ayala RP, García RR, Rojas JD, Bermúdez M, Bernal J. Identification of messenger RNA of fetoplacental source in maternal plasma of women with normal pregnancies and pregnancies with intrauterine growth restriction. Colomb. Med. 2012; 43(3) 184-88