



Original Article

Frequency and variability of dental morphology in deciduous and permanent dentition of a Nasa indigenous group in the municipality of Morales, Cauca, Colombia

Frecuencia y variabilidad de la morfología dental en dentición temporal y permanente de un grupo de indígenas Nasa del municipio de Morales (Cauca, Colombia)

Eider Díaz¹, Lorena García¹, Michelle Hernández¹, Lesly Palacio¹, Diana Ruiz¹, Nataly Velandia¹, Judy Villavicencio², Freddy Moreno^{*3}

¹Dental Scientist at the Universidad del Valle (Cali, Colombia).

²School of Dentistry at the Universidad del Valle (Cali, Colombia).

³School of Basic Sciences at the Universidad del Valle (Cali, Colombia), Professor, School of Health Sciences at the Pontificia Universidad Javeriana (Cali, Colombia).

Díaz E, García L, Fernández M, Palacio L, Ruiz D, Velandia N, Villavicencio J, Moreno F. Frequency and variability of dental morphology in deciduous and permanent dentition of a Nasa indigenous group in the municipality of Morales, Cauca, Colombia. *Colomb Med.* 45(1): 15-24.

© 2014 Universidad del Valle. This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Article history:

Received: 18 January 2013
Revised: 9 January 2014
Accepted: 26 February 2014

Keywords:

Dental anthropology; dental morphology; non-metric dental traits; indigenous Nasa.

Palabras clave:

Antropología dental; morfología dental; rasgos morfológicos dentales coronales; indígenas nasa.

Abstract

Objectives: To determine the frequency, variability, sexual dimorphism and bilateral symmetry of fourteen dental crown traits in the deciduous and permanent dentition of 60 dental models (35 women and 25 men) obtained from a native, indigenous group of Nasa school children of the Musse Ukue group in the municipality of Morales, Department of Cauca, Colombia.

Methods: This is a quantitative, descriptive, cross-sectional study that characterizes dental morphology by means of the systems for temporary dentition from Dahlberg (winging), and ASUDAS (crowding, reduction of hypocone, metaconule and cusp 6), Hanihara (central and lateral incisors in shovel-shape and cusp 7), Sciulli (double bit, layered fold protostylid, cusp pattern and cusp number) and Grine (Carabelli trait); and in permanent dentition from ASUDAS (Winging, crowding, central and lateral incisors in shovel-shape and double shovel-shape, Carabelli trait, hypocone reduction, metaconule, cusp pattern, cusp number, layered fold protostylid, cusp 6 and cusp 7).

Results: The most frequent dental crown features were the shovel-shaped form, grooved and fossa forms of the Carabelli trait, metaconule, cusp pattern Y6, layered fold, protostylid (point P) and cusp 6. Sexual dimorphism was not observed and there was bilateral symmetry in the expression of these features.

Conclusions: The sample studied presented a great affinity with ethnic groups belonging to the Mongoloid Dental Complex due to the frequency (expression) and variability (gradation) of the tooth crown traits, upper incisors, the Carabelli trait, the protostylid, cusp 6 and cusp 7. The influence of the Caucasoide Dental Complex associated with ethno-historical processes cannot be ruled out.

Resumen

Objetivos: Determinar la frecuencia, la variabilidad, el dimorfismo sexual y la simetría bilateral de 14 Rasgos Morfológicos Dentales Coronales (RMDC) en dentición temporal y permanente de 60 modelos (35 mujeres y 25 hombres) obtenidos de un grupo de indígenas escolares Nasa del cabildo Musse Ukue del municipio de Morales, departamento del Cauca.

Métodos: Estudio descriptivo transversal cuantitativo que caracteriza la morfología dental mediante los sistemas en dentición temporal Dahlberg (winging), Asudas (crowding, reducción de hipocono, metaconulo y cúspide 6), Hanihara (incisivos centrales y laterales en pala y cúspide 7), Sciulli (doble pala, protostilido, pliegue acodado, patrón cuspeo y número de cúspides) y Grine (cúspide de Carabelli); y en dentición permanente Asudas (Winging, crowding, incisivos centrales y laterales en pala, doble pala, cúspide de Carabelli, reducción del hipocono, metaconulo, patrón cuspeo, número de cúspides, pliegue acodado, protostilido, cúspide 6 y cúspide 7).

Resultados: Los RMDC más frecuentes fueron la forma de pala, formas surco y fosa de la cúspide de Carabelli, metaconulo, patrón cuspeo Y6, pliegue acodado, protostilido (punto P) y cúspide 6. No se observó dimorfismo sexual y hubo simetría bilateral en la expresión de los rasgos.

Conclusiones: La muestra estudiada presenta una alta afinidad con grupos étnicos pertenecientes al Complejo Dental Mongoloide debido a la frecuencia (expresión) y variabilidad (gradación) de los RMDC, incisivos en pala, la cúspide Carabelli, el Protostilido, la cúspide 6 y la cúspide 7; sin embargo, no se puede descartar influencia del Complejo Dental Caucasoide asociado con procesos etno-históricos.

*Corresponding author:

Moreno F. School of Dentistry at the Universidad del Valle (Cali, Colombia)
Calle 3 A # 36B-00 Edificio 132 PBX: 554 2469 - 5580759
E-mail: freddy.a.moreno@correounivalle.edu.co

Introduction

Analysis of dental morphology in the context of dental anthropology seeks to understand the manner in which the frequency, sexual dimorphism and bilateral symmetry of Tooth Crown Morphological Traits (TCMT) present in deciduous and permanent teeth¹. The TCMT consist of phenotypic enamel forms expressed and regulated by the genome of an individual and population during odontogenesis. Structures can be positive (tubercular) or negative (inter-tuberculars and fosomorfos) that have the potential to be present, or not, at a specific site (frequency) and in a different form (variability) in one or more members of a population group²⁻⁶.

Based on the TCMT, dental morphology is studied from an interdisciplinary viewpoint (biology, anthropology, dentistry, paleopathology, archeology, forensic science) because teeth can be used in the estimation of biological relationships between populations. This is accomplished by comparatively analyzing past and present human groups in an attempt to clarify the historical, cultural and biological macro and micro-evolutionary processes that lead to an understanding of the origin, formation, contacts, displacements, migrations pathways and isolates that have led to the populating of the planet and ethnic variation of humanity^{1,3,7,8}. It similarly constitutes an accurate means of recognizing individuals whose death makes it difficult to distinguish them by other processes which are part of the individual reconstruction of osteobiography (odontography) or for that of the general population^{1,8,9}.

This research seeks to observe the expression of 14 TCMT's from a group of Nasa indigenous people from the Musse Ukue council of in the municipality of Morales, province of Cauca, for the purpose of further processing of dental morphological markers that contribute to the ethnographic description of Southwestern Colombia. This is an essential process in forensic dental identification to document the elements of the basic identification quatrains (sex, age, ethnicity and height)⁹.

Materials and Methods

Population and sampling

This is a quantitative, cross-sectional, descriptive study of the frequency, variability, sexual dimorphism and bilateral symmetry of 14 TCMT's in deciduous and permanent dentition from study models obtained from a group of school children of the indigenous Nasa tribe of the Musse Ukue council of the municipality of Morales, Cauca, Colombia. The school children were from Colombian Nasa parents and grandparents and were between the ages of 5 to 18 years. Students were selected from a survey and interview examination who met the inclusion criteria (with clinically healthy teeth where the TCMT were taken into account) and who were asked to sign an informed consent and had parents that authorized their participation via an informed consent agreement. The sample number was selected by convenience based on the central limit theorem which considers a sample of 30 to be significant because "regardless of the functional form of the population from which the sample is drawn, the distribution of sample means calculated with samples of size n drawn from a population with a mean m and finite variance s^2 , approaches a normal distribution with mean m and standard error s^2/n , when n

increases. If the n is large, $n \geq 30$, the distribution of sample means can approximate a normal distribution"¹⁰.

Standardization of morphological analysis

The six observers learned to manage the odontoscopic systems (Asudas-Arizona State University Dental Anthropology System-, Dahlberg, Hanihara, Sciulli and Grine) with a standardization protocol and double blind in nature to control bias and achieve a unified observational criteria, according to that described by Nichol and Turner II¹¹.

Observation

Once the research gained the support of the Research Ethics Committee for Human Subjects of the School of Health, Universidad del Valle and in accordance with Article 11 of Resolution 008430 of the Ministry of Social Protection¹² and with the Helsinki Declaration¹³, who classified this study as being without risk, we then proceeded to the observation of 14 TCMT in plaster models using 10-power magnification from a Hu-Friedy® fine point explorer and from gradations of Dahlberg (winging) deciduous dentition system, Asudas (crowding, reduced hypocone, metaconules and cuspids 6) , Hanihara (central and lateral incisors in shovel-shape and cusp 7), Sciulli (double shovel-shaped, protostylid, layered fold, groove pattern and number of cusps) and Grine (Carabelli cusp); and in permanent dentition through Asudas (Winging, crowding, central and lateral incisors in shovel-shaped, double shovel-shaped, Carabelli cusp, hypocone reduction, metaconules, groove pattern, number of cusps, layered fold, protostylid , cusp 6 and cusp 7) (Table 1).

Statistical analysis

The data obtained from TCMT observation were entered into a template on Excel® and processed with the SPSS®, 17.0 Software through descriptive tests and by univariate and bivariate analysis tests using nonparametric tests (Wilcoxon and Mann-Whitney U) for each of the morphological features. A probability level of <0.05 was considered statistically significant. To determine biological distances the distance matrix was used starting with the hierarchical cluster classification by means of the Euclidean distance squared, while the dendrogram was obtained by using the Ward method.

Results

Estimating the degree of concordance for standardization of observers was done by means of the Kappa test with Stata® 6.0 Software. Criteria for acceptable inter-observer reliability (observer vs. rater) were 93%, 80%, 85%, 92%, 92% and 87% for the five observers; and intra-observer (observer vs. observer) were 90%, 81%, 81%, 81%, 87% and 85%, respectively.

Frequencies were obtained for each of the TCMTs for the 60 models studied (35 females and 25 males). The most common features were the shovel-shaped central and lateral incisors, high frequencies of the groove and fossa forms of Carabelli's cusp, the metaconules, the Y6 cusp pattern, layered fold, protostylid and cusp 6. None of the features presented sexual dimorphism ($p < 0.05$) except for the upper incisors in the deciduous teeth, and all TCMTs are expressed bilaterally ($p < 0.05$) (Table 2). With respect to biological distance in the distance matrix (Table 3) as well as in

Table 1. Methods of observation of MCDT.

Feature	Tooth	Gradation	Degree of expression		Method of reference
			Rank	Presence	
Temporary Dentition					
Winging	ui1	0. Absent 1. Unilateral 2. Bilateral	0-2	2	Dahlberg (7)
Crowding	Ui2	0. Absent 1. Vestibular 2. Palatine	0-3	1-2	ASUDAS (7)
Upper central incisors	ui1	0. Absent 1. Semi-shovel 2. Shovel shaped 3. Strong shovel	0-3	2-3	Hanihara (16)
Upper lateral incisors	ui2	0. Absent 1. Semi-shovel 2. Shovel shaped 3. Strong shovel	0-3	2-3	Hanihara (16)
Double shovel	ui1	0. Absent 1. Mesal 2. Distal 3. Both	0-3	1-3	Sciulli (25)
Carabelli cusp	um2	0. Absent 1. Depression in form of U or Y 2. Two parallel grooves 3. Small cusp 4. Free vertex cusp	0-4	3-4	Grine (26)
Reduction of the hypocone	um2	0. Absent 1. Free vertex cusp 2. Presents a smooth cuspule 3. Presents a small cuspule 4. Presents a medium-sized cusp 5. Presents a normal cusp 6. Presents a larger cusp of grade 5	0-1	1	ASUDAS (7)
Metaconules	um2	0. Absent 1. Presents a smooth cuspule 2. Presents a well-marked cuspule 3. Presents a small well-defined cuspule 4. Small cuspid 5. Intermediate sized cuspid	0-5	1-5	ASUDAS (7)
Protostylid	um2	0. Absent. 1. Fossa or groove 2. Cuspid	0-2	2	Sciulli (25)
Layered fold	Im2	0. Absent. 1. Present.	0-1	1	Sciulli (25)
Cusp pattern	Im2	+. Cusps 1, 2, 3&y 4 in contact X. Cusps 1 & 4 in contact Y. cusps 2 & 3 in contact	Y, +, X	Y	Sciulli (25)
Number of cusps	Im2	2. MI & Mv. 3. MI, Mv & Dv. 4. MI, Mv, Dv & DI. 5. MI, Mv, Dv & DI & D. 6. MI, Mv, Dv & DI, D & cusp 6. 7. MI, Mv, Dv & DI, D, cusp 6 & cúsp7. 8. More than 7 cusps	2-Aug	5	Sciulli (25)
Cusp 6	Im2	0. Absent 1. Cusp 6 << Cusp 5 2. Cusp 6 < Cusp 5 3. Cusp 6 = Cusp 5 4. Cusp 6 > Cusp 5 5. Cusp 6 >> Cusp 5	0-5	1-5	ASUDAS (7)
Cusp 7	im2	0. Absent 1. Trace 2. Small cusp 3. Well developed	0-3	1-3	Hanihara (16)
Permanent Dentition					
Winging	UI1	0. Absent 1. Bilateral 2. Unilateral 3. Strong unilateral 4. Contralateral	0-4	1-3	ASUDAS (7)

Continued Table 1. Methods of observation of MCDT.

Feature	Tooth	Gradation	Degree of expression		Method of reference
			Ranko	Presence	
Crowding	ui2	0. Absent 1. Vestibular 2. Palatine	0-3	1-2	ASUDAS (7)
Upper incisors	UI1	0. Absent 1. Smooth 2. Well-marked 3. Semi-shovel shape 4. Marked semi-shovel shaped 5. Shovel-shaped 6. Marked shovel-shape 7. Barrel form	0-7	2-7	ASUDAS (7)
Shovel-shaped Incisors	UI2	0. Absent 1. Smooth 2. Well-marked 3. Semi-shovel shape 4. Marked semi-shovel shaped 5. Shovel-shaped 6. Marked shovel-shape 7. Barrel form	0-7	2-7	ASUDAS (7)
Double shovel	UI1	0. Absent 1. Smooth 2. Well-marked 3. Semi-double shovel 4. Double shovel 5. Marked double shovel 6. External doublé shovel	0-6	2-6	ASUDAS (7)
Carabelli cusp	UM1/UM2	0. Absent 1. Groove present 2. Fossa present 3. Small depression in form of Y 4. Large depression in form of Y 5. Small cusp 6. Medium cusp 7. Free vertex cuspid	0-7	2-7	ASUDAS (7)
Reduction of the hypocone	UM1/UM2	0. Absent 1. Free vertex cuspid 2. A smooth cuspule is presented 3. A small cuspid is presented 4. A moderate-sized cuspid is presented 5. A normal cusp is presented 6. A cuspid larger than grade 5 is presented	0-6	1	ASUDAS (7)
Metaconules	UM1/UM2	0. Absent 1. Presents a smooth cuspule 2. Presents a well-marked cúspule 3. Presents a small well-defined cuspule 4. Small cuspid 5. Intermediate sized cuspid	0-5	1-5	ASUDAS (7)
Protostylid	LM1/LM2	0. Absent. 1. A small hole is presented 2. Curved distal buccal groove is presented 3. A second groove is presented from the buccal groove that extends mesially 4. The second groove is slightly more pronounced than the first 5. The second groove is very deep 6. A small, blunt cusp 7. A mid-sized cusp presents with free point	0-7	2-7	ASUDAS (7)
Layered fold	LM1/LM2	0. Absent 1. Little pronounced constriction 2. Marked constriction 3. Form of L	0-3	2-3	ASUDAS (7)
Cusp pattern	LM1/LM2	+. Cusps 1, 2, 3 & 4 in contact X. Cusps 1 & 4 in contact Y. Cusps 2 & 3 in contact	Y, +, X	Y	ASUDAS (7)
Number of cusps	LM1/LM2	4 Cusps 1, 2, 3 & 4 5 Cusps 1, 2, 3, 4 & 5 6 Cusps 1, 2, 3, 4, 5 & 6 7 Cusps 1, 2, 3, 4, 5, 6 & 7	4-Jul	5	ASUDAS (7)
Cusp 6	LM1/LM2	0. Absent 1. Cusp 6 << Cusp 5 2. Cusp6 < Cusp 5 3. Cusp 6 = Cusp 5 4. Cusp 6 > Cusp 5 5. Cusp 6 >> Cusp 5	0-5	1-5	ASUDAS (7)
Cusp 7	LM1/LM2	0. Absent 1. Lines 1.5. Marked lines 2. Small cuspid 3. Medium-sized cuspid 4. Large cuspid	0-4	1-4	ASUDAS (7)

their respective dendrograms (Fig.1) it can be seen that the sample has a high affinity for the Mongoloid Dental Complex due to the frequency of the TCMT incisors with shovel shape, Carabelli's cusp, the protostylid, the cusp 6 and cusp 7. The distance matrix has a sensitivity (Bootstrapping – 1000 resampling method) of $p=0.49$.

Discussion

Dental crown morphological traits Winging, Crowding, central and lateral incisors in shovel and double shovel shape

Winging involves rotating one or both upper central incisors relative to the midline. Scott and Turner⁷ relate this feature with the absence of space in the dental arch that prevents the proper alignment of the incisors. According to Rodríguez⁸, the meso-lingual rotation of both incisors is considered the product of genetic factors characteristic of native American populations while the rotation of a single tooth or both in a disto-lingual direction is caused by crowding. In a population of Amazonian Indians this feature was observed with high frequency¹⁴. In this study for the deciduous dentition winging was absent, while in permanent dentition a relative frequency of the trait was observed, represented in grade 1 expression (disto-buccal rotation of both incisors). There was no sexual dimorphism.

Regarding crowding or overcrowding of the upper lateral incisors, it is more common in Mongoloid populations and less frequent in Caucasian populations⁹, but it was absent in temporary dentition and was low in frequency with permanent dentition. Sexual dimorphism was associated with an increased number of females.

The shovel shape showed a high frequency, mainly in the middle grades (semi-shovel and marked semi-shovel). Also, bilateral symmetry was evident along with the absence of sexual dimorphism. Turner¹⁴ observed this trait in 98.8% of East Asians and 99.8% of South American Indians. Among indigenous people from Colombia, it varies between 50% and 100%⁹. Aragón *et al.*¹⁴ observed high frequencies among indigenous tribes in the Amazon basin.

In the case of the double-shovel form, its low frequency is highlighted (smooth grades and lines), bilateral symmetry and sexual dimorphism associated with a greater number of females. Aragón *et al.*¹⁴ also observed high frequencies in Amazonian indigenous groups.

Carabelli's cusp, reduction of the hypocone and metaconules

The cusp or tubercle of Carabelli was considered virtually absent from the sample. However, it is important to note that in both dentitions the groove and fossa shapes predominated over the cusp forms with dichotomous expression, which is characteristic of American indigenous peoples⁹. Hanihara¹⁶ observed low frequencies of this trait among Japanese and high averages in Blacks and White Americans and found that it distinguishes Caucasians from Asian populations. In the latter case it is predominately in the groove and fossa shapes. Turner¹⁵ found significant expressions among sinodonts, South American Indians, and Northeast Europeans. Among indigenous Colombians, the frequency varies between 20% and 90%⁹ and the groove and fossa shapes predominate over the cusp form⁴.

Severe reduction and the absence of the disto-lingual cusp (hypocone) is a valued trend from the first upper molar to the second upper molar⁹ and is associated with a simplification of the dental morphology and reduced size. In this study, the reduction of the hypocone resulted in the absence of the bilateral form in both sexes. Hypocone is most prevalent in grades 0 and 1. In indigenous Colombians, it is between 80% and 100%. The global variation is between 13% and 95%, with a minimum found among Australian aborigines and a maximum among Mongoloids, which was true in regard to this study⁹.

Regarding metaconules, or small cuspid with a free tip located at the distal edge of the upper molars between the disto-buccal and disto-palatal cuspid, they were absent from the sample. This is consistent with that reported in the literature since in American populations, whether prehispanic indigenous or contemporary, the trend is toward decreasing expression⁷.

Protostylid and layered fold

The frequency of this feature in deciduous dentition was significant from the expression of thin cusps with blunted tips on the labial surface of the meso-buccal cusp of the lower second molars. In permanent teeth this trait was practically absent, but significant expression was found in grade 1 (point P or foramen caecum) in permanent teeth, which has been defined as an americanoide trait. There was no sexual dimorphism; however there was bilateral symmetry^{7,9,17}.

Concerning the layered fold, which describes the way in which the meso-lingual cusp is directed toward the central fossa of the temporary and permanent lower molars, a high frequency in both dentitions is reported with bilateral symmetry and no sexual dimorphism. The TCMT is an important marker in sinodont populations, and with a high frequency in indigenous Colombians between 60% and 100%⁹.

Groove pattern and number of cusps

The groove pattern of the second lower deciduous molars and first lower permanent molars describes the configuration of contact of the cusps and their number. The classic pattern is "Y", or "Driopitecino" originating from past Asian populations, along with "X" configurations and the "+" or "cruciform" which are all considered as reductions that are frequently observed in Caucasoid groups⁹. Among Colombian indigenous populations a predominance of Y⁶ and Y⁷ configuration is reported (temporary dentition) wherein the Y cusp pattern has a strong genetic control that allows it to be retained from the last Asian populations that inhabited the Americas by way of the Bering land bridge from which prehispanic and current Amerindian populations are derived¹⁴. In this study, in temporary dentition the most frequent cusp pattern is Y⁵, while in permanent dentition, the first molar is +5 and the second molars are +4 which may be associated with miscegenation. This trait does not present with sexual dimorphism and had bilateral expression.

Cusp 6 and 7

The frequency and variability of cusp 6 (located between the disto-buccal and disto-lingual cusps of the lower molars) and cusp 7 (located on the marginal edge between the meso-lingual and disto-lingual cusps of the lower molars) are present in both

Table 2. Relative frequencies of dental morphological features

Features	Frequency		Sexual dimorphism		Bilateral symmetry		
	n	Tooth	%	Tooth	Mann-Whitney p<0.05	Teeth	Wilcoxon p<0.05
Winging	8	51-61	0	51-61	0.2	-	-
	46	11-21	45.6	11-21	0.4	-	-
Crowding	15	52-62	0	52-62	1.0	-	-
	41	12-22	29.3	12-22	0.0	-	-
Shovel-shaped Incisors	43	11	90.7	11	0.8	11-21	1.0
	43	21	90.7	21	0.8	-	-
	38	12	81.5	12	0.2	12-22	0.0
	37	22	86.4	22	0.1	-	-
Double shovel-shaped incisors	43	11	20.9	11	0.0	11-21	0.0
	38	12	5.3	12	0.1	-	-
	42	21	19	21	0.0	12-22	0.0
	36	22	8.4	22	0.1	-	-
Carabelli cusp	49	16	49	16	0.5	16-26	0.0
	50	26	28	26	0.0	-	-
	32	55	6.3	55	0.6	55-65	0.8
	29	65	6.9	65	0.3	-	-
Reduction of the hypocone	17	17	29.4	17	0.4	17-27	0.5
	18	27	27.8	27	0.3	-	-
Metaconules	47	16	31.9	16	0.9	16-26	0.2
	47	26	38.3	26	0.4	-	-
	12	17	0	17	1.0	17-27	0.2
	17	27	11.8	27	0.3	-	-
	26	55	7.7	55	1.0	55-65	0.6
	29	65	6.9	65	0.1	-	-
Protostylid	39	36	2.6	36	0.0	36-46	0.5
	42	46	9.5	46	0.2	-	-
	24	37	4.2	37	0.6	37-47	0.1
	15	47	0	47	0.3	-	-
	26	76	42.3	76	0.4	76-86	0.3
	26	86	53.8	86	0.7	-	-
Layered fold	51	36	41.1	36	0.5	36-46	0.6
	54	46	42.6	46	0.6	-	-
	26	37	0	37	0.7	37-47	0.6
	26	47	3.8	47	0.8	-	-
	23	76	78.3	76	0.8	76-86	1.0
	19	86	73.7	86	0.1	-	-
Cusp pattern	39	36	17.9	36	0.1	37-47	0.1
	39	37	12	37	0.4	-	-
	25	46	12.8	46	0.0	36-46	0.6
	23	47	0	47	0.3	-	-
	14	75	78.6	75	0.0	75-85	0.6
	12	85	33.3	85	0.1	-	-
Number of cusps	47	37	34.8	37	0.7	37-47	0.2
	46	36	100	36	0.7	-	-
	23	47	36.3	47	0.1	36-46	0.6
	22	46	100	46	0.2	-	-
	19	75	100	75	0.2	75-85	0.5
	21	85	100	85	0.1	-	-
Cusp 6	47	36	38.3	36	0.2	36-46	0.9
	46	46	34.8	46	0.4	-	-
	24	37	12.5	37	0.3	37-47	0.1
	23	47	21.7	47	0.8	-	-
	23	75	17.4	75	0.1	75-85	0.7
	23	85	12.9	85	0.3	-	-
Cusp7	53	36	2.8	36	0.2	36-46	0.7
	53	46	5.7	46	0.4	-	-
	25	37	0	37	1.0	37-47	1.0
	24	47	0	47	1.0	-	-
	20	75	15	75	0.2	75-85	0.1
	22	85	0	85	1.0	-	-

dentitions. They are, not considered to be at significant levels if you consider that cusp 6 is characteristic of Asian populations and cusp 7 of Negro populations¹⁴.

Biological distance and dental complexes

Because of the frequency and variability of TCMTs, human populations can be associated with geographical distributions, and different researchers have ethnographically classified human beings in complex populations from dental morphology. The first of these complexes was defined by Hanihara in 1966¹⁶ as the Mongoloid Dental Complex, which brings together different populations from East Asia that are characterized by having a complex dental morphology represented in a high frequency of shovel-shaped, layered fold and cusp pattern 6 incisors. Later, Turner in 1984¹⁵ divided the Mongoloid Dental Complex into two groups. The first subdivision, or Sinodont, composed of Northeast Asian populations, is characterized by the addition and enhancement of some TCMTs, such as the shovel-shaped and layered fold incisors, along with the Y6 cusp pattern, the protostylid (cusp forms) and winging. The second subdivision or sundadont covers Southeast Asian populations that have retained an ancestral condition and have simplified the expression of some morphological traits.


On the other hand, Zoubov¹⁶ proposed a dental delimitation of global populations into two complexes: the Eastern Dental Complex, the equivalent of the Mongoloid Dental Complex proposed by Hanihara; and the Western Dental Complex,

consisting of Northern Caucasoid and Negroid populations (Southern Caucasoid populations) characterized by the high frequency of the Carabelli cusp (cusp forms), of the cusp pattern X, the groove pattern + and the cusp 7 more prevalent in Negroid populations. Irish¹⁸ subdivided the Negroid populations of Southern Africa (Western Dental Complex) into the Sub-Saharan Dental Complex and North African Dental Complex.

Edgar (2007)¹⁹ grouped humans into five clusters: the Mongoloid Dental Complex formed by sinodont and sundadont groups; the Caucasoid Dental Complex formed by Western Eurasian groups (Europe, North Africa, Middle East and India); the Saharan African Dental Complex (composed of West African and South African subgroups closer to sundadont populations of the South Pacific), several Pacific groups of Sahul or Oceania and American Paleo-Indians who present frequencies and morphological variations that take them out of the complexes described.

In the case of the American populations, currently the model proposed by Turner in 1984¹⁵ is accepted. It suggests that the populating of the American continent initially occurred by human sinodonts who migrated from Northern China and crossed Bering land bridge. This is affirmed by past and present American indigenous groups having a sinodont dental morphology, and therefore, should be included in the Mongoloid Dental Complex, which is in agreement with the miscegenation with other ethnic groups. The high frequency of shovel-shaped, layered fold incisors, protostylid and cusp pattern Y⁶ support the thesis of the Northeast

Figure 1. Dendrogram obtained from the distance matrix among world populations based on squared Euclidean distance



Asian origin for the first settlers of the American continent. For Zoubov¹⁷, the high frequency of the protostylid in grade 1 (point P) as a unique feature of American populations that allows him to propose the existence of a Amerindoid Dental Complex, consisting of all the American Paleo-Indian and the contemporary populations that are derived from them.

Regarding the Colombian population, the study of dental morphology and its association with dental complexes reviewed through ethno-historical processes that have occurred in the country is quite complicated. Rodríguez⁹ states that past indigenous peoples are characterized by presenting high frequencies of winging, crowding, reduced hypocone, layered fold and P point of the protostylid, which places them closer to the Paleo-Indian sinodontes of the Mongoloid Dental Complex. However, in the case of the contemporary indigenous people, the situation varies mainly due to the miscegenation that occurred with the arrival of Northern Caucasoid human groups from Western Europe (Western Dental Complex). They populated American territory in three historical periods recognized as the discovery, conquest and colonization. These groups were characterized by a high frequency of cusp expression of the cusp of Carabelli, the X cusp pattern and the + cusp pattern. However, the dental morphology is simplified given the low frequency of shovel-shaped incisors and double-shovel pattern, the layered fold, and cusps 6 and 7^{17,18}.

Leon and Riaño²⁰ analyzed a series of plaster casts of several current Colombian indigenous populations obtained by the Human Expedition conducted by the Pontificia Universidad Javeriana. The results indicate a high frequency of shovel-shaped incisors, a low frequency of the Carabelli cusp, a high proportion of hypocone reduction, variable frequency of sixth cusp, a high frequency percentage of the seventh cusp, high percentages of the upper layered fold and high frequencies of the protostylid. Cerón²¹ compared some discrete dental and cranial traits among a group of Andean mestizos and a native Embera population, noting that the latter group had high frequencies of shovel-shaped incisors and cusp 6. Similarly, Negroid human groups (Southern Caucasoids from the Western Dental Complex) were brought as slaves to the South American continent and were distributed throughout different regions of Colombia. For this reason, in virtue of macro-evolutionary process represented by numerous migrations, contacts and isolations, the multiethnic, multicultural and polygenic nature of the Colombian population arose as described by Yunis *et al.*, in 1992 and Ramos *et al.*, in 1993 - cited by J.V. Rodríguez⁹ - so that the tri-ethnic genetic composition average in Colombian remains constituted by Caucasoid genes (62%), Mongoloid (26%) and Negroid (12%), which were distributed differentially throughout differing regions of the country.

Perhaps it is in the region of Southwestern Colombia where these processes are accentuated and in which the greatest number of studies have been carried out in populations described as Caucasoid Mestizos, Indigenous and Afro-Colombian. The Research Group of Oral and Maxillofacial Surgery at the University del Valle has studied several populations from the southern part of the department (i.e. province) of Valle del Cauca and the northern part of the department of Cauca, and inferred that the frequency of TCMTs reflect historical miscegenation and dominance of these phenotypic expressions. This occurred in such a way that a group

of Mestizos in the city of Cali was characterized in a simplification of the dental morphology as having a low frequency of the cusp of Carabelli (this trait was expressed ambiguously in their fossa forms - characteristic of sinodonts - and cuspids of medium size - a characteristic of Caucasoids), a reduced hypocone (a property of Western Caucasoid groups) and the high frequency of the point P of the protostylid (an exclusive feature of Amerindian populations)⁴.

Similarly, by studying two groups of Afro-Colombians from the Department of Cauca, one from the city of Puerto Tejada²² and the other from the city Villarica²², high frequencies of Carabelli's cusp were observed of a medium size, groove pattern +, "X" cusp pattern, and a high frequency of cusp 7, suggesting notable influence by the Western Dental Complex. Concerning the Afro-Colombian populations, Delgado²⁴ indicates that the populations were derived from Africans who came to the American continent as slaves from West Africa, Central West (Sub-Saharan Africa), South-East and North, all of which are ranked in the Western Dental Complex (Southern Negroid).

The term "dental complex" refers to the characterization of large population groups, according to a specific combination of TCMTs, and since modern human groups have the same number of these features in both dentitions, the only detectable difference is in the frequencies of these traits. That is why it is necessary to cover a wider range of population groups with significant samples.

According to the results obtained from this study, it can be said that the indigenous group of Nasa who formed the sample for this study have high frequencies of TCMT characteristic of the Mongoloid Dental Complex. Specifically with the sinodonts, and similarly with other Colombian and American indigenous groups, the findings are consistent with those reported by Turner¹⁵, Hanihara¹⁶, Zoubov¹⁷ and Rodríguez⁹, which also coincide with the theory of the Mongoloid origin of the indigenous tribes of South America. However, the significant frequencies of some features suggest the influence of contemporary Caucasoid human groups (Fig. 1).

Conclusions

The most frequent TCMT in both dentitions were shovel-shaped forms, the groove shapes and fossa forms, Carabelli cusp, the metaconules, the Y⁶ cusp pattern, the layered fold, protostylid (point P) and the cusp 6. The TCMT observed did not display sexual dimorphism and had bilateral symmetry in their expression. The Nasa indigenous group who constituted the sample had a high affinity with ethnic groups belonging to the Mongoloid Dental Complex. This can be corroborated with the distance matrix and dendrogram obtained from the frequencies of the TCMT shovel-shaped incisors, Carabelli cusp, the protostylid, the cusp 6 and cusp 7. However, one cannot rule out the influence of Caucasoid Dental Complex associated with ethno-historical processes that occurred in Southwestern Colombia.

This research found new elements of invaluable ethnographic value from the analysis of dental morphology that eventually will allow us to understand the human diversity of this region of Southwestern Colombia, the establishment of regional linkages associated with macro-evolutionary events that occurred

Table 3. Distance matrix among world populations based on Euclidean squared distances

Population	Sun	Sin	NEA	NES	Esk	Ale	INA	ISA	NE	Ch	Jap	An	Eur	NA	SA	AS	IC	Wau	Nuk	Chim	MM	Pae	Guam	Core	Gua	W	Guan	Ama	Emb	AT	AP	AV	AG	AFT	Mear	MCal	Na	
Sun	0	0.8	0.5	1.1	1.4	2.7	2.2	1.6	4.3	1.5	2.9	8.5	6.0	6.4	18.6	7.1	2.3	6.2	9.6	17.8	7.8	2.0	4.8	2.4	10.7	16.6	3.9	17.0	2.7	14.3	14.0	5.7	8.0	7.7	8.6	7.2	1.6	
Sin		0	0.2	0.4	1.6	4.3	1.3	0.4	8.6	2.5	4.3	12.8	10.6	10.7	20.1	11.1	2.5	9.0	12.4	18.1	10.8	3.9	6.9	2.5	12.4	14.1	6.1	14.2	2.0	17.1	17.2	9.8	12.6	10.4	7.9	11.9	2.4	
NEA			0	0.5	1.1	3.1	1.5	0.8	7.3	1.7	3.1	11.0	9.5	9.7	20.6	9.9	2.4	7.1	12.7	17.7	9.0	3.5	5.3	2.1	11.6	15.6	5.6	15.5	2.2	17.0	16.7	8.4	11.4	9.6	9.2	10.3	1.9	
NES				0	1.0	4.3	0.5	1.2	9.2	2.3	3.3	12.9	11.9	12.5	23.2	13.2	3.9	10.4	14.6	23.2	12.9	4.1	7.3	3.3	14.7	16.9	8.5	17.9	3.7	21.6	21.8	11.4	14.7	11.8	10.8	13.1	3.0	
Esk					0	4.4	1.8	2.6	7.5	0.6	0.9	8.3	10.5	12.1	21.8	10.9	3.7	7.2	17.3	23.9	8.6	2.7	3.7	2.0	10.9	18.5	8.0	22.7	4.4	22.3	21.9	8.6	12.1	8.5	14.4	9.4	1.7	
Ale						0	4.4	6.8	5.8	4.4	4.3	11.3	8.4	8.0	31.1	10.1	7.0	6.4	13.8	19.3	9.7	7.9	5.8	6.9	17.9	27.9	7.5	20.1	8.5	20.1	18.3	9.0	12.1	14.8	17.0	10.5	5.9	
INA							0	2.8	11.0	3.9	4.3	14.1	14.4	14.7	25.7	15.1	4.9	13.5	15.8	24.2	15.4	6.0	8.1	4.1	16.8	17.6	10.5	17.1	6.5	24.7	24.8	14.4	17.7	13.8	12.7	15.6	5.5	
ISA								0	10.2	3.3	5.8	14.6	11.8	11.9	18.5	12.2	2.9	9.6	13.2	18.4	11.6	4.1	8.6	2.9	12.1	12.5	6.5	14.5	0.9	16.4	16.9	10.3	13.1	10.5	6.6	12.9	2.4	
NE									0	5.4	6.4	4.1	0.5	1.3	18.2	2.5	6.7	6.0	7.4	20.5	5.8	3.5	5.9	7.8	11.1	26.3	3.3	27.3	9.9	11.0	9.9	1.5	1.9	6.2	14.6	1.5	5.8	
Ch										0	0.6	6.9	7.8	9.6	21.4	9.0	4.3	4.2	16.7	24.0	6.3	2.1	3.3	2.8	10.2	21.2	6.8	25.9	3.9	20.1	19.3	5.6	9.0	7.5	15.0	6.7	0.6	
Jap											0	6.7	9.7	11.9	26.1	11.0	6.4	5.2	20.4	28.0	7.5	3.7	3.1	4.1	12.5	25.8	9.5	30.1	7.2	25.3	24.1	7.5	11.4	9.4	20.3	7.9	2.2	
AN												0	5.8	7.8	14.4	3.4	7.1	9.3	15.7	22.5	3.8	4.4	2.7	6.3	5.2	23.4	6.6	32.3	14.5	16.7	15.5	4.0	4.7	2.6	21.2	1.9	8.5	
Eur													0	0.5	16.7	2.2	8.0	7.1	6.0	19.7	6.8	4.7	8.5	9.9	11.7	26.6	3.0	27.7	10.6	8.2	7.3	1.3	0.1	6.7	13.4	1.7	7.3	
NA															0	16.7	2.0	7.7	8.4	3.8	15.5	7.6	6.2	9.6	10.5	12.5	25.0	2.2	22.8	10.7	5.8	4.9	2.4	1.7	8.0	11.3	3.2	8.9
SA																0	10.1	9.8	27.6	13.7	17.8	15.9	11.4	18.3	12.4	5.3	6.9	10.1	22.9	16.7	9.3	11.4	14.4	11.9	5.3	10	13.6	19.1
AS																	0	5.0	8.5	6.4	11.7	3.5	4.9	5.5	6.8	5.2	18.5	1.3	21.3	10.7	5.4	4.5	1.8	1.2	3.0	11.7	1.5	8.6
IC																		0	9.7	8.8	10.8	6.1	2.4	4.0	0.6	4.5	7.8	2.4	11.4	3.5	10.0	10.2	6.5	7.7	3.8	5.7	7.2	4.1
Wau																		0	20.3	21.3	3.4	7.6	5.3	8.6	12.9	31.9	7.1	32.1	7.1	17.8	15.4	3.8	7.3	11.2	21.2	6.0	3.5	
Nuk																			0	14.4	17.2	10.0	17.6	13.5	16.7	17.9	4.7	14.4	13.6	4.8	5.6	10.1	7.7	12.1	5.5	11.0	15.5	
Chim																			0	13.6	20.9	16.8	14.3	13.3	14.5	14.5	8.5	6.9	16.2	7.7	6.8	17.8	17.5	17.0	11.8	20.1	21.8	
MM																				0	6.2	2.2	5.5	4.2	22.0	4.2	26.5	9.0	12.3	10.3	2.7	4.6	4.4	18.2	3.1	6.0		
Pae																					0	4.4	2.4	6.1	14.8	3.8	23.0	4.5	13.0	13.3	3.5	4.8	2.8	9.7	3.7	2.1		
Guam																						0	2.5	5.3	19.9	5.7	23.9	8.7	18.0	16.3	5.3	8.0	4.8	18.4	4.9	4.8		
Core																							0	4.4	9.8	4.4	15.3	3.7	14.8	14.6	7.2	9.5	4.1	9.4	7.7	3.1		
Gua																								0	10.1	5.7	22.3	10.4	11.4	11.3	7.1	7.6	1.3	13.6	6.4	9.5		
W																								0	13.7	9.8	13.2	16.3	18.8	23.9	23.5	12.4	6.6	24.7	19.4			
Guam																								0	14.3	5.3	3.9	3.5	2.5	2.6	4.1	6.1	3.7	5.6				
Ama																								0	16.2	15.8	16.8	28.4	28.4	7.6	31.2	28.4						
Emb																								0	13.1	13.4	8.0	10.7	9.4	6.5	11.3	1.9						
AT																								0	0.3	8.6	6.2	10.3	6.6	10.5	16.7							
AP																								0	7.5	5.4	10.5	8.5	9.3	16.3								
AV																								0	0.6	4.1	14.2	0.6	4.8									
AG																								0	4.1	13.4	0.8	7.9										
AFT																								0	12.1	3.1	7.2											
MCnr																								0	16.9	12.2												
MCal																								0	6.7													
Na																								0	0													

* Sun=Sundadonts (15), Sin=Sinodonts (15), NEA=Northeast Asia (15), NES=Northeast Siberia (15), Ale=Aleutians (15), INA=North American Indians (15), ISA=South American Indians (15), NE=Northwestern Europe (15), Ch=Chinese (15), Jap=Japanese (15), AN=American Negrós (15), Eur=Europe (27), NA=North Africa (27), SA=South-Africa (27), ASS=Africa Sub-Sahara (27), IC=Indigenous Colombians (9), Wau=Waunana (20), MM=Murui-Muinane (20), Pae=Parézes (20), Gua=Guahibo (20), Core=Coreguaje (20), W=Wayúú (20), Guan=Guane (20), Ama=Amazonas (14), Emb=Emberá (21), AT=Afrodscendientes de Tumaco (24), AV=Afrodscendientes de Villavieja (24), AC=Afrodscendientes Guaji (24), A PT=Afrodscendientes de Puerto Tejada (22), MCar=Mestizos from Cartagena (23), MCal=Mestizos from Cali (4), Na=Nasa (this research).

since the settlement of America, and the concomitant mixing between populations with Mongoloid origin (American Indians), Caucasoid (Western Europeans), and Negroid (African descent).

Acknowledgements

The authors wish to thank the Nasa community council of Musse Ukue, the town of Morales, Cauca Department, for allowing us to obtain the sample needed for this study.

Conflict of interest:

None declared

References

1. Rodríguez CD. Antropología dental en Colombia. Comienzos, estado actual y perspectivas de investigación. *Antropo*. 2003; 4:17-27.
2. Rodríguez CD. La antropología dental y su importancia en el estudio de los grupos humanos. *Rev Fac Odont Univ Ant*. 2005; 16 (1 y 2):52-59.
3. Rodríguez CD. Antropología dental prehispánica: variación y distancias biológicas en la población enterrada en el cementerio prehispánico de Obando, Valle del Cauca, Colombia, entre los siglos VIII y XIII d.C. Syllaba Press. Miami; 2003.
4. Moreno F, Moreno SM, Díaz CA, Bustos EA, Rodríguez JV. Prevalencia y variabilidad de ocho rasgos morfológicos dentales en jóvenes de tres colegios de Cali, 2002. *Colomb Med*. 2004; 35 (Suppl 1):16-23.
5. Moreno SM, Moreno F. Eight Non-Metric dental traits in alive racially mixed population from Cali, Colombia. *Inter J Dental Anthropol*. 2005; 6:14-25.
6. Aguirre L, Castillo D, Solarte D, Moreno F. Frequency and Variability of five non-metric dental crown traits in the primary and permanent dentitions of a racially mixed population from Cali, Colombia. *Dental Anthropology*. 2006; 19(2):39-47.
7. Scott GC, Turner II CG. The anthropology of modern human teeth: dental morphology and its variation in recent human populations. London: Cambridge University Press; 1997.
8. Rodríguez JV. Introducción a la antropología dental. *Cuad Antropol*. 1989; 19: 1-41.
9. Ruiz A, Gómez C, Londoño D. Investigación Clínica: Epidemiología Clínica Aplicada. Pontificia Universidad Javeriana. Centro Editorial Javeriano: Bogotá; 2002.
10. Rodríguez JV. La antropología forense en la identificación humana. Universidad Nacional de Colombia. Bogotá; 2004.
11. Nichol CR, Turner II CG. Intra and inter-observer concordance in classifying dental morphology. *Am J Phys Anthropol*. 1986; 69: 299-315.
12. Ministerio de Salud. Resolución por la cual se establecen las normas científicas, técnicas y administrativas para la investigación en salud. Resolución 008430/1993 de 4 de octubre (acceded October 2006). Available from: <http://www.minproteccionsocial.gov.co/vbecontnt/library/documents/DocNewsNo267711.pdf>.

13. Asociación Médica Mundial. Principios éticos para las investigaciones médicas en seres humanos, Declaración de Helsinki. Finlandia, junio 1964 (accedido en enero de 2005). Disponible en URL: <http://www.wma.net/s/policy/b3.htm>.
14. Aragón N, Bastidas C, Bedón L, Duque P, Sánchez M, Rivera S, Triana F, Bedoya N, Moreno F. Rasgos Morfológicos Dentales Coronales en dentición temporal y permanente: Distancia biológica entre tres grupos indígenas del Amazonas Colombiano. *Revista Odontológica Mexicana*. 2008; 12(1): 13-28.
15. Turner II CG. Advances in the dental Search for native American origins. *Acta Anthropogen*. 1984; 8: 23-78.
16. Hanihara T. Dental and cranial affinities among populations of East Asia and the Pacific. *Am J Phys Anthropol*. 1992; 88: 163-82.
17. Zoubov AA. La antropología dental y la práctica forense. *Maguaré*. 1998; 13: 243-52.
18. Irish JD. Characteristic high- and low-frequency dental traits in Sub-Saharan African populations. *Am J Phys Anthropol*. 1997; 102(4): 455-67.
19. Edgar HJH. Microevolution of African American Dental Morphology. *Am J Phys Anthropol*. 2007; 132: 535-44.
20. León CF, Riaño C. Frecuencia de ocho rasgos morfológicos dentales en población indígena de Colombia, comparada con poblaciones indígenas americanas, europeas y asiáticas (tesis de postgrado). Santa Fe de Bogotá: Fundación Centro de Investigaciones y Estudios Odontológicos (CIEO), Universidad Militar Nueva Granada; 1997.
21. Estudio comparativo de las características cefalométricas, dentales y genéticas entre un grupo de indígenas Emberá del río Saija y un grupo de Santa Fe de Bogotá. VIII Congreso de Antropología en Colombia, Universidad Nacional de Colombia, Departamento de Antropología. Santa Fe de Bogotá 5, 6 y 7 de diciembre de 1997.
22. Rocha L, Rivas H, Moreno F. Frecuencia y variabilidad de la morfología dental en niños afro-colombianos de una institución educativa de Puerto Tejada, Cauca, Colombia. *Colomb Med*. 2007; 38: 210-21.
23. Marcovich I, Prado E, Díaz P, Ortiz Y. Análisis de la morfología dental en escolares afro-colombianos de Villarica, Cauca, Colombia. *Rev Fac Odontol Univ Antioq*. 2012; 24(1): 37-61.
24. Delgado-Burbano ME. Population affinities of African Colombians to Sub-Saharan Africans based on dental morphology. *HOMO*. 2007; 58: 329-56.
25. Sculli PW. Evolution of Dentition in Prehistoric Ohio Valley Native Americans: II. Morphology of the Deciduous Dentition. *Am J Phys Anthropol*. 1998; 106: 189-205.
26. Grine F. E. 1986. Anthropological Aspects of the Deciduous Teeth of African Blacks. En Singer L. y Lundy J. K. (Eds) *Variation, Culture, and Evolution in African Populations*. Johannesburg: Witwatersrand University Press.
27. Irish JD. Ancestral dental traits in recent Sub-Saharan Africans and the origins of modern humans. *J Hum Evol*. 1998; 34: 81-98.