

## ORIGINAL RESEARCH ARTICLE

# Correlates of Postpartum Sexual Activity and Contraceptive Use in Kano, Northern Nigeria

DOI: 10.29063/ajrh2018/v22i1.10

Zubairu Iliyasu<sup>1,5\*</sup>, Hadiza S. Galadanci<sup>2</sup>, Khadija M. Danlami<sup>1</sup>, Hamisu M. Salihu<sup>3</sup> and Muktar H. Aliyu<sup>4</sup>

Department of Community Medicine, Bayero University Kano, Nigeria<sup>1</sup>; Department of Obstetrics and Gynaecology, Bayero University Kano, Nigeria<sup>2</sup>; Department of Family and Community Medicine, Baylor College of Medicine, Houston, Texas, USA<sup>3</sup>; Departments of Health Policy and Medicine & Vanderbilt Institute for Global Health, Vanderbilt University, Nashville, Tennessee, USA<sup>4</sup>; Section of Public Health, School of Health and Related Research (SchHARR), The University of Sheffield, UK<sup>5</sup>

\*For Correspondence: E-mail: [ziliyasu@yahoo.com](mailto:ziliyasu@yahoo.com); Phone: +2348035868293

## Abstract

Practices related to resumption of coitus after childbirth remains poorly documented in Nigeria. This study examined factors associated with sexual intercourse, delivery-coitus interval, and contraceptive use among postpartum women attending a tertiary centre in Kano, northern Nigeria. A cross section of 317 women attending immunization, postnatal and family planning clinics within 12 months of childbirth was interviewed using a structured questionnaire. Vaginal intercourse was resumed by most women ( $n=212$ ; 66.9%, 95% Confidence Interval (CI) = 61.8%, 72.2%) with delivery-coitus resumption interval (mean  $\pm$  SD) of  $9.6 \pm 5.2$  weeks postpartum. The majority (67.9%) resumed sexual activity within 8 weeks of delivery. Nearly two-thirds 65.6% ( $n=139/212$ ) of the sexually active women reported current use of contraceptives. Onset of postpartum sexual activity was independently associated with mode of delivery adjusted odds ratio (AOR) (95%CI)= 1.10 (1.03,1.78), baby's age AOR (95%CI) =2.10 (1.27, 8.70), number of living children AOR (95%CI)=1.21 (1.07,1.79), onset of menstruation AOR (95%CI)=0.34 (0.17,0.69) and co-habitation AOR (95%CI)=0.47 (0.016, 0.14). Contraceptive use was predicted by educational status, sexual activity, baby's age and menstruation. Most women attending maternal and child health clinics resumed sexual intercourse within 2 months of delivery, but only two-thirds used modern contraceptive methods. Contraceptive counseling should commence early, preferably during pregnancy. (*Afr J Reprod Health 2018; 22[1]:103-112*).

**Keywords:** sexual intercourse, postpartum, contraceptive use, predictors, Nigeria

## Résumé

Les pratiques liées à la reprise du coït après l'accouchement restent peu documentées au Nigeria. Cette étude a examiné les facteurs associés aux rapports sexuels, à l'intervalle entre accouchement et à l'utilisation de contraceptifs chez les femmes post-partum qui fréquentent un centre tertiaire à Kano, dans le nord du Nigéria. Un échantillon de 317 femmes qui fréquentent des cliniques de vaccination, postnatales et de planification familiale dans les 12 mois suivant l'accouchement a été interviewé à l'aide d'un questionnaire structuré. Les rapports sexuels par voie vaginale ont été repris par la plupart des femmes ( $n = 212$ , 66,9%, 95% Intervalle de confiance (IC) = 61,8%, 72,2%) avec intervalle de reprise de l'accouchement-coït (moyenne  $\pm$  écart-type) de  $9,6 \pm 5,2$  semaines après l'accouchement. La majorité (67,9%) a repris l'activité sexuelle dans les 8 semaines suivant l'accouchement. Près des deux tiers 65,6% ( $n = 139/212$ ) des femmes sexuellement actives ont déclaré avoir utilisé des contraceptifs. Le début de l'activité sexuelle post-partum était indépendamment associé au mode d'accouchement modifié odds ratio (95% CI) = 1,10 (1,03,1,78), l'âge de bébé AOR (IC à 95%) = 2,10 (1,27-8,70), le nombre de enfants AOR (IC 95%) = 1,21 (1,07,1,79), début de la menstruation AOR (IC à 95%) = 0,34 (0,17-0,69) et AOR de cohabitation (IC à 95%) = 0,47 (0,016, 0,14). L'utilisation de la contraception a été prédite par le statut éducatif, l'activité sexuelle, l'âge du bébé et la menstruation. La plupart des femmes qui fréquentent les centres de santé maternelle et infantile ont repris leurs relations sexuelles dans les deux mois suivant l'accouchement, mais seulement les deux tiers ont utilisé des méthodes contraceptives modernes. La consultation de contraceptif devrait commencer tôt, de préférence pendant la grossesse. (*Afr J Reprod Health 2018; 22[1]: 103-112*).

**Mots-clés:** rapports sexuels, post-partum, utilisation de contraceptifs, indice, Nigeria

## Introduction

Childbirth is a unique but challenging experience. For the mother it exerts profound physical, mental and emotional strain, which could influence health and the time to resumption of sexual activities<sup>1,2</sup>. Yet the postnatal period extending up to six weeks (42 days) after birth receives less attention from health care providers than pregnancy and childbirth<sup>3</sup>. Postpartum sexual changes may not be easy to discuss, for both patient and provider. Yet sexual concerns are common, and the majority of patients welcome help from their primary care provider<sup>4</sup>.

Previously, women were culturally obliged to stay away from their husbands to breastfeed for as long as 2-3 years<sup>5</sup>. Recent reports indicate that duration of postpartum abstinence is becoming considerably shortened to an average of six to eight weeks<sup>5,6</sup>. Most women resumed intercourse in the early postnatal period while a small proportion delayed till up to six months<sup>5,6</sup>. Many gynecologists now recommend waiting for 4 to 6 weeks after delivery to allow cessation of lochial loss and healing of episiotomies, tears or other birth-related perineal injuries<sup>7,8</sup>. Although the delivery-coitus resumption interval has been reported from different countries, including Nigeria<sup>5,9</sup>, we are not aware of any such study from Kano, a highly populous and culturally distinct city in northern Nigeria, where polygamy is common, and the region has consistently reported the highest fertility preferences, total fertility rate and lowest contraceptive uptakes in the country<sup>10-12</sup>.

This study determined the prevalence of postpartum sexual activity, delivery-coitus interval, and their determinants among women who delivered within 12 months of the study and attended the postnatal/family planning and child welfare/immunization clinics at Aminu Kano Teaching Hospital in Kano, Nigeria. In addition, the study assessed sexual morbidity, contraceptive use and associated predictive factors. The findings could inform health workers' counseling protocols and treatment readiness.

## Methods

### *Setting/study population*

The study was conducted among recently delivered women ( $\leq 12$  months of study initiation) attending the postnatal clinic, family planning and immunization/child welfare clinics of Aminu Kano Teaching Hospital (AKTH), Kano, Nigeria from January to February 2015. Immunization services are offered free of charge, but mothers attending the other clinics pay consultation fees and need not to have delivered at AKTH. AKTH is a 500-bed tertiary facility that serves as a referral centre for more than 9 million people in Kano and neighboring states. Majority of the patients are from the middle socioeconomic strata and are of the indigenous Hausa-Fulani ethnic group, although Igbo and Yoruba groups also constitute a substantial proportion of attendees<sup>13</sup>.

### *Study design/population*

A cross-sectional study design was used. The minimum sample size of 337 was estimated using the hypothesis testing method<sup>14</sup>, prevalence of sexual intercourse in the postpartum period in a previous study (67.6%)<sup>9</sup> and error rate of 5%. The estimated sample size was increased by 10% to account for anticipated subject non-response, giving a final sample size of 371. Mothers attending the postnatal/family planning clinics and those who brought their children for immunization were eligible. Women who 1) had stillbirths or 2) were medically advised to abstain from sexual intercourse or 3) who delivered  $>12$  months earlier were excluded.

A multistage sampling method was used. In the first stage, sample sizes were proportionately allocated based on attendance at the postnatal, family planning and immunization/child welfare clinics. In the second stage a systematic sampling technique was used in each clinic to select the study subjects. A sampling interval was first determined using the clinic registers as sampling frame and the assigned sample size.

Sequential serial numbers were then allocated to the women as they registered in the three clinics. Women whose serial numbers coincided with the sampling process were invited for an interview after obtaining informed consent. The first respondent in each clinic was selected at random between 1 and the clinic's sampling interval. Thereafter, subsequent respondents were identified by adding the sampling interval to the serial number of the previous respondent. Four Hausa-speaking female medical students were trained to conduct the interviews. The training included reading through the questionnaire, standardization of translations, interview techniques and the process of sampling and obtaining informed consent. Ethical clearance was obtained from the AKTH ethics committee.

### ***Instruments/data collection***

A pre-tested structured interviewer-administered questionnaire adapted from previous studies<sup>9,12</sup> was used to collect data. The modified questionnaire had five sections. The first section sought information on socio-demographic characteristics. The second section elicited obstetric history including parity, breastfeeding status and onset of menstruation after the delivery. The third section enquired about resumption of vaginal intercourse and if so, the interval between delivery and resumption of coitus. Participants were also asked about current modern contraceptive use. Motivations for resumption of sexual intercourse or abstinence, sexual morbidity/problem(s) encountered and health seeking behavior were also ascertained. The questionnaire was translated from English Language into the local (Hausa) language and back-translated to check for consistency. Female medical students were trained to conduct the interviews in a culturally acceptable manner. The questionnaire was validated by pre-testing on 30 postpartum women attending postnatal care in a different hospital (Murtala Muhammad Specialist Hospital) to ascertain the appropriateness of the items included in the different measures.

### ***Measurements***

Although, sexual activity refers to engagement in any form of act intended to arouse or gratify the sexual interest of an individual such as touching, kissing, vaginal intercourse and other mutually pleasurable techniques<sup>15</sup>, in order to avoid ambiguity and ensure cultural sensitivity this study focused on resumption of vaginal intercourse. Contraception was also considered as informed voluntary adoption of methods of spacing, deferment or limiting child birth by the woman or her partner<sup>16</sup>.

Measurements were operationalized as used previously<sup>9,12</sup> as follows: Respondents were categorized as having resumed sexual intercourse if they report having had vaginal intercourse following the last delivery. The delivery-coitus interval was recorded in weeks. The current use of modern contraceptive methods had 'Yes' or 'No' response options. A woman was said to have sexual morbidity if she reported one or more sexual problems on resumption of sexual intercourse (e.g., vaginal discharge, vaginal irritation, lack of libido, dyspareunia, etc.). The response options for motivation for resumption of sexual intercourse were categorized as 'husband's request', 'initiation by self', 'cultural demand' and 'feeling it was convenient'.

### ***Data analysis***

Data was analyzed using SPSS (version 21)<sup>17</sup>. Incomplete, inaccurate and inconsistent data were excluded pre-analysis following data cleaning. Frequencies and simple percentages were used to describe categorical variables. Measures of central tendency and variability were employed to describe quantitative variables, depending on the distribution of the variable. Bivariate analysis was performed using Pearson's Chi-square test, Chi-square test for trend or Fisher Exact test as appropriate. Differences between means were tested using the Student t-test. Type I error was fixed at 5% for all tests of significance. Variables with  $P < 0.05$  on bivariate analysis were

included in a multivariate logistic model to identify independent predictors of sexual intercourse after the last child birth. Adjusted odds ratios generated from the logistic models were used to quantify magnitude of relationships/associations.

## Results

Of the 371 women approached, 317 completed the interviews, giving a response rate of 85.4%. Participants' mean age  $\pm$  standard deviation (SD) was  $27.0 \pm 5.4$  years. Participants were predominantly Hausa (51.1%,  $n=162$ ) or Fulani (25.2%,  $n=80$ ) and Muslim (82.0%,  $n=260$ ). Of those that were married (89.0%,  $n=282$ ), a third (33%,  $n=94$ ) were in polygamous unions. Median (range) duration of marriage was 6 years (1-29 years) and majority (89.7%,  $n=253/282$ ) were cohabiting with their husbands at the time of the study. About a third of respondents had secondary (32%,  $n=101$ ) or post-secondary (31%,  $n=98$ ) education. Almost half (48%,  $n=152$ ) of the respondents were stay at home mothers.

### Obstetric characteristics

Median parity (range) was two (one to nine) with mean ( $\pm$ SD) age of babies of  $4.9 \pm 3.3$  months. Nearly two-thirds (61%  $n=193$ ) of the participants had spontaneous vaginal delivery, 23% ( $n=73$ ) had caesarean section while 16.1% ( $n=51$ ) had instrumental vaginal delivery. More than half (52.1%,  $n=165$ ) of the participants had episiotomies done. In addition, (15.1%,  $n=48$ ) women sustained genital injuries. The most common injuries were vaginal lacerations (70.8%,  $n=34/48$ ) and perineal tears (25.0%,  $n=12/48$ ). More than three-fourths (78%,  $n=247$ ) of the women were breastfeeding their babies at the time of the study and most of the women (61%,  $n=193$ ) had resumed menstruation.

### Resumption of sexual intercourse postpartum

Two thirds of the respondents (66.9%,  $n=212/317$ ) (95% CI=61.8% to 72.2%) had resumed sexual

intercourse after their last delivery. The interval between delivery and resumption of coitus ranged from 4 weeks to 28 weeks (mean $\pm$ SD=9.6 $\pm$ 5.2 weeks). Of those who had sexual intercourse after delivery ( $n=212$ ), the proportions resuming <6 weeks, 6 to <8 weeks, 8 to <12 weeks, 12 to  $\leq$ 28 weeks were 21.7%, 46.2%, 16.5%, and 15.6% respectively. Motivations for resumption of coitus were mainly 'husband's demand' (67.5%,  $n=143/212$ ), and self-wish (14.6%,  $n=31$ ). In contrast, the main reasons proffered by sexually abstinent women (33.0%,  $n=105$ ) included 'baby too young' (81.0%,  $n=85$ ), 'don't want to get pregnant just yet' (13.3%,  $n=14$ ), 'not interested' (2.9%,  $n=3$ ) and 'partner not available as a result of divorce or widowhood' (2.9%,  $n=3$ ).

### Sexual morbidity

Almost two-thirds 64.2% ( $n=136/212$ ) of respondents who had resumed sexual intercourse reported a sexual problem at some point. These included reduced sexual desire (31.6%,  $n=67$ ), dyspareunia (32.5%,  $n=69$ ), vaginal dryness/soreness (15.6%,  $n=33$ ), vaginal 'looseness' (15.6%,  $n=33$ ) and vaginal discharge (6.1%,  $n=13$ ). However, less than two-thirds of those who experienced sexual morbidity (63.2%,  $n=86/136$ ) sought help. The main sources of help were friends 70.9% ( $n=61/86$ ), mother 29.1% ( $n=25/86$ ), doctor 11.6% ( $n=10/86$ ), or the internet 8.1% ( $n=7/86$ ). Reasons for not seeking advice included shyness (42.0%  $n=21/50$ ), problem resolved (52.0%,  $n=26/50$ ), cultural/religious factors (26.0%,  $n=13/50$ ), and 'no female doctor' (16.0%,  $n=8/50$ ).

### Contraceptive use

Of respondents that resumed coitus ( $n=212$ ), almost two-thirds 65.6% ( $n=139$ ) were currently using modern contraceptives. The popular methods were: injectables (33.1%,  $n=46/139$ ), implants (21.6%,  $n=30/139$ ), oral contraceptives (20.1%,  $n=28/139$ ), and intrauterine device (10.8%,  $n=15/139$ ). Less popular ones included the rhythm method (5.0%,  $n=7/139$ ), male condom

**Table 1:** Socio-demographic and Obstetric Characteristics of Postpartum Women in Kano, Nigeria

| Characteristics               | Frequency   |
|-------------------------------|-------------|
| <b>Age groups (years)</b>     | No. (%)     |
| <25                           | 115 (36.2)  |
| 25-29                         | 101 (31.9)  |
| ≥30                           | 101 (31.9)  |
| Total                         | 317 (100)   |
| <b>Education</b>              |             |
| No formal                     | 65 (20.5)   |
| Primary                       | 51 (16.1)   |
| Secondary                     | 103 (32.5)  |
| Tertiary                      | 98 (30.9)   |
| Total                         | 317 (100.0) |
| <b>Ethnicity</b>              |             |
| Hausa                         | 162 (51.1)  |
| Fulani                        | 80 (25.2)   |
| Igbo                          | 29 (9.1)    |
| Yoruba                        | 25 (7.9)    |
| Others                        | 21 (6.6)    |
| Total                         | 317 (100.0) |
| <b>Religion</b>               |             |
| Muslims                       | 260 (82.0)  |
| Christians                    | 57 (18.0)   |
| Total                         | 317 (100.0) |
| <b>Marital Status</b>         |             |
| Married                       | 282 (88.9)  |
| Widowed/Divorced              | 35 (11.1)   |
| Total                         | 317 (100.0) |
| <b>Occupation</b>             |             |
| Stay at home mothers          | 152 (48.0)  |
| Traders                       | 92 (29.0)   |
| Civil servant                 | 25 (8.0)    |
| Others                        | 48 (15.0)   |
| Total                         | 317 (100.0) |
| <b>No. of living children</b> |             |
| 0-1                           | 90 (28.4)   |
| 2-4                           | 183 (57.7)  |
| ≥5                            | 44 (13.9)   |
| Total                         | 317 (100.0) |
| <b>Mode of delivery</b>       |             |
| Spontaneous vaginal           | 199 (62.8)  |
| Caesarean section             | 73 (23.0)   |
| Instrumental                  | 45 (14.2)   |
| Total                         | 317 (100.0) |
| <b>Baby's age (months)</b>    |             |
| <1                            | 47 (14.8)   |
| 1-6                           | 169 (53.3)  |
| >6-12                         | 101 (39.9)  |
| Total                         | 317 (100.0) |

(3.6%,  $n=5/139$ ), female condom (2.9%,  $n=4/139$ ), and female sterilization (2.9%,  $n=4/139$ ).

### **Predictors of postpartum resumption of sexual intercourse and contraceptive use**

At bivariate level, postpartum sexual intercourse was significantly associated with respondent's age,

education, co-habitation, duration of marriage, number of living children, mode of delivery, baby's age and resumption of menstruation ( $P<0.05$ ). After adjusting for confounders using logistic regression analysis, number of living children, mode of delivery, baby's age, resumption of menstruation and co-habitation remained significant predictors (Table 2).

At bivariate level contraceptive use was significantly associated with respondent's age, education, duration of marriage, number of living children, resumption of sexual intercourse, baby's age, and resumption of menstruation ( $P<0.05$ ). After adjusting for potential confounders, education, baby's age, resumption of sexual intercourse, and menstruation remained independent predictors (Table 3).

## **Discussion**

The proportion of women that had resumed sexual intercourse after delivery (66.9%) was comparable with reports from other parts of Nigeria [Jos (67.6%)<sup>9</sup>, Sagamu (65%)<sup>5</sup>] and from Kampala, Uganda (66.4%)<sup>18</sup>. However, it is higher than the figures found in Ghana (23.8%)<sup>19</sup> and Germany (47%)<sup>20</sup> but lower than the those from the UK (89%)<sup>21</sup> and USA (90%)<sup>22</sup>. The mean time to resumption of coitus postpartum was longer than the reported figures from Uganda (7.06 weeks)<sup>18</sup> and Turkey (7.87 weeks)<sup>23</sup>. Most women in this study had resumed sexual intercourse within a fortnight after puerperium.

Apart from variations in follow up period (range; 3 to 12 months) and study populations, the differences between our findings and others likely reflect tradition, culture or religious peculiarities.

Sexual intercourse within the puerperium (<6 weeks post-delivery) was not rare (21.7%), as was earlier reported from Ekiti (27.6%), south west Nigeria<sup>24</sup>. An even higher proportion of women resumed sexual intercourse soon after the puerperium in Nnewi (48.7%), south east Nigeria<sup>25</sup>. The puerperium is usually followed by a postnatal check commonly at 6 weeks when women are examined to ensure that episiotomies and perineal tears are healed, thereby re-assuring couples that all is well<sup>26</sup>. With this re-assurance, couples are more likely to resume sexual activity soon afterwards. The tendency towards early

**Table 2:** Predictors of Postpartum Sexual Intercourse in Women in Kano, Nigeria

| Characteristics                     | Sexual activity |             | Crude OR (95% CI) | Adjusted OR(95%CI) | P-value |
|-------------------------------------|-----------------|-------------|-------------------|--------------------|---------|
|                                     | No. (%) Resumed | Not resumed |                   |                    |         |
| <b>Age Group</b>                    |                 |             |                   |                    |         |
| <25                                 | 56(48.7)        | 59(51.3)    | 1.00              |                    |         |
| 25-29                               | 72 (71.3)       | 29 (28.7)   | 1.46 (1.17-1.83)* | 2.47 (0.83-7.34)   | 0.10    |
| ≥30                                 | 78 (77.2)       | 23 (22.8)   | 1.59 (1.28-1.97)* | 1.63 (0.60-4.48)   | 0.34    |
| <b>Education</b>                    |                 |             |                   |                    |         |
| No formal                           | 35(53.8)        | 30(46.2)    | 1.00              |                    |         |
| Primary                             | 28 (54.9)       | 23 (45.1)   | 1.02 (0.73-1.43)  | 1.50 (0.59-3.80)   | 0.40    |
| Secondary                           | 69 (67.0)       | 34(33.0)    | 1.24 (0.96-1.62)  | 0.82 (0.28-2.38)   | 0.71    |
| Tertiary                            | 74 (75.5)       | 24(24.5)    | 1.40 (1.09-1.80)* | 1.11 (0.49-2.50)   | 0.80    |
| <b>Co-habitation with husband</b>   |                 |             |                   |                    |         |
| Yes                                 | 194 (68.3)      | 90(31.7)    | 1.00              |                    |         |
| No                                  | 12 (36.4)       | 21(63.6)    | 0.53 (0.34-0.84)* | 0.47 (0.016-0.14)  | 0.001*  |
| <b>Mode of delivery</b>             |                 |             |                   |                    |         |
| Spontaneous vaginal                 | 139(69.8)       | 60(30.2)    | 1.43 (1.04-1.95)* | 1.10 (1.03-1.78)   | 0.05*   |
| Caesarean                           | 45 (61.6)       | 28(38.4)    | 1.26 (0.89-1.79)  | 0.46 (0.19-1.11)   | 0.084   |
| Instrumental                        | 22(48.9)        | 23(51.1)    | 1.00              |                    |         |
| <b>Baby's age (months)</b>          |                 |             |                   |                    |         |
| <1                                  | 17(36.2)        | 30(63.8)    | 1.00              |                    |         |
| 1-6                                 | 104(61.5)       | 65(38.5)    | 1.70 (1.14-2.53)* | 1.53 (1.22-7.52)   | 0.017*  |
| >6-12                               | 85(84.2)        | 16(15.8)    | 2.33 (1.58-3.43)* | 2.10 (1.27-8.70)   | 0.001*  |
| <b>Duration of marriage (years)</b> |                 |             |                   |                    |         |
| ≤2                                  | 31(42.5)        | 42(57.5)    | 1.00              |                    |         |
| >2-10                               | 137(72.1)       | 53(27.9)    | 1.70 (1.28-2.25)* | 1.06 (0.25-4.41)   | 0.94    |
| >10                                 | 38(70.4)        | 16(29.6)    | 1.66 (1.21-2.28)* | 0.35 (0.11-1.07)   | 0.065   |
| <b>No. of living children</b>       |                 |             |                   |                    |         |
| 0-1                                 | 47(52.2)        | 43(47.8)    | 1.00              |                    |         |
| 2-4                                 | 126(68.9)       | 57(31.1)    | 1.32 (1.06-1.64)* | 1.10 (0.53-2.29)   | 0.79    |
| ≥5                                  | 33(75.0)        | 11(25.0)    | 1.44 (1.11-1.86)* | 1.21 (1.07-1.79)   | 0.034*  |
| <b>Resumption of menstruation</b>   |                 |             |                   |                    |         |
| Yes                                 | 152 (80.4)      | 37 (19.6)   | 1.00              |                    |         |
| No                                  | 54 (42.2)       | 74 (57.8)   | 0.52 (0.42-0.65)* | 0.34 (0.17-0.69)   | 0.003*  |

OR=Odds Ratio; 95%CI=95% Confidence Interval; \*=Significant at P≤0.05

coital resumption following child birth contrasts with the long established culture of prolonged abstinence and 1-2 years of breastfeeding which protected against pregnancy in traditional cultures<sup>5,27</sup>. While most gynecologists recommend waiting for 4 to 6 weeks after delivery to allow cessation of lochial loss and healing of episiotomies, tears or other birth-related perineal injuries<sup>7,8</sup>, the importance of preventing unplanned pregnancy soon after childbirth cannot be stressed enough. Although the uptake of contraceptives seems to be higher than in the general populace, the effects of selection factors cannot be ruled out,

whereby women presenting to tertiary centers were more likely to be educated and of higher socioeconomic status. The link between education/socioeconomic status and contraceptive use is consistent<sup>28-30</sup>.

Apart from maintaining marital harmony, some of our respondents considered early resumption of sexual intercourse as a strategy to prevent their partners from seeking sexual gratification elsewhere, a logical presumption. However, sexual intercourse increases the risk of unwanted pregnancy in the absence of contraception. Predominance of husband-initiated

**Table 3:** Predictors of Postpartum Contraceptive use in Women in Kano, Nigeria

| Characteristics                     | Contraceptive use |            | Crude OR (95% CI) | Adjusted OR(95%CI) | P-value |
|-------------------------------------|-------------------|------------|-------------------|--------------------|---------|
|                                     | Yes               | No         |                   |                    |         |
| <b>Age group</b>                    |                   |            |                   |                    |         |
| <25                                 | 22 (19.1)         | 93 (80.9)  | 1.00              |                    |         |
| 25-29                               | 44 (43.6)         | 57 (56.4)  | 2.28 (1.47-3.52)* | 0.96 (0.46-1.99)   | 0.90    |
| ≥30                                 | 53 (52.5)         | 48 (47.5)  | 2.74 (1.80-4.17)* | 2.05 (0.87-4.84)   | 0.10    |
| <b>Education</b>                    |                   |            |                   |                    |         |
| No formal                           | 18 (27.7)         | 47 (72.3)  | 1.00              |                    |         |
| Primary                             | 12 (23.5)         | 39 (76.5)  | 0.85 (0.45-1.60)  | 1.78 (0.79-4.03)   | 0.16    |
| Secondary                           | 40 (38.8)         | 63 (61.2)  | 1.40 (0.88-2.23)  | 1.35 (0.70-2.58)   | 0.37    |
| Tertiary                            | 49 (50.0)         | 49 (50.0)  | 1.81 (1.16-2.80)* | 2.62 (1.03-6.69)   | 0.043*  |
| <b>Resumed sexual activity</b>      |                   |            |                   |                    |         |
| Yes                                 | 107 (51.9)        | 99 (48.1)  | 1.00              |                    |         |
| No                                  | 12 (10.8)         | 99 (89.2)  | 0.21 (0.12-0.36)* | 0.18 (0.088-0.38)  | 0.001*  |
| <b>Baby's age (months)</b>          |                   |            |                   |                    |         |
| ≤1                                  | 12 (25.5)         | 35 (74.5)  | 1.00              |                    |         |
| >1-6                                | 50 (29.6)         | 119 (70.4) | 1.16 (0.67-1.99)  | 1.34 (0.50-3.61)   | 0.56    |
| >6-12                               | 57 (56.4)         | 44 (43.6)  | 2.21 (1.31-3.71)* | 1.93 (1.06-3.49)   | 0.031*  |
| <b>Duration of marriage (years)</b> |                   |            |                   |                    |         |
| ≤2                                  | 14 (19.2)         | 59 (80.8)  | 1.00              |                    |         |
| >2-10                               | 76 (40.0)         | 114 (60.0) | 2.09 (1.26-3.45)* | 2.82 (0.82-9.68)   | 0.10    |
| >10                                 | 29 (53.7)         | 25 (46.3)  | 2.80 (1.64-4.77)* | 2.18 (0.79-5.96)   | 0.13    |
| <b>No. of living children</b>       |                   |            |                   |                    |         |
| 0-1                                 | 21 (23.3)         | 69 (76.7)  | 1.00              |                    |         |
| 2-4                                 | 76 (41.5)         | 107 (58.5) | 1.78 (1.18-2.69)* | 0.99 (0.3-3.30)    | 0.99    |
| ≥5                                  | 22 (50.0)         | 22 (50.0)  | 2.14 (1.33-3.45)* | 0.73 (0.27-2.02)   | 0.54    |
| <b>Resumption of menstruation</b>   |                   |            |                   |                    |         |
| Yes                                 | 96 (50.8)         | 93 (49.2)  | 1.00              |                    |         |
| No                                  | 23 (18.0)         | 105 (82.0) | 0.35 (0.24-0.53)* | 0.40 (0.21-0.75)   | 0.004*  |

OR=Odds Ratio; 95%CI=95% Confidence Interval; \*=Significant at  $P \leq 0.05$

resumption of coitus (77.4%) in the present study concurs with reports from Jos, Nigeria<sup>9</sup> and from Uganda<sup>18</sup> and Cote d'Ivoire<sup>31</sup>. It is culturally untenable for such requests to be turned down in our study setting where 22.7% of women agreed that a husband is justified in hitting or beating his wife when she refuses to have sexual intercourse with him<sup>12</sup>. This contrasts with the developed world where women are more assertive.

Men's interest in early resumption of sexual activity highlights the need to engage them in discussions about appropriate timing of sexual activities after child birth, unplanned pregnancy, and contraceptive use well before the birth of the baby. Such couple-centered prenatal counseling could allay anxiety and encourage informed choices about when to safely and comfortably

resume sexual activity. Health workers also need the capacity and confidence to initiate such sensitive discussions. Secondly, a tenth of the women reportedly resumed sexual intercourse on their own accord. This is unusual as sexual norms and assumptions in many African settings do not expect women to want sex or find it pleasurable<sup>9</sup>. Nonetheless, effects of cultural inhibitions are by no means uniform even in the same part of the country<sup>9,32</sup>.

A third of women were yet to resume sexual intercourse after delivery, mostly because the baby was too young or they were afraid of getting pregnant too soon. This contrasts with the reasons proffered by women in Jos, Nigeria<sup>9</sup> and Kampala, Uganda<sup>18</sup>. In both instances the main reason was non-availability of the husband.

Elsewhere, in Ontario, Canada<sup>33</sup>, decline in sexual desire was the main reason for non-resumption of coitus postpartum. This difference may be due to the patriarchal nature of the study setting, where women are less assertive compared to developed countries where women are empowered and aware of their sexual and reproductive rights<sup>21, 33</sup>. The other reasons given for abstinence from coitus are like earlier studies<sup>18,25</sup>.

The proportion of women who encountered at least one sexual problem on resumption (40%) is higher than the figures reported from Uganda (22.2%)<sup>18</sup> and the Gambia (27%)<sup>34</sup>, but less than the numbers from Turkey (58.3%)<sup>23</sup>, Australia (64.3%)<sup>35</sup>, the UK (83%)<sup>21</sup>, and Canada (96.0%)<sup>33</sup>. Apart from differences in study populations, methods and measurements, these figures could have been influenced by under-reporting occasioned by cultural restraints and self-limiting nature of some of the symptoms. Cultural inhibition and lack of encouragement on the part of health providers could further explain the proportion of women who reportedly discussed the symptoms with health providers<sup>5,9,18</sup>. The range of symptoms was in concordance with earlier studies<sup>5,18,25</sup>. Health care providers should routinely counsel and encourage couples, especially women, to voice out sexual concerns during the postpartum period so that they can be effectively addressed. Existing guidelines could be adopted to reflect local cultural sensitivities and individual preferences<sup>36</sup>.

More than a third (38%) of the women in this study were currently using a modern method. This is much higher than figures reported from population-based surveys (10%)<sup>12</sup> and postpartum women attending other Nigerian hospitals in Jos (19.1%)<sup>9</sup>, and Enugu (14.7%)<sup>37</sup>. Apart from variations in study population, methods and measures of the studies, this unusually high level of contraceptive uptake could be attributed to the correspondingly high proportion of women with formal education and the dominance of women from middle and high socioeconomic strata among our study participants. These factors are known to influence contraceptive use<sup>12</sup>. Regardless of the relatively higher contraceptive uptake rate in this study compared to other studies, the fact that one-third of respondents were not using any modern

contraceptive method despite early resumption of sexual intercourse in the postpartum highlights the need for prenatal counseling on family planning options.

The popular methods among the respondents were similar to the findings of the 2013 Demographic Health Survey, where injectables were the favorite among married women<sup>12</sup>. The higher contraceptive prevalence among women that had resumed menstruation could be due to the use of menstruation as an external sign of fecundity by health care providers and lay public alike<sup>38</sup>. It is a common practice to ask women to return for family planning services after resumption of menstruation<sup>39</sup>.

The predictive effects of mode of delivery, baby's age, number of living children, menstruation, and co-habitation on resumption of coitus postpartum have been similarly reported<sup>5,18,38</sup>. The differences in periods of recuperation following different modes of delivery, pain and tissue injury that accompany instrumental and caesarean delivery and pre-morbid conditions of the women could account for variations in time of resumption of sexual activity.

Baby's age corresponds to the time since delivery. The longer this duration, the more likely that episiotomies and vaginal lacerations have healed. In addition, the older the baby, the more hopeful the mother is that her child would survive even if another pregnancy occurs. Sexual desire is also expected to improve the longer the interval from time of delivery. Women of high parity are more experienced and are more likely to have developed a timeline from previous births.

Unsurprisingly, contraceptive use was significantly associated with educational status, resumption of sexual intercourse, baby's age, and resumption of menstruation. The influence of these factors has been reported earlier<sup>9</sup>. Education empowers women to plan their lives while the remaining three factors increase the risk of pregnancy. Therefore, girl-child education and economic empowerment are important long-term measures for preventing unwanted and unplanned pregnancies and their consequences.

This study was limited by at least two factors. First, selection factors at the study clinics restricted the extent to which the results could be



generalized. Being a tertiary centre, rural dwelling women and those of low literacy and socio-economic status were under-represented. Secondly, sexual issues are still taboo in the study setting. This is more so in the postpartum period when women are expected to be abstinent and to concentrate on the care of the baby. Therefore, socially desirable responses may be given. However, one-on-one confidential interviews conducted by trained female interviewers from the same culture reduced the social barrier.

## Conclusion

In conclusion, majority of women in this study setting were sexually active postpartum. The time of resumption of sexual activity varied widely among couples, and was mainly influenced by partner's predisposition, socio-demographic and obstetric factors. Contraceptive uptake was higher than in the general populace and was also strongly influenced by educational status and obstetric characteristics. Couple-centered counseling and discussion about postpartum sexuality, pregnancy risks, and contraception could allay anxiety and inform individual choices about timing of sexual activity following child birth. A detailed assessment of sexual function using a standardized tool and qualitative interviews could shed more light on postpartum sexual behavior in this setting.

## Contribution of Authors

Zubairu Iliyasu, Hadiza S Galadanci and Khadija M. Danlami conceived and designed the study, Khadija M. Danlami and Hadiza S Galadanci collected the data while Zubairu Iliyasu, Hadiza S Galadanci, Hamisu M Salihu and Muktar H Aliyu analysed the data and prepared the draft manuscript. All authors reviewed and approved of the final manuscript.

## References

1. Connolly A, Thorp J and Pahal L. Effects of pregnancy and childbirth on postpartum sexual function: a longitudinal prospective study. *IntUrogynecol J Pelvic Floor Dysfunct*2005 16(4):263-7
2. Handa VL. Sexual function and childbirth. *SeminPerinatol*2006;30:253-6.
3. Fort, AL, Kothari, MT and Abderrahim N. Postpartum care: levels and determinants in developing countries. Calverton, Maryland, USA, Macro International, 2006 ([http://www.measuredhs.com/pubs/pub\\_details.cfm?ID=676](http://www.measuredhs.com/pubs/pub_details.cfm?ID=676)) accessed 12<sup>th</sup> February 2015
4. Nusbaum MR, Gamble G, Skinner B and Heiman J. The high prevalence of sexual concerns among women seeking routine gynecological care. *J Fam Pract* 2000;49(3):229-32.
5. Dada OA, Akesode FA, Olanrewaju DM, Olowu OA, Sule-Odu O, Fakoya TA, Oluwole, FA and Odunlami BV. Infant feeding and lactational amenorrhoea in Sagamu, Nigeria. *AfrJ Reprod Health*2002;6:39-50.
6. Abdool Z, Thakar R and Sultan AH. Postpartum female sexual function. *Eur J ObstetGynaecol Repro Biol.*2009;145:133-7.
7. Johnson CE. Sexual health during pregnancy and the postpartum. *The Journal of Sexual Medicine.*2011;8:1267.
8. Leeman LM and Rogers RG. Sex after childbirth: Postpartum sexual function. *Obstetrics and Gynecology.* 2012;119:647.
9. Anzaku A and Mikah S. Postpartum resumption of sexual activity, sexual morbidity and use of modern contraceptives among Nigerian women in Jos. *Ann Med Health Sci Res.* 2014 Mar;4(2):210-6. doi: 10.4103/2141-9248.129044.
10. National Population Commission (NPC) [Nigeria] and ORC Macro (2004). Nigeria Demographic and Health Survey 2003. Calverton, Maryland: National Population Commission and ORC Macro.
11. National Population Commission (NPC) [Nigeria] and ICF Macro (2009). Nigeria Demographic and Health Survey 2008. Abuja, Nigeria: National Population Commission and ICF Macro.
12. National Population Commission (NPC) [Nigeria] and ICF International (2014). Nigeria Demographic and Health Survey 2013. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF International.
13. Iliyasu Z and Babashani M. Prevalence and predictors of tuberculosis co-infection among HIV-seropositive patients attending the Aminu Kano Teaching Hospital, northern Nigeria. *J Epidemiol.* 2009;19(2):81-7.
14. Lwanga SK and Lemeshow S. Sample size determination in health studies: a practical manual. World Health Organization 1991:29-32
15. WHO. Defining Sexual Health. [http://www.who.int/reproductivehealth/topics/sexual\\_health/sh\\_definitions/en/](http://www.who.int/reproductivehealth/topics/sexual_health/sh_definitions/en/) accessed 18<sup>th</sup> February 2015
16. WHO. Contraceptive prevalence. [http://www.who.int/reproductivehealth/topics/family\\_planning/contraceptive\\_prevalence/en/](http://www.who.int/reproductivehealth/topics/family_planning/contraceptive_prevalence/en/) accessed 19<sup>th</sup> February 2015
17. IBM Corp. (2012). IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY, USA: IBM Corporation.
18. Odar E, Wandabwa J and Kiondo P. Sexual practices of

- women within six months of childbirth in Mulago hospital, Uganda. *Afr Health Sci* 2003;3:117-23.
19. Borda MR, Winfrey W and McKaig C. Return to sexual activity and modern family planning use in the extended postpartum period: An analysis of findings from seventeen countries. *Afr J Reprod Health* 2010;14:72-9.
  20. Buhling KJ, Schmidt S, Robinson JN, Klapp C, Siebert G and Dudenhausen JW. Rate of dyspareunia after delivery in primiparae according to mode of delivery. *Eur J Obstet Gynecol Reprod Biol* 2006;124:42-6.
  21. Barret G, Pendry E, Peacock J, Victor C, Thakar R and Manyonda I. Women's sexual health after childbirth. *BJOG* 2000;107:186-95.
  22. Brubaker L, Handa VL, Bradley CS, Connolly A, Moalli P, Brown MB and Weber A. Sexual function 6 months after first delivery. *Obstet Gynecol* 2008;111:1040-4.
  23. Acele EÖ and Karaçam Z. Sexual problems in women during the first postpartum year and related conditions. *J Clin Nurs* 2012;21:929-37.
  24. Adanikin AI, Awoleke JO, Adeyiolu A, Alao O and Adanikin PO. Resumption of intercourse after childbirth in southwest Nigeria. *The European Journal of Contraception and Reproductive Health Care* 2015;20(4):241-248.
  25. Adinma JI. Sexual activity during and after pregnancy. *Adv Contracept* 1996;12:769-75.
  26. Olsson A, Robertson E, Falk K and Nissen E. Assessing women's sexual life after childbirth: the role of the postnatal check. *Midwifery*. 2011 Apr;27(2):195-202. doi: 10.1016/j.midw.2009.04.003.
  27. Isenaiumhe AE and Oviawe O. The changing pattern of Post-Partum Sexual Abstinence in a Nigerian Rural community. *Soc. Sci. Med.* 1986, 23 (7): 683 – 686
  28. Olatoregun O, Fagbamigbe AF, Akinyemi OJ, Yusuf OB and Bamgboye EA. A comparative analysis of fertility differentials in Ghana and Nigeria. *Afr J Reprod Health*. 2014 Sep;18(3):36-47.
  29. Lamidi EO. State variations in women's socioeconomic status and use of modern contraceptives in Nigeria. *PLoS One*. 2015;10 (8):e0135172.
  30. Larsson C and Stanfors M. Women's Education, Empowerment, and Contraceptive Use in sub-Saharan Africa: Findings from Recent Demographic and Health Surveys. *African Population Studies* 2014; 28 (2) Suppl: 1022-1034.
  31. Desgrees-du-Lou A and Brou H. Resumption of sexual relations following childbirth: Norms, practices and reproductive health issues in Abidjan, Côte d'Ivoire. *Reprod Health Matters* 2005;13:155-63.
  32. Orisaremi TC. The influence of breastfeeding beliefs on the sexual behavior of the Tarok in north-central Nigeria. *Sex Reprod Healthc.* 2013;4(4):153-60. doi: 10.1016/j.srhc.2013.10.005.
  33. Rowland M, Foxcroft L, Hopman WM and Patel R. Breastfeeding and sexuality immediately post partum. *Can Fam Physician* 2005;51:1366-7.
  34. Walraven G, Scherf C, West B, Ekpo G, Paine K, Coleman R, Bailey R and Morison L. The burden of reproductive-organ disease in rural women in The Gambia, West Africa. *Lancet* 2001;357(9263):1161-7.
  35. Khajehei M, Doherty M, Tilley PJ and Sauer K. Prevalence and risk factors of sexual dysfunction in postpartum Australian women. *J Sex Med.* 2015 Jun;12(6):1415-26. doi: 10.1111/jsm.12901.
  36. Lamont J. Female sexual health consensus clinical guidelines. *J Obstet Gynaecol Can* 2012;34:769-75.
  37. Okeke T, Ugwu E, Ezenyeaku C, Ikeako L and Okezie O. Postpartum practices of parturient women in enugu, South East Nigeria. *Ann Med Health Sci Res.* 2013;3(1):47-50. doi: 10.4103/2141-9248.109486.
  38. Sathiyasusuman A, Bado A and Lailulo YA. Promoting family planning use after childbirth and desire to limit childbearing in Ethiopia. *Reprod Health.* 2014;11:53. doi: 10.1186/1742-4755-11-53.
  39. Wang IY and Fraser IS. Reproductive function and contraception in the postpartum period. *Obstet Gynecol Surv.* 1994 Jan;49(1):56-63.
  40. Ezebialu IU and Eke AC. Resumption of vaginal intercourse in the early postpartum period: determinants and considerations for child spacing in a Nigerian population. *J Obstet Gynaecol.* 2012 May;32(4):353-6. doi: 10.3109/01443615.2011.652701.