

Erratum to: Biosorption of Cu(II) and Pb(II) from aqueous solutions by chemically modified spent coffee grains

F. J. Cerino-Córdova · P. E. Díaz-Flores · R. B. García-Reyes · E. Soto-Regalado ·
R. Gómez-González · M. T. Garza-González · E. Bustamante-Alcántara

Received: 11 March 2013 / Published online: 24 April 2013
© Islamic Azad University (IAU) 2013

Erratum to: Int J Environ Sci Technol
DOI:10.1007/s13762-013-0198-z

The original version of this article unfortunately contained mistakes. The presentations of Figs. 1 and 2 were incorrect.

The correct versions of the figures are given in following page.

The online version of the original article can be found under
doi:[10.1007/s13762-013-0198-z](https://doi.org/10.1007/s13762-013-0198-z).

F. J. Cerino-Córdova (✉) · R. B. García-Reyes ·
E. Soto-Regalado · R. Gómez-González ·
M. T. Garza-González · E. Bustamante-Alcántara
Universidad Autónoma de Nuevo León, UANL, Facultad de
Ciencias Químicas, Av. Universidad S/N, Cd. Universitaria,
66451 San Nicolás de los Garza, NL, Mexico
e-mail: felipejccuanl@yahoo.com.mx

P. E. Díaz-Flores
Universidad Autónoma de San Luis Potosí, Coordinación,
Académica Región Altiplano, Carretera a Cedral Km 5+600 San
José de las Trojes, 78700 Matehuala, SLP, Mexico

Fig. 1 Speciation diagram of **a** Pb(II) and **b** Cu(II) for a total concentration of 1.20 and 4.72 mmol/L, respectively




Fig. 2 ATR-FTIR spectra of non-modified spent coffee grains (NMSCG), and modified spent coffee grains with 0.1 M and 0.6 M citric acid (MSCG-0.1 M and MSCG-0.6 M, respectively)

