

NUTRITIONAL QUALITY OF BEEF PRODUCED IN CHILE FROM DIFFERENT PRODUCTION SYSTEMS

Rodrigo Morales^{1*}, Carolina Folch¹, Sergio Iraira¹, Nolberto Teuber¹, and Carolina E. Realini²

In recent years, beef industry has improved production processes to ensure quality and certification for overseas meat markets. However, there is limited scientific information about the nutritional quality of beef from Chile. The aim of the present study was to evaluate the nutritional quality of beef produced in Chile from different production systems classified according to the type of finishing diet. Two-hundred and five animals from 13 livestock farms were used, 80 steers were finished on grazing pasture, 79 on pasture supplemented with grain or concentrate and 46 steers were finished on a typical feedlot system in the central zone of Chile. *Longissimus thoracis* intramuscular fat, fatty acid composition and cholesterol content were determined. Beef from pasture-fed animals showed higher content of vaccenic acid (C18:1 *trans*-11) rumenic acid (CLA *cis*-9 *trans*-11) and *n*-3 fatty acids, and lower *n*-6:*n*-3 ratio than beef from feedlot animals. However, Chilean beef from the three production systems showed similar intramuscular fat percentages and a *n*-6:*n*-3 ratio lower than 4.0. Results indicate that the *Longissimus* muscle from Chilean beef could be recommended as a source of red meat for a healthy diet.

Key words: Fatty acids, pasture, feedlot, Chilean beef quality.

Producers and consumers are both interested in the nutritional quality of beef, due to the added value producers could obtain for their product and for consumers' willingness to eat healthier food.

It is well documented that the animal finishing diet affects the fat content and fatty acid composition of beef (Scollan *et al.*, 2001; Latimori *et al.*, 2008). Latimori *et al.* (2008) showed that beef from animals raised on pasture has lower fat and cholesterol concentrations, and more polyunsaturated fatty acids (PUFA) than beef from feedlot animals. Other authors have shown that beef from animals fed on pasture contains more *n*-3 fatty acids and conjugated linoleic acid (CLA) than beef from feedlot animals, due to the high content of linolenic acid (*n*-3) in pasture lipids (Christie, 1981; Steen and Porter, 2003; Elgersma *et al.*, 2003). Conjugated linoleic acid is an isomer of linoleic acid and is deposited in beef by direct (ruminal biohydrogenation and isomerization) or indirect (endogenous synthesis) pathways (Ip *et al.*, 1994). Whetsell *et al.* (2003) reported that CLA C18:2 *cis*-9 *trans*-11 (rumenic acid) possess anticarcinogenic and immunostimulant properties. It has also been shown

that CLA exhibits antioxidant properties, inhibiting the discoloration of meat during storage (Du *et al.*, 2000).

In most developed countries and in Chile, heart disease, diabetes and some types of cancer are the principal causes of death, which are associated with high consumption of saturated and *trans* fatty acids, and refined carbohydrates, stress and a sedentary lifestyle (DEIS, 2007). Thus, there is consumer interest in increasing healthy and natural food and reducing processed foods in the diet to decrease the damaging impact of saturated fatty acids on health. Therefore, consumers are demanding healthier food, including beef (Ip *et al.*, 1994; Whetsell *et al.*, 2003; Wood *et al.*, 2003; Grunert *et al.*, 2004).

In Chile, there are different beef production systems according to the finishing diet. In the central part of Chile (32° to 38° S), animals are fed in feedlots using rations produced in the farms, such as corn silage, alfalfa hay and grain soiling. Waste by-products from different agro-industries, such as poultry feces as well as fruit and vegetable by-products, are normally used as feed in Chilean feedlots (Claro and González, 2005). Grain usually represents between 20 and 40% of the total diet in Chilean feedlots. In contrast, beef production in the southern regions of Chile (38° to 41° S) relies on direct grazing, where there is a temperate rainy climate and pasture is the main feeding source. In these regions, over 50% of beef and over 70% of milk are produced nationally (INE, 2007). The dairy industry is an important source of animals for beef production because only 20% of the

¹Instituto de Investigaciones Agropecuarias INIA, Ruta 5 Norte km 8, P.O. Box 24-0, Osorno, Chile.

*Corresponding author (rmorales@inia.cl).

²Institut de Recerca i Tecnologia Agroalimentaries (IRTA), Finca Camps i Armet s/n, E-17121, Monells (Girona), Spain.

Received: 20 June 2011.

Accepted: 4 January 2012.

animals produced in Chile are from beef cattle breeds, whereas the rest come from dairy systems (Goic, 2001). There is another type of semi-intensive production system in Chile that combines pasture grazing with supplement.

The beef production system based on pasture has a lower environmental impact and causes lower animal stress than other systems (IICA, 2004). Additionally, consumers positively value beef from animals fed on pasture and raised outdoors (Schnettler *et al.*, 2008). The beef industry plays an important role in the economy of the southern regions of Chile where more than one million hectares of pastures are available for animal production and the country relies on an exceptional sanitary status. In recent years, the beef industry has improved production processes to ensure quality and certification for overseas meat markets. However, there is limited scientific information about the nutritional quality of beef from Chile. In addition, limited research has been focused on the influence of diet from different Chilean production systems on beef fatty acid profile and cholesterol content. The aim of this study was to evaluate the nutritional quality (intramuscular fat and cholesterol content and fatty acid composition) of beef produced in Chile under different production systems classified according to the type of finishing diet.

MATERIALS AND METHODS

Animals and diets

Two-hundred and five steers from 13 livestock farms were evaluated. Animals were classified into three production systems according to the type of finishing diet (the last 30 d before slaughter). The predominant animal breed was Friesian and Friesian-crosses from dairy farms whereas the rest were British breeds. The individual animal breed information was not available at the abattoir and some farmers had different breeds in the same slaughter group.

Eighty steers were finished on pasture (P), which consisted of typical native and improved pastures from Los Ríos and Los Lagos Regions in Southern Chile (39°50' to 41°30' S). The predominant species in these grasslands

were *Lolium perenne* L., *Trifolium repens* L., *Bromus* spp. and *Holcus lanatus* L. (Siebald, 2005). Seventy nine steers were finished on pastures and supplemented with grain (wheat or corn) or concentrate (PG). Forty six steers were finished on a typical feedlot system (F) in the central zone of Chile (32° to 38° S). The number of farms and animals, and feed components of the finishing diets from the three production systems are shown in Table 1, and the chemical and fatty acid composition of the finishing diets in Table 2.

In the diet from Chilean feedlots, a low grain percentage is normally used between 20 and 40% DM (in the present study the grain level was approximately 24%, Table 2), in contrast to that of other countries such as USA (Lawrence *et al.*, 2007). In addition, different sources of fiber are used, such as alfalfa silage and hay, and many different by-products from the fruit and vegetable agro-industries (Claro and González, 2005). The type of by-product used in the feedlot depends on the price, season, and localization of the source. In the present study a typical Chilean feedlot was selected that was representative of the beef produced in central Chile under this system.

Sampling procedures

Animals between 18 and 24-mo of age and with hot carcass weight ranging from 250 to 300 kg were slaughtered. At 24 h post-mortem, 200 g of *Longissimus thoracis* were removed from the 8th to the 9th vertebra and transported to the Meat Science Laboratory of the Remehue Regional Research Centre at the Instituto de Investigaciones Agropecuarias (INIA) in Osorno, Chile. Muscle samples were vacuum packed and stored at -18 ± 2 °C until analysis.

Chemical analysis was performed on 100 g of the feedstuff and meat samples. Samples of feedstuff were collected days previous to slaughter. Grass samples were taken from pasture on the animals were grazing whereas grain and concentrate samples were obtained from storage silo. In order to quantify intramuscular fat (IMF) from meat samples, all external fat was removed.

Table 1. Number of farms and animals, and feed components of the finishing diets from the three production systems.

Production system	Farms (n)	Animals (n)	Feed type
Pasture (P)	6	80	Pasture (predominantly <i>Lolium perenne</i> L., <i>Trifolium repens</i> L., <i>Bromus</i> spp., and <i>Holcus lanatus</i> L.)
Pasture plus supplement (PG)	1	10	Pasture + 3.0 kg corn
	1	8	Pasture + 1.5 kg wheat
	2	35	Pasture + 1.5 kg concentrate
	2	26	Pasture + 3.5 kg concentrate
Feedlot (F)	1	46	Ration composition (kg): 1 crushed oat 0.5 corn 3 sifted oat 6 chicken feces 5 chopped hay 3 broad bean pod

Table 2. Average chemical composition of the finishing diets (last 30 d).

	Pasture plus supplement (PG)				Feedlot (F)
	Pasture (P) (n = 39)	Corn grain (n = 3)	Wheat grain (n = 3)	Concentrate (n = 15)	(n = 3)
Dry mater, %	15.9	71.5	88.2	88.8	74.8
Crude protein, %	22.6	5.3	16.1	17.6	18.1
Metabolizable energy, Mcal kg ⁻¹	2.8	3.5	2.9	2.6	2.2
Ash, %	9.5	1.2	1.7	6.4	15.9
Ether extract, %	2.7	2.5	1.7	4.3	2.3
C18:1 n-9, %	2.14	25.27	25.31	18.54	29.28
C18:2 n-6, %	12.01	58.00	57.30	57.21	31.70
C18:3 n-3, %	66.18	1.20	1.15	3.40	7.46
SFA, %	19.31	14.51	15.53	18.4	25.86
MUFA, %	2.29	26.15	26.02	20.56	30.46
PUFA, %	78.43	59.34	58.45	61.01	43.68
P:S	4.22	4.09	2.86	3.42	1.72
n-6:n-3	0.19	48.40	49.82	17.65	4.26

SFA: Saturated fatty acids (C14:0 + C16:0 + C18:0); MUFA: monounsaturated fatty acids (C18:1 c-9); PUFA: polyunsaturated fatty acids (C18:2 n-6; C18:3 n-3); P:S: Polyunsaturated: saturated fatty acid ratio; n-6:n-3 fatty acid ratio.

Samples were dried at $60 \pm 2^\circ\text{C}$ by 48 h and ground before analysis. Sample moisture was determined by drying at $105 \pm 2^\circ\text{C}$ until constant weight (Harris, 1970) and total fat and IMF was measured by Soxhlet extraction 920.39 method (AOAC, 1990). Total N was measured by the Kjeldahl 984.13 method (AOAC, 1990) and ash following the 942.05 method (AOAC, 1990).

Fatty acid composition

Fat extraction was performed according to that by Bligh and Dyer (1959) and Lumley and Colwell (1991). Previous to analysis 20 g of fresh sample were thawed and grounded and posteriorly extracted using methanol, chloroform and water (80:50:32 mL). After, the samples were homogenized for 30 min. Afterwards, the samples were filtrated through filter paper in a glass funnel and was added water until to observe a biphasic separation. The fat was concentrated in the chloroform layer. The chloroform phase was collected and removed by evaporation and 2.0 mL of n-hexane were added to the extract, which was then stored at $-18 \pm 2^\circ\text{C}$ until analysis. Approximately 0.5 g fat was obtained with this extraction. Before the fatty acid C23:0 (Nu-Chek Prep, Elysian, Minnesota, USA) was added to the samples as an internal standard. Transmethylation was carried out according to the method described by Ichihara *et al.* (1996). One hundred μL of KOH in 2 N methanol were added and the mixture (sample + 2.0 mL) was agitated for 3 min at room temperature. After phase separation, the supernatant was collected and analyzed by gas chromatography. The fatty acid profile was determined in a gas chromatograph (Clarus 600 Perkin Elmer®, Waltham, Massachusetts, USA) equipped with a flame ionization detector (FID). A capillary column SP-2560™ (Sigma-Aldrich Co., Bellefonte, Pennsylvania, USA) of $100\text{ m} \times 0.25\text{ mm} \times 0.25\text{ }\mu\text{m}$ film was used. Helium was used as the carrier gas at 1.0 mL min^{-1} and inlet pressure of 15 psi, and the method of injection was split (100:1). The injector temperature was fixed at 250°C and the detector temperature at 260°C . The injected sample volume was $1.0\text{ }\mu\text{L}$ and the oven temperature was programmed to increase from 140°C (held for 5 min) to 240°C (held for 15 min) at 4°C min^{-1} . Fatty acids were identified by comparing the retention times of the chromatograph peaks with those of the methyl esters from a mixture prepared with a FAME Mix standard of 37 components (Standard: 47885-U, SigmaAldrich Co, St. Louis, Missouri, USA), C18:1 *t*-11 methyl ester standard (Standard: 46905-U, SigmaAldrich) and octadecadienoic conjugated methyl acid (Standard: O5632, Sigma Aldrich).

Cholesterol content

Cholesterol extraction was performed in 2 g of sample by direct saponification of 2% KOH in 50 mL of absolute ethanol (Mazalli *et al.*, 2003) with subsequent chromatographic analysis using the following conditions:

the injector temperature was fixed at 270°C and the detector temperature at 300°C . The sample volume injected was $1.0\text{ }\mu\text{L}$. The oven temperature was programmed to increase from 160°C (held for 1 min) to 300°C (held for 7 min) at $10^\circ\text{C min}^{-1}$. A capillary column Elite-1 (Perkin Elmer®, USA) of $30\text{ m} \times 0.25\text{ mm} \times 0.25\text{ }\mu\text{m}$ film was used. Cholesterol was quantified by the standard internal method using 5- α -cholestane (Supelco Analytical, Bellefonte, Pennsylvania, USA) as reference.

Statistical analysis

Data were analyzed by ANOVA using the General Lineal Model (GLM) procedure of SAS (SAS Institute, 2001) with the type of finishing diet (pasture, pasture plus supplement, and feedlot) as a fixed effect in the model. Least-square means were separated by Tukey's Studentized range test.

RESULTS AND DISCUSSION

Carcass weight, intramuscular fat percentage and cholesterol content of *Longissimus thoracis*

Hot carcass weight was higher ($P < 0.05$) for steers finished on the feedlot diet than those from animals finished on pasture or pasture plus supplement (Table 3). Animals from P and PG production systems in Chile are usually slaughtered, between 470 and 550 kg live weight depending on the breed, whereas steers from the Chilean feedlots are slaughtered at above 550 kg live weight (Goic and Rojas, 2004; Claro and González, 2005).

No differences ($P > 0.05$) were found among groups for IMF percentage despite differences found in carcass weight between the F and the other two P and PG groups. Intramuscular fat values from this study (2.3%, 2.1% and 2.3% for P, PG, and F, respectively) were similar to those reported by Contreras (2006) for Chilean steers (1.9%).

De la Fuente *et al.* (2009) found similar IMF values to those from the present study in beef from European and Uruguayan animals finished on pasture (1.76% to 2.36%), but higher values in beef from animals fed on pasture with supplement (2.92% to 2.95%). Other authors have reported higher IMF content of Argentinean (Garcia *et al.*, 2008; Schor *et al.*, 2008) and Brazilian (Padre *et al.*, 2006)

Table 3. Least-squares means for carcass weight, intramuscular fat and cholesterol content of *Longissimus thoracis* from steers finished on pasture, pasture plus supplement and feedlot diet.

	Finishing diet			RMSE
	Pasture (P) (n = 80)	Pasture plus supplement (PG) (n = 79)	Feedlot (F) (n = 46)	
Hot carcass weight, kg	260.57 $\pm$ 30.53b	269.07 $\pm$ 58.57b	292.39 $\pm$ 35.65a	43.304
Intramuscular fat, %	2.33 $\pm$ 1.06	2.09 $\pm$ 0.79	2.29 $\pm$ 1.11	0.959
Cholesterol 100 mg g ⁻¹	73.14 $\pm$ 15.61ab	66.89 $\pm$ 18.64b	79.87 $\pm$ 15.16a	16.834

RMSE: Root mean squared error.

Common letter within a row indicates no significant difference ($P > 0.05$).

beef than those reported here. These authors reported IMF values from 2.86% to 3.38% for beef from animals fed on pasture and 3.5% to 4.2% for those from feedlot animals. Besides diet, animal breed, age and weight at slaughter can influence fat deposition among other factors (Lawrence and Fowler, 1997), and it is difficult to draw conclusions regarding the main factor determining differences in IMF deposition among studies.

Beef from animals fed on pasture-based systems showed lower cholesterol values (Table 3) compared to that from feedlot animals (PG: 66.9 vs. F: 79.9 mg 100 g⁻¹, $P < 0.05$; and P: 73.1 vs. F: 79.9 mg 100 g⁻¹, $P > 0.05$, numerical difference). Similar cholesterol values have been observed by other authors (Schor *et al.*, 2008; Orellana *et al.*, 2009), whereas lower cholesterol values (45.6 to 52.3 mg 100 g⁻¹) have been reported for beef from steers fed on pasture by Padre *et al.* (2006) and feedlot by Rule *et al.* (2002). However, some authors have indicated that the cholesterol content of muscle does not differ significantly between the feeding regimes of cattle (Wheeler *et al.*, 1987; Taylor and Smith, 1990; Rule *et al.*, 1997).

***Longissimus thoracis* intramuscular fatty acid composition**

Intramuscular fat from steers fed P and PG diets had higher ($P < 0.05$) proportions of C20:0 and C24:0 compared to beef from feedlot animals, with no differences between treatments in myristic (C14:0) fatty acid (Table 4). Intramuscular fat from feedlot animals showed higher values of all saturated fatty acid (SFA), C15:0, C16:0 (except for PG), C17:0 and C18:0 compared to beef from pasture-based systems. The sum of all monounsaturated fatty acid (MUFA), and particularly oleic acid (C18:1 *c*-9), was higher in intramuscular fat from animals fed P and PG diets compared with those fed the feedlot diet (43.3% and 42.5% vs. 39.8% MUFA, and 36.1% and 37.2% vs. 34.2% C18:1 *c*-9, respectively).

Longissimus thoracis muscle from P- and PG-fed animals showed lower percentages of C14:1, C15:1, C17:1, C20:1 (except for PG) and C24:1 and a higher percentage of C16:1 compared with F ($P < 0.05$). There were no differences ($P > 0.05$) among treatments in the percentage of all PUFA. However, beef from P and PG showed lower percentages of C18:2 *n*-6, and C20:3 *n*-3, and higher percentages of C18:3 *n*-3 and C20:3 *n*-6 (except for P which did not differ from F) than beef from the F production system ($P < 0.05$). The percentage of all PUFAs was higher than those reported in other studies using different animal feed sources (Schor *et al.*, 2008), but similar to those found by Padre *et al.* (2006) and Garcia *et al.* (2008) for Brazilian and Argentinean beef, respectively.

Intramuscular fat from animals fed on pasture only showed a higher percentage of C18:1 *t*-11 (vaccenic acid) than the two other finishing diets ($P < 0.05$). In

Table 4. Least-squares means for fatty acid composition (%) of *Longissimus thoracis* from steers finished on pasture, pasture plus supplement and feedlot diet.

Fatty acid	Finishing diet			RMSE
	Pasture (P) (n = 80)	Pasture plus supplement (PG) (n = 79)	Feedlot (F) (n = 46)	
SFA				
C14:0	2.21 ± 0.63	2.19 ± 0.59	2.30 ± 0.40	0.569
C15:0	0.49 ± 0.11b	0.42 ± 0.09c	0.56 ± 0.10a	0.103
C16:0	23.49 ± 2.22b	24.13 ± 2.36ab	24.78a ± 1.34	2.115
C17:0	2.52 ± 0.65b	2.49 ± 0.86b	2.80 ± 0.42a	0.698
C18:0	20.47 ± 2.27b	20.23 ± 2.30b	21.80 ± 2.20a	2.267
C20:0	0.08 ± 0.05a	0.06 ± 0.06a	0.02 ± 0.03b	0.050
C24:0	0.09 ± 0.07a	0.09 ± 0.07a	0.04 ± 0.04b	0.063
MUFA				
C14:1	0.31 ± 0.12b	0.33 ± 0.12b	0.42 ± 0.12a	0.119
C15:1	0.19 ± 0.06b	0.17 ± 0.05b	0.35 ± 0.08a	0.061
C16:1	0.76 ± 0.53a	0.72 ± 0.65a	0.48 ± 0.08b	0.520
C17:1	1.08 ± 0.23a	0.96 ± 0.26b	1.17 ± 0.10a	0.223
C18:1 <i>t</i> -11	4.46 ± 3.27a	2.90 ± 2.73b	2.75 ± 1.85b	2.806
C18:1 <i>c</i> -9	36.10 ± 4.09a	37.21 ± 4.23a	34.18 ± 3.77b	4.075
C20:1	0.06 ± 0.05b	0.08 ± 0.04ab	0.09 ± 0.02a	0.042
C24:1	0.32 ± 0.28b	0.16 ± 0.29b	0.34 ± 0.24a	0.029
PUFA				
C18:2 <i>n</i> -6	2.02 ± 0.98c	2.61 ± 1.82b	3.24 ± 1.05a	1.368
CLA C18:2 <i>c</i> -9 <i>t</i> -11	0.88 ± 0.25a	0.70 ± 0.29b	0.61 ± 0.16b	0.246
C18:3 <i>n</i> -3	1.03 ± 0.38a	1.02 ± 0.32a	0.71 ± 0.36b	0.354
C20:2 <i>c</i> -11 <i>c</i> -14	0.08 ± 0.13	0.07 ± 0.13	0.06 ± 0.03	0.113
C20:3 <i>n</i> -6	0.18 ± 0.09ab	0.19 ± 0.01a	0.15 ± 0.05b	0.090
C20:3 <i>n</i> -3	0.16 ± 0.12b	0.15 ± 0.13b	0.32 ± 0.15a	0.127
C20:4 <i>n</i> -6	0.04 ± 0.05	0.03 ± 0.04	0.04 ± 0.02	0.041
C20:5 <i>n</i> -3, EPA	0.09 ± 0.25	0.22 ± 0.24	0.03 ± 0.05	0.048
C22:5 <i>n</i> -3, DPA	0.64 ± 0.43	0.68 ± 0.40	0.57 ± 0.30	0.393
Unidentified	2.24 ± 0.39	2.18 ± 0.51	2.21 ± 0.37	0.432
SFA	49.34 ± 2.74b	49.61 ± 3.15b	52.30 ± 1.81a	2.731
MUFA	43.29 ± 2.34a	42.53 ± 3.40a	39.78 ± 2.21b	2.756
PUFA	5.13 ± 1.93	5.65 ± 2.36	5.71 ± 1.64	2.040
P:S	0.11 ± 0.04	0.11 ± 0.05	0.11 ± 0.03	0.044
<i>n</i> -6	3.13 ± 1.02b	3.54 ± 1.79b	4.03 ± 1.06a	1.379
<i>n</i> -3	1.92 ± 0.96ab	2.08 ± 0.87a	1.63 ± 0.72b	0.879
<i>n</i> -6: <i>n</i> -3	1.75 ± 0.36b	1.84 ± 0.89b	2.79 ± 1.08a	0.789

RMSE: Root mean squared error; CLA: conjugated linoleic acid; EPA: Eicosapentaenoic acid; DPA: Docosapentaenoic acid; SFA: saturated fatty acids: C14:0 + C16:0 + C17:0 + C18:0 + C20:0 + C24:0; MUFA: monounsaturated fatty acids: C14:1 + C15:1 + C16:1 + C17:1 + C18:1 *t*-11 + C18:1 *c*-9 + C20:1 + C24:1; PUFA: polyunsaturated fatty acids: C18:2 *n*-6 + C18:3 *n*-3 + C20:2 *c*-11 *c*-14 + C20:3 *n*-6 + C20:3 *n*-3 + C20:4 *n*-6 + C20:5 *n*-3 + C22:5 *n*-3 + CLA C18:2 *c*-9 *t*-11. P:S: polyunsaturated:saturated fatty acid ratio; *n*-6:*n*-3: fatty acid ratio.

Common letter within a row indicates no significant difference ($P > 0.05$).

this sense, other studies have also found that steers fed on grass had higher C18:1 *t*-11 than those fed on grain-based diets (Leheska *et al.*, 2008; Garcia *et al.*, 2008). Moreover, some studies suggest a linear increase in CLA *c*-9, *t*-11 synthesis as the C18:1 *t*-11 content of the diet increased in human subjects (Salminen *et al.*, 1998; Turpeinen *et al.*, 2002a), whereas, the rate of conversion of C18:1 *t*-11 to CLA *c*-9, *t*-11 was estimated to range from 5 to 12% in rodents to 19 to 30% in humans (Turpeinen *et al.*, 2002b).

CLA *c*-9, *t*-11 was higher ($P < 0.05$) for steers fed pasture only (1.32% and 0.88%) than for steers fed pasture plus supplement (1.08% and 0.70%) or a feedlot diet (0.94% and 0.61%). CLA *c*-9, *t*-11 is produced as a result of the biohydrogenation process occurring in the rumen, where unsaturated fatty acids (mainly C18:3 *n*-3 and C18:2 *n*-6) from the diet are first isomerized and partially saturated later (Christie, 1981). As was

mentioned, CLA *c*-9, *t*-11 is also synthesized by endogenous conversion of C18:1 *t*-11 (*trans*-vaccenic acid) by the enzyme Δ -9-desaturase in the adipose tissue and the mammary gland (Griinari and Bauman, 1999). Biohydrogenation efficiency decreases as the concentrate in the animal diet increases (Sauvant and Bas, 2001) and the CLA *c*-9, *t*-11 concentration in the adipose tissue is higher when animals are fed on pasture than those fed on stored forages or grain (French *et al.*, 2000). Other studies have also shown that steers finished on pasture had higher CLA *c*-9, *t*-11 than those fed on grain-based diets (French *et al.*, 2000; Realini *et al.*, 2004; Nuernberg *et al.*, 2005; Garcia *et al.*, 2008; Leheska *et al.*, 2008; De la Fuente *et al.*, 2009).

There were no differences among treatments in the P:S ratio of the intramuscular fat ($P > 0.05$). A value of 0.4 or higher is recommended for the P:S ratio (Department of Health, 1994). However, De la Fuente *et al.* (2009) indicated that the P:S ratio is of limited significance because not all saturated fatty acids increase cholesterol. Moreover, the positive effect of monounsaturated fatty acids such as C18:1 *c*-9 (Lee *et al.*, 1998) for human health is not considered when this ratio is used. C18:1 *c*-9 increases human HDL-cholesterol and decreases LDL-cholesterol concentrations (Katan *et al.*, 1994), and there is a positive relationship between LDL-cholesterol levels and human cardiovascular diseases. In contrast, HDL-cholesterol reduces the risk of cardiovascular diseases (Kwiterovich, 1997). Beef from P- and PG-fed animals showed higher values of C18:1 *c*-9 than that from F-fed animals (Table 4). Considerable attention has been paid to the relative proportion of *n*-6 and *n*-3 fatty acids, as diets with high *n*-6:*n*-3 ratios have been highlighted as risk factors in certain cancers and coronary heart diseases (Hibbeln *et al.*, 2006). A value of 4.0 or less for a diet is recommended for the *n*-6:*n*-3 ratio (Department of Health, 1994).

Beef from all production systems evaluated showed *n*-6:*n*-3 ratios lower than 4.0. However, beef from steers fed P and PG showed ratio values lower than 2.0 (Table 4), whereas beef from feedlot steers showed a mean ratio value of 2.79 ($P < 0.05$). Similar results have been reported by other authors (Enser *et al.*, 1998; Nuernberg *et al.*, 2002; Schor *et al.*, 2008; De la Fuente *et al.*, 2009) who have also found that the *n*-6:*n*-3 ratio increases as the grain or concentrate content in the diet increases.

The quantitative fatty acid composition in mg of fatty acids per 100 g of muscle is shown in Table 5. Although seafood is the major dietary source of *n*-3 fatty acids, red meat also constitutes a significant source of *n*-3 fatty acids for some populations with reduced intake of seafood products (Sinclair *et al.*, 1994). Beef from animals fed on pasture showed higher levels of *n*-3 fatty acids than that from feedlot animals. Two hundred and fifty grams of beef from pasture-fed animals provided 112.5 mg of *n*-3 fatty acids, roughly 5.6% of the dietary reference intake

Table 5. Least-squares means for fatty acid composition (mg 100 g⁻¹) of *Longissimus thoracis* from steers finished on pasture, pasture plus supplement and feedlot diet.

Selected fatty acids	Finishing diet			RMSE
	Pasture (P) (n = 80)	Pasture plus supplement (PG) (n = 79)	Feedlot (F) (n = 46)	
C18:2 <i>n</i> -6	47.4 ± 34.9b	49.3 ± 32.6b	68.6 ± 26.2a	32.23
C18:3 <i>n</i> -3	24.2 ± 16.7a	20.5 ± 9.6a	14.3 ± 5.9b	12.43
CLA	20.0 ± 10.4a	14.9 ± 9.0b	14.0 ± 7.7b	9.33
C18:2 <i>c</i> -9, <i>t</i> -11				
SFA	1148.2 ± 514.0	1020.9 ± 409.3	1202.4 ± 583.9	495.44
MUFA	1011.2 ± 469.7	886.3 ± 358.2	928.6 ± 497.8	438.59
PUFA	119.3 ± 77.1	111.3 ± 55.0	119.7 ± 36.0	61.73
<i>n</i> -6	72.7 ± 43.3ab	68.6 ± 36.3b	86.4 ± 32.2a	38.44
<i>n</i> -3	45.0 ± 34.7a	41.3 ± 22.7ab	32.2 ± 8.4b	26.34

RMSE: Root mean squared error; CLA: conjugated linoleic acid; SFA: saturated fatty acids; C14:0 + C16:0 + C17:0 + C18:0 + C20:0 + C24:0; MUFA: monounsaturated fatty acids; C14:1 + C15:1 + C16:1 + C18:1 *c*-9 + C18:1 *t*-11 + C20:1 + C24:1; PUFA: polyunsaturated fatty acids; C18:2 *n*-6 + C18:3 *n*-3 + C20:2 *c*-11 + C14 + C20:3 *n*-6 + C20:3 *n*-3 + C20:4 *n*-6 + C20:5 *n*-3 + C22:5 *n*-3 + CLA C18:2 *c*-9 *t*-11.

Common letter within a row indicates no significant difference ($P > 0.05$).

for *n*-3 fatty acids (FAO/WHO, 2008), whereas beef from feedlot animals supplied 80.5 mg of *n*-3 fatty acids, which would correspond to 4.1% of the recommended value.

Based on the amount of CLA *c*-9, *t*-11 needed to reduce cancer in experimental animal studies (Ritzenthaler *et al.*, 2001), humans would need to consume around 620 mg CLA *c*-9, *t*-11 d⁻¹ for men and 441 mg CLA *c*-9, *t*-11 d⁻¹ for women to have a cancer protective effect (Ritzenthaler *et al.*, 2001). Results from the present study show that 250 g of Chilean beef from pasture-fed animals would provide 50 mg (8.1% to 11.3%) of the CLA *c*-9, *t*-11 requirements, whereas PG and F would contribute 37.8 mg (6.1% to 8.6%) and 35 mg (5.6% to 7.9%), respectively. Thus, Chilean beef from pasture-fed animals, coupled with the consumption of grass-fed dairy products, would provide a higher concentration of CLA *c*-9, *t*-11 than beef from feedlot cattle to achieve dietary CLA *c*-9, *t*-11 levels closer to the recommended values.

There is limited nutritional information available characterizing the nutritional quality of Chilean beef. Results from the present study provide quantitative information for human nutritionists to provide more accurate consumer recommendations with values from local beef, rather than using data from international nutritional tables. In addition, objective information on the nutritional quality of Chilean beef from different production systems will be valuable for promoting beef consumption in local and international trade.

CONCLUSIONS

Beef from pasture-fed animals showed higher contents of C18:1 *t*-11, CLA *c*-9, *t*-11 and *n*-3 fatty acids, and a lower *n*-6:*n*-3 ratio than beef from feedlot animals. Beef from animals fed on pasture supplemented with grain showed a similar composition to that from pasture-fed animals, and intermediate values for some fatty acids between pasture and feedlot diets. However, Chilean beef from

the three production systems evaluated showed similar levels of intramuscular fat and a *n-6:n-3* ratio lower than 4.0. Results indicate that the *Longissimus* muscle from Chilean beef could be recommended as a source of red meat for a healthy diet.

ACKNOWLEDGEMENTS

This work was funded by the “Inter-American Development Bank (IDB)” through a “Bien Club” negotiated by the “Agencia Regional de Desarrollo Productivo de la Región de Los Lagos” and supported by “Cluster de carne bovina” and “La Corporación de la Carne”. We would like to acknowledge the CONICYT (PDA08) for financing Rodrigo Morales’s post-doctorate studies with a “Programa Bicentenario de Ciencia y Tecnología (PBCT)” scholarship. We thank Nelly Delgado and Paulina Flores for sample analyses. We are grateful to the slaughterhouse staff for their cooperation in the present study: Jaime Fuchslocher, Rodrigo Soto, Rodrigo Vergara, and Astrid Hernández at FRIGOSORNO S.A.; Italo Mencarini, Héctor Mimica, Javier Ibáñez, and Consuelo Rice at FRIVAL; and Javier Navarro at FRIMA S.A.; José Godoy, Juan Carlos Letelier, and Deborah Alarcón at AASA.

Calidad nutricional de carne obtenida de diferentes sistemas productivos.

En los últimos años, el sector de carne bovina en Chile ha desarrollado un notable avance a nivel de mejoramiento de procesos de producción, aseguramiento de calidad, y certificación exportadora. Sin embargo, existe poca información disponible sobre la calidad nutricional de la carne producida en Chile. En el presente estudio se evaluó la calidad nutricional de la carne producida en Chile en diferentes sistemas de producción clasificados de acuerdo con el tipo de acabado. Se evaluó la carne de 205 animales de 13 predios: 80 novillos se finalizaron a pastoreo, 79 fueron finalizados a pastoreo más suplementación, mientras que 46 se terminaron en un típico feedlot de la zona central de Chile. En el músculo *Longissimus thoracis* se determinó la cantidad de grasa intramuscular, el perfil de ácidos grasos, y el colesterol. La carne de los animales finalizados a pradera muestra altos niveles de ácido vaccénico (C18:1 *trans*-11), ácido ruménico (CLA *cis*-9 *trans*-11) y ácidos grasos *n-3*, además de un bajo nivel de la relación *n-6:n-3* en comparación con los animales de feedlot. Sin embargo, la carne de los tres sistemas de producción muestra similares niveles de grasa intramuscular y una relación *n-6:n-3* bajo 4.0. Los resultados indican que la carne chilena podría ser recomendada como fuente alimenticia para una dieta saludable.

Palabras clave: ácidos grasos, praderas, calidad de carne chilena.

LITERATURE CITED

- AOAC. 1990. Official methods of analysis of the Association of Official Analytical Chemists. 15th ed. Vol. II. p. 1298. Association of Official Analytical Chemists (AOAC), Arlington, Virginia, USA.
- Bligh, E.G., and W.J. Dyer. 1959. A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology* 37:911-917.
- Christie, W. 1981. Lipid metabolism in ruminant animals. p. 452. Pergamon Press Ltd., Oxford, UK.
- Claro, D., y M. González. 2005. Producción de carne bovina en la zona central. p. 323-349. In A. Catrileo (ed.) Producción y manejo de carne bovina en Chile. Instituto de Investigaciones Agropecuarias INIA, Temuco, Chile.
- Contreras, S. 2006. Efecto de la suplementación con cereales y afrecho de soya en la engorda de novillos sobre las características de canal y calidad de carne. Tesis de Medicina Veterinaria. Universidad Austral de Chile, Facultad de Ciencias Veterinarias, Valdivia, Chile.
- DEIS. 2007. Indicadores básicos de salud 2007. Departamento de estadísticas e Información de Salud (DEIS), Ministerio de Salud, Gobierno de Chile, Santiago, Chile.
- De la Fuente, J., M.T. Díaz, I. Álvarez, M.A. Oliver, M. Font i Furnols, C. Sañudo, *et al.* 2009. Fatty acid and vitamin E composition of intramuscular fat in cattle reared in different production systems. *Meat Science* 82:331-337.
- Department of Health. 1994. Nutritional aspects of cardiovascular disease. Report on Health and Social Subjects N° 46. Her Majesty's Stationery Office (HMSO), The National Archives, London, UK.
- Du, M., D.U. Ahn, and J.L. Sell. 2000. Effects of dietary conjugated linoleic acid and linoleic:linolenic acid ratio on polyunsaturated fatty acid status in laying hens. *Poultry Science* 79:1749-1756.
- Elgersma, A., G. Ellen, H. van der Horst, B.G. Muuse, H. Boer, and S. Tamminga. 2003. Comparison of the fatty acid composition of fresh and ensiled perennial ryegrass (*Lolium perenne* L.), affected by cultivar and regrowth interval. *Animal Feed Science and Technology* 108:191-205.
- Enser, M., K.G. Hallett, B. Hewitt, G.A. Fursey, J.D. Wood, and G. Harrington. 1998. Fatty acid content and composition of UK beef and lamb muscle in relation to production system and implications for human nutrition. *Meat Science* 49:329-341.
- FAO/WHO. 2008. Fats and fatty acids in human nutrition report of an expert consultation. Food and Agriculture Organization of the United Nations and the World Health Organization. FAO Food Nutrition Papers 91:1-91.
- French, P., C. Stanton, F. Lawless, E.G. O'Riordan, F.J. Monahan, P.J. Caffrey, and A.P. Moloney. 2000. Fatty acid composition, including conjugated linoleic acid of intramuscular fat from steers offered grazed grass, grass silage or concentrate-based diets. *Journal of Animal Science* 78:2849-2855.
- García, P.T., N.A. Pensel, A.M. Sancho, N.J. Latimori, A.M. Kloster, M.A. Amigone, and J.J. Casal. 2008. Beef lipids in relation to animal breed and nutrition in Argentina. *Meat Science* 79:500-508.
- Goic, L. 2001. Descripción y perspectivas de la producción de carne bovina en Chile. Informe final. p. 136-148. In Catrileo (ed.) Taller internacional: Limitaciones y perspectivas del sector cárnico bovino en Chile y el Mercosur. FAO, Universidad Mayor, CECUM, Santiago, Chile.
- Goic, L., y C. Rojas. 2004. Sistemas de crianza, recría y engorda en la zona sur. p. 107-120. In A. Catrileo (ed.) Manual de producción de bovinos de carne para la VIII, IX y X regiones. Instituto de Investigaciones Agropecuarias INIA, Temuco, Chile.
- Griinari, J.M., and D.E. Bauman. 1999. Biosynthesis of conjugated linoleic acid and its incorporation into meat and milk in ruminants. p. 180-200. In Yurawecz, M.P., M.M. Mossoba, J.K.G. Kramer, M.W. Pariza, and G.J. Nelson (eds.) Advances in conjugated linoleic acid research. Vol. I. AOCS Press, Champaign, Illinois, USA.

- Grunert, K.G., L. Bredahl, and K. Brunsø. 2004. Consumer perception of meat quality and implications for product development in the meat sector—a review. *Meat Science* 66:259-272.
- Harris, L.E. 1970. Nutrition research techniques for domestic and wild animals. p. 182. Vol. I. Utah State University, Logan, Utah, USA.
- Hibbeln, J.R., L.R. Nieminen, T.L. Blasbalg, J.A. Riggs, and W.E. Lands. 2006. Healthy intakes of n-3 and n-6 fatty acids: estimations considering worldwide diversity. *American Journal Clinical Nutrition* 83:1483S-1493.
- Ichihara, K., A. Shibahara, K. Yamamoto, and T. Nakayama. 1996. An improved method for rapid analysis of the fatty acids of glycerolipids. *Lipids* 31:535-539.
- IICA. 2004. Estudio comparativo de dos sistemas de producción de leche: pastoreo y confinamiento. 62 p. Instituto Interamericano de Cooperación para la Agricultura IICA, Oficina Nicaragua, Nicaragua.
- INE. 2007. VII Censo Agropecuario y Forestal, Instituto Nacional de Estadísticas (INE), Santiago, Chile. Available at <http://www.censoagropecuario.cl/> (accessed 24 March 2011).
- Ip, C., J.A. Scimeca, and H.J. Thompson. 1994. Conjugated linoleic acid: A powerful anticarcinogen from animal fat sources. *Cancer* 74:1050-1054.
- Katan, M.B., P.L. Zock, and R.P. Mesink. 1994. Effects of fats and fatty acid on blood lipids in humans: an overview. *American Journal of Clinical Nutrition* 60(suppl. 1):1017-1022.
- Kwiterovich, P.O. 1997. The effect of dietary fat, antioxidants and pro-oxidants on blood lipids, lipoproteins and atherosclerosis. *Journal of American Dietetic Association* 97(supplement):31-41.
- Latimori, N.J., A.M. Kloster, P.T. García, F.J. Carduza, G. Grigioni, and N.A. Pensel. 2008. Diet and genotype effects on the quality index of beef produced in the Argentine Pampeana region. *Meat Science* 79:463-469.
- Lawrence, T.L.J., and V.R. Fowler. 1997. Growth of farm animals. 337 p. 2nd ed. CABI Publishing, Wallingford, UK.
- Lawrence, J., S. Shouse, W. Edwards, D. Loy, J. Lally, and R.E. Martin. 2007. Beef feedlot systems manual. 28 p. Iowa State University, Iowa Beef Centre, Ames, Iowa, USA.
- Lee, K.N., M.W. Pariza, and J.M. Ntambi. 1998. Conjugated linoleic acid decreases hepatic stearyl-CoA desaturase mRNA expression. *Biochemical Biophysical Research Communications* 248:817-821.
- Leheska, J.M., L.D. Thompson, J.C. Howe, E. Hentges, J. Boyce, J.C. Brooks, *et al.* 2008. Effects of conventional and grass-feeding systems on the nutrient composition of beef. *Journal of Animal Science* 86:3575-3585.
- Lumley, I.D., and R.K. Colwell. 1991. Extraction of fats from fatty foods and determination of fat content. p. 227-245. In Rossell, J.B., and J.L.R. Pritchard (eds.) *Analysis of oilseeds, fats and fatty foods*. Elsevier Applied Science, London, UK.
- Mazalli, M.R., T. Saldanha, e N. Bragagnolo. 2003. Determinação de colesterol em ovos: comparação entre método enzimático e um método por cromatografia líquida de alta eficiência. *Revista do Instituto Adolfo Lutz, Sao Paulo* 62:49-53.
- Nuernberg, K., D. Dannenberger, G. Nuernberg, K. Ender, J. Voigt, N.D. Scollan, *et al.* 2005. Effect of grass-based and a concentrate feeding system on meat quality characteristics and fatty acid composition of *Longissimus* muscle in different cattle breeds. *Livestock Production Science* 94:137-147.
- Nuernberg, K., G. Nuernberg, K. Ender, S. Lorenz, K. Winkler, R. Rickert, and H. Steinhart. 2002. N-3 fatty acids and conjugated linoleic acids of *Longissimus* muscle in beef cattle. *European Journal of Lipid Science and Technology* 104:463-471.
- Orellana, C., F. Pena, A. Garcia, J. Perea, J. Martos, V. Domenech, and R. Acero. 2009. Carcass characteristics, fatty acid composition, and meat quality of Criollo Argentino and Braford steers raised on forage in a semi-tropical region of Argentina. *Meat Science* 81:57-64.
- Padre, R.G., J.A. Aricetti, F.B. Moreira, I.Y. Mizubitu, I.N. Prado, J. Visentainer, *et al.* 2006. Fatty acid profile, and chemical composition of *Longissimus* muscle of bovine steers and bulls finished in pasture system. *Meat Science* 74:242-248.
- Realini, C.E., S.K. Duckett, G.W. Brito, M. Dalla Rizza, and D. De Mattos. 2004. Effect of pasture vs. concentrate feeding with or without antioxidants on carcass characteristics, fatty acid composition, and quality of Uruguayan beef. *Meat Science* 66:567-577.
- Ritzenthaler, K., M.K. McGuire, R. Falen, T.D. Shultz, N. Dasgupta, and M.A. McGuire. 2001. Estimation of conjugated linoleic acid intake by written dietary assessment methodologies underestimates actual intake evaluated by food duplicate methodology. *Journal of Nutrition* 131:1548-1554.
- Rule, D.C., K.S. Broughton, S.M. Shellito, and G. Maiorano. 2002. Comparison of muscle fatty acid profiles and cholesterol concentrations of bison, beef cattle, elk, and chicken. *Journal of Animal Science* 80:1201-1211.
- Rule, D.C., M.D. MacNeil, and R.E. Short. 1997. Influence of sire growth potential, time on feed and growing-finishing strategy on cholesterol and fatty acids of ground carcass and *Longissimus* muscle of beef steers. *Journal of Animal Science* 75:1525-1533.
- SAS Institute. 2001. Statistical analysis system release 8.01. SAS Institute Inc., Cary, North Carolina, USA.
- Salminen, M.M., M. Mutanen, M. Jauhiainen, and A. Aro. 1998. Dietary trans-fatty acids increase conjugated linoleic acid levels in human serum. *Journal Nutrition Biochemistry* 9:93-98.
- Sauvant, D., et P. Bas. 2001. La digestion des lipides chez le ruminant. *INRA Productions Animales* 14:303-310.
- Schnettler, B., R. Silva, L. Vallejos, and N. Sepúlveda. 2008. Consumer perception of animal welfare and livestock production in the Araucania Region, Chile. *Chilean Journal of Agricultural Research* 68:80-93.
- Schor, A., M.E. Cossu, A. Picallo, J. Martínez Ferrer, J.J. Grigera Naón, and D. Colombatto. 2008. Nutritional and eating quality of Argentinean beef: A review. *Meat Science* 79:408-422.
- Scollan, N., N-J. Choi, E. Kurt, A. Fisher, M. Enser, and J.D. Wood. 2001. Manipulating the fatty acid composition of muscle and adipose tissue in beef cattle. *British Journal of Nutrition* 85:115-124.
- Siebold, E. 2005. Praderas naturalizadas para la producción de carne bovina. p. 61-74. In A. Catrileo (ed.) *Producción y manejo de carne bovina en Chile*. Instituto de Investigaciones Agropecuarias INIA, Temuco, Chile.
- Sinclair, A.J., L. Johnson, K. O'Dea, and R.T. Holman. 1994. Diet rich in lean beef increase arachidonic acid and long-chain omega 3 polyunsaturated fatty acids levels in plasma phospholipids. *Lipids* 29:337-343.
- Steen, R., and M. Porter. 2003. The effects of high-concentrate diets and pasture on the concentration of conjugated linoleic acid in beef muscle and subcutaneous fat. *Grass and Forage Science* 58:50-57.
- Taylor, D.G., and L.W. Smith. 1990. The influence of breed and type of feed on the cholesterol content of the M. *longissimus dorsi* of steers. *Australian Journal of Experimental Agriculture* 30:797-799.
- Turpeinen, A.M., M. Mutanen, A.S.I. Aro, S.P.D. Basu, and J.M. Griinar. 2002a. Bioconversion of vaccenic acid to conjugated linoleic acid in humans. *American Journal of Clinical Nutrition* 76:504-10.
- Turpeinen A.M., M. Mautanen, A. Aro, I. Salminen, S. Basu, and D.L. Palmquist. 2002b. Bioconversion of vaccenic acid to conjugated linoleic acid in humans. *American Journal of Clinical Nutrition* 76:504-10.
- Wheeler, T.L., G.W. Davis, B.J. Stoecker, and C.J. Harmon. 1987. Cholesterol concentration of *longissimus* muscle, subcutaneous fat and serum of two beef cattle breed types. *Journal of Animal Science* 65:1531-1537.
- Whetsell, M., E. Raybun, and J.D. Dossier. 2003. Human health effects of fatty acids in beef. *Pasture-Based Beef Systems for Appalachia Project, Extension Service*. West Virginia University, Morgantown, Virginia, USA.
- Wood, J.D., R.I. Richardson, G.R. Nute, A.V. Fisher, M.M. Campo, E. Kasapidou, *et al.* 2003. Effects of fatty acids on meat quality: A review. *Meat Science* 66:21-32.