

COVID-19 vaccine hesitancy - A challenge to vaccine uptake among youth in Saudi Arabia

Authors: B. S. Alasmri^{1*}; S. E. Mahmood²

Affiliations: ¹Saudi Board Preventive Medicine, Abha, Saudi Arabia; ²Department of Family and Community Medicine, King Khalid University, Abha, Saudi Arabia

EDITORIAL

The coronavirus disease (COVID-19 emerged in Wuhan, China, in late 2019 and by February 12th, 2020, it had spread across all countries, becoming a Public Health Emergency of International Concern (PHEIC) [1]. The World Health Organization (WHO) declared it a global pandemic on March 11th, 2020 [2].

The causative virus is a novel type of coronavirus known as Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) and the disease was named COVID-19 [3], severely affecting elderly people and people with comorbidities the most [4,5].

Preventive measures (e.g., physical social distancing, lockdown, and mask-wearing) were implemented by all countries, including Kingdom of Saudi Arabia (KSA), to control the spread of the disease [2,6,7]. Although these measures limit COVID-19 transmission, vaccination is one of the mainstays to prevent the spread of and eradicate the SARS-CoV-2 infection and the COVID-19 pandemic [8].

Extensive research has been conducted to develop COVID-19 vaccines rapidly and by late 2020, the first vaccines were ready to be rolled out, starting with the most vulnerable groups [6,9,10]. The Saudi Ministry of Health launched a vaccine

campaign on December 17th, 2020, to provide free vaccination to all citizens and residents through about 600 vaccination centers around the country [11]. However, vaccination hesitancy, the reluctance or refusal to vaccinate, despite the availability of vaccines, is one of the barriers to COVID-19 vaccination acceptance [12].

Even if COVID-19 vaccine success depends on high uptake levels, vaccine hesitancy poses significant risks for the hesitant individual and the community [13]. Vaccine hesitancy has been observed even before the COVID-19 pandemic, regardless of education level, socioeconomic status, and nationality. It has also been reported for healthcare workers. This is because vaccine acceptance depends on an interest in personal protection against the disease balanced with concerns about safety [12].

Studies showed the safety and efficacy of the vaccine as the top reasons for hesitancy together with religious reasons [14]. Despite many reports highlighting that the SARS-CoV-2 vaccine is safe, for some people, the rapid development of the vaccine raises safety concerns and it is not regarded as a safe scientific breakthrough. This is a unique characteristic attitude compared to hesitancy toward other vaccines [13]. Apart from that, new mutations of the virus, the need for continued engagement in preventive measures after people

***Corresponding author:** Bushra Saeed Alasmri, Saudi Board Preventive Medicine, Abha, Saudi Arabia, email: Boshraasmri99@hotmail.com; **Potential Conflicts of Interest (Col):** All authors: no potential conflicts of interest disclosed; **Funding:** All authors: All authors: no funding has been sought or gained for this project; **Academic Integrity.** All authors confirm that they have made substantial academic contributions to this manuscript as defined by the ICMJE; **Ethics of human subject participation:** The study was approved by the local Institutional Review Board. Informed consent was sought and gained where applicable; **Originality:** All authors: this manuscript is original has not been published elsewhere; **Review:** This manuscript was peer-reviewed by three reviewers in a double-blind review process; **Type-editor:** Ahmed (USA).

Received: 25th February 2022; **Initial decision given:** 25th February 2022; **Revised manuscript received:** 26th February 2022; **Accepted:** 26th February 2022.

Copyright: © The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY-NC-ND) ([click here](#)) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. **Publisher:** Rwanda Biomedical Centre (RBC)/Rwanda Health Communication Center, P. O. Box 4586, Kigali. ISSN: 2079-097X (print); 2410-8626 (online)

Citation for this article: B. S. Alasmri; S. E. Mahmood. COVID-19 vaccine hesitancy- A challenge to vaccine uptake among youth in Saudi Arabia. Rwanda Medical Journal, Vol. 79, no. 1, p. 57-59, 2022. <https://dx.doi.org/10.4314/rmj.v79i1.8>

have been vaccinated and the diversity in response policy changes worldwide contribute to people's reluctance to be vaccinated [15]. Continuously changing new information about new mutations, the symptoms, severity, and mortality coupled with misinformation, rumors and false conspiracy theories circulated in the media resulted in confusion and uncertainty about the effectiveness of the developed vaccines [16].

Before the availability of COVID-19 vaccines, studies that assessed attitudes of the general public towards vaccines revealed that higher-income regions were the least certain regarding vaccine safety and that most people in lower-income areas agreed that vaccines are safe. The studies showed similar findings regarding vaccine effectiveness [9], but when more information was available about the vaccine development process, a lot of misinformation was circulated, especially on social media and resulted in lowering the public trust in the vaccine safety and effectiveness. The Middle East was among the regions with the lowest COVID-19 vaccine acceptance rates globally. Such low rates were most probably related to the region's widespread beliefs in conspiracy theories [15]. Misinformation and conspiracy theories widely circulated included that vaccines are used to track personal data and impact fertility. The misinformation about severe adverse effects like anaphylaxis and blood clots have also increased skepticism further lowered public trust [17].

In the KSA, a study showed that only 67% of respondents intended to receive the vaccine, and 7% were hesitant and most hesitant respondents (64.3%) were young adults (aged 18–29 years). A

study in South Africa also found younger adults to be more concerned and less accepting of COVID-19 vaccination than older people [16]. Another study by Fadhel et al. found the prevalence of vaccine hesitancy to be 20.6% in Saudi Arabia with dominance among females and young people [17], with 70% of the participants concerned about the safety and efficacy. Hesitancy and mistrust in COVID-19 vaccines could undermine the KSA efforts in combatting the pandemic because other studies have reported similar concerns about the safety and effectiveness of the vaccines in Saudi Arabia and the findings of the age and gender groups concerned remained consistent [8,18–20].

Therefore, education programs to raise awareness should be tailored depending on the level of health, scientific, general literacy and socio-demographics of the targeted audiences focusing on young Saudis and strategies to tackle vaccination reluctance should be designed to eradicate community-specific anti-vaccination misconceptions [21,22,23]. The involvement of trusted non-government organizations and community-based groups is crucial to building trust in COVID-19 vaccination to facilitate communication with people at community levels and help them make fully informed decisions about the vaccine. The evidence-based and behaviorally informed strategies to implement the vaccination program is also necessary to achieve high vaccine acceptance [15].

Trust-building measures based on dialogue and transparency, educating the young population about the social benefits of vaccination and engaging youth leaders have been shown to be effective.

REFERENCES

- [1] World Health Organization, "COVID 19 Public Health Emergency of International Concern (PHEIC) Global Research and Innovation Forum: Towards a Research Roadmap," *Glob. Res. Collab. Infect. Dis. Prep.*, pp. 1–10, 2020, [Online]. Available: [https://www.who.int/publications/m/item/covid-19-public-health-emergency-of-international-concern-\(pheic\)-global-research-and-innovation-forum](https://www.who.int/publications/m/item/covid-19-public-health-emergency-of-international-concern-(pheic)-global-research-and-innovation-forum).
- [2] C. Musanabagnwa, L. Munir, J. B. Mazarati, C. M. Muvunyi, S. Nsanzimana, and L. Mutesa, "Lockdown Restrictions during COVID-19 Outbreak in Rwanda," *Rw. Public Heal. Bul*, vol. 2, no. 2, pp. 24–29, 2020.
- [3] Y. Wu et al., "SARS-CoV-2 is an

appropriate name for the new coronavirus," *Lancet*, vol. 395, no. 10228, pp. 949–950, 2020, doi: 10.1016/S0140-6736(20)30557-2.

- [4] V. Jain and J. M. Yuan, "Predictive symptoms and comorbidities for severe COVID-19 and intensive care unit admission: a systematic review and meta-analysis," *Int. J. Public Health*, vol. 65, no. 5, pp. 533–546, 2020, doi: 10.1007/s00038-020-01390-7.
- [5] C. Smith, "The structural vulnerability of healthcare workers during COVID-19: Observations on the social context of risk and the equitable distribution of resources," *Soc. Sci. Med. J.*, no. January, 2020.
- [6] C. Nsanzabaganwa, "Global Trend of COVID-19 Treatment. 2(2): 13-24.," *Rw. Public Heal. Bul*. 2020, pp. 13–17, 2020.

- [7] A. K. B. Abdulrahman, K. A. B. Abdulrahman, and R. M. Nouh, "Response of Saudi population to strict preventive measures against COVID-19," *Int. J. Environ. Res. Public Health*, vol. 18, no. 24, 2021, doi: 10.3390/ijerph182413424.
- [8] M. Al-Mohaithef and B. K. Padhi, "Determinants of COVID-19 vaccine acceptance in Saudi Arabia: A web-based national survey," *J. Multidiscip. Healthc.*, vol. 13, pp. 1657–1663, 2020, doi: 10.2147/JMDH.S276771.
- [9] X. Tang et al., "COVID-19 vaccination intention during early vaccine rollout in Canada: a nationwide online survey," *Lancet Reg. Heal. - Am.*, no. January, 2020.
- [10] B. Noella, "Rwanda COVID-19 Vaccination Success Story," *Public Heal. Bul.*, pp. 10–12, 2021.
- [11] Y. S. Alqahtani, "Acceptability of the COVID-19 Vaccine among Adults in Saudi Arabia: A Cross-Sectional Study of the General Population in the Southern Region of Saudi Arabia," *Vaccines*, vol. 10, no. 1, 2022, doi: 10.3390/vaccines10010041.
- [12] K. Matsui, "Vaccination Hesitancy for COVID-19," *JMA J.*, vol. 4, no. 4, pp. 443–444, 2021, doi: 10.31662/jmaj.2021-0157.
- [13] K. G. M. Danabal, S. S. Magesh, S. Saravanan, and V. Gopichandran, "Attitude towards COVID 19 vaccines and vaccine hesitancy in urban and rural communities in Tamil Nadu, India – a community based survey," *BMC Health Serv. Res.*, vol. 21, no. 1, pp. 1–10, 2021, doi: 10.1186/s12913-021-07037-4.
- [14] S. Lawes-Wickwar et al., "A rapid systematic review of public responses to health messages encouraging vaccination against infectious diseases in a pandemic or epidemic," *Vaccines*, vol. 9, no. 2, pp. 1–26, 2021, doi: 10.3390/vaccines9020072.
- [15] S. A. Nossier, "Vaccine hesitancy: the greatest threat to COVID-19 vaccination programs," *J. Egypt. Public Health Assoc.*, vol. 96, no. 1, pp. 18–20, 2021, doi: 10.1186/s42506-021-00081-2.
- [16] S. Cooper, H. van Rooyen, and C. S. Wiysonge, "COVID-19 vaccine hesitancy in South Africa: A complex social phenomenon," *South African Med. J.*, vol. 111, no. 8, pp. 702–703, 2021, doi: 10.7196/SAMJ.2021.v111i8.15800.
- [17] F. H. Fadhel, "Vaccine hesitancy and acceptance: an examination of predictive factors in COVID-19 vaccination in Saudi Arabia," *Health Promot. Int.*, pp. 1–13, 2021, doi: 10.1093/heapro/daab209.
- [18] D. Almaghaslah, A. Alsayari, G. Kandasamy, and R. Vasudevan, "COVID-19 vaccine hesitancy among young adults in Saudi Arabia: A cross-sectional web-based study," *Vaccines*, vol. 9, no. 4, pp. 1–8, 2021, doi: 10.3390/vaccines9040330.
- [19] M. Mahmoud, M. M. Alanazi, M. S. Albarrak, N. K. Aljarba, and N. G. Almutairi, "The Percentage of Vaccine Hesitancy among Married Individuals in Times of the COVID-19 Pandemic: A Cross Sectional Study in Riyadh City, Kingdom of Saudi Arabia," *Saudi J. Heal. Syst. Res.*, pp. 1–7, 2021, doi: 10.1159/000520681.
- [20] S. S. Othman et al., "Association between social media use and the acceptance of COVID-19 vaccination among the general population in Saudi Arabia - a cross-sectional study," *BMC Public Health*, vol. 22, no. 1, p. 375, 2022, doi: 10.1186/s12889-022-12757-1.
- [21] J. V. Lazarus et al., "A global survey of potential acceptance of a COVID-19 vaccine," *Nat. Med.*, vol. 27, no. 2, pp. 225–228, 2021, doi: 10.1038/s41591-020-1124-9.
- [22] C. Lin, P. Tu, and L. M. Beitsch, "Confidence and receptivity for COVID-19 vaccines: A rapid systematic review," *Vaccines*, vol. 9, no. 1, pp. 1–32, 2021, doi: 10.3390/vaccines9010016.
- [23] E. A. Largent, G. Persad, S. Sangenito, A. Glickman, C. Boyle, and E. J. Emanuel, "US Public Attitudes Toward COVID-19 Vaccine Mandates," *JAMA Netw. Open*, vol. 3, no. 12, pp. 2020–2023, 2020, doi: 10.1001/jamanetworkopen.2020.33324.